

Diffusions with a nonlinear irregular drift coefficient and probabilistic interpretation of generalized Burgers' equations

B.Jourdain*

December 11, 1996

Abstract

We prove existence and uniqueness for two classes of diffusions which are the sum of a Brownian motion and a nonlinear irregular drift. Within each class, we construct diffusions linked with generalized Burgers' equations.

Introduction

In this paper, we are interested in diffusions given by two nonlinear martingale problems. Each problem is closely linked to the nonlinear partial differential equation satisfied by the time marginals of any solution. Under our assumptions on the diffusion and the drift coefficients, the time marginals are absolutely continuous (for $t > 0$) and the partial differential equation provides a nice evolution equation for the densities. Our proofs for existence and uniqueness are based on fixed-point methods for this evolution equation.

The first section is devoted to a mean field martingale problem. For F a bounded measurable \mathbb{R}^d valued function on $[0, +\infty) \times \mathbb{R}^d \times \mathcal{P}(\mathbb{R}^d)$, Lipschitz continuous in its last variable for the total variation metric, we say that $P \in \mathcal{P}(C([0, +\infty), \mathbb{R}^d))$ with time marginals $(P_t)_{t \geq 0}$ solves the nonlinear martingale problem (MP1) starting at $m \in \mathcal{P}(\mathbb{R}^d)$ if $P_0 = m$ and for any $\phi \in C_b^2(\mathbb{R}^d)$

$$\phi(X_t) - \phi(X_0) - \int_0^t \left(\frac{1}{2} \Delta \phi(X_s) + F(s, X_s, P_s) \cdot \nabla \phi(X_s) \right) ds \text{ is a } P\text{-martingale}$$

where X denotes the canonical process on $C([0, +\infty), \mathbb{R}^d)$. We prove existence and uniqueness for (MP1).

If the drift coefficient F was Lipschitz continuous in its second and last variables for the sum of the Fortet-Mourier metric on $\mathcal{P}(\mathbb{R}^d)$ ($\rho(\mu, \mu') = \sup\{\int \phi d\mu - \int \phi d\mu'; |\phi(x) - \phi(y)| \leq |x - y| \wedge 1\}$)

*ENPC-CERMICS, La Courtine, 93167 Noisy le Grand Cedex France and Laboratoire de Probabilités, Université Paris 6, 4 place Jussieu, F-75252 Paris Cedex 05 France - e-mail : jourdain@cermics.enpc.fr

and the Euclidian metric on \mathbb{R}^d , we could apply classical existence and uniqueness results for nonlinear diffusions, which are proved by sample-path couplings (see for example Graham [4]). But our assumptions are much weaker since we do not suppose any continuity in the second variable and the Fortet-Mourier metric is obviously smaller than the total variation metric. The counterpart is that the diffusion coefficient is linear and the drift coefficient F is bounded. By a fixed-point method, we prove that the evolution equation satisfied by the densities of the time marginals of any solution of (MP1) admits a unique solution. The results for the martingale problem itself follow quite immediately.

By our theorem, for $d = 1$ and $F(s, x, \mu) = (\int_{\mathbb{R}} H(x - y)\mu(dy))^q$ where $q \geq 1$ and H denotes the Heaviside function ($H(x) = 1_{\{x \geq 0\}}$), the martingale problem (MP1) starting at m admits a unique solution P . Let $V(t, x)$ and $v(x)$ be the distribution functions of P_t and m . Generalizing results given by Bossy and Talay in [1] for Burgers' equation ($q = 1$), we prove that V is a weak solution of

$$\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2} - \frac{1}{q+1} \frac{\partial(u^{q+1})}{\partial x}$$

with initial condition v and obtain P as the propagation of chaos limit of a sequence of weakly interacting particle systems. Our propagation of chaos result is trajectorial and stronger than the one proved by Bossy and Talay.

The second section deals with a moderate martingale problem in which the drift coefficient depends on the densities of the time marginals. Thus the nonlinearity is more ticklish. For F a bounded measurable \mathbb{R}^d valued function on $[0, +\infty) \times \mathbb{R}^d \times \mathbb{R}$, satisfying

$$\forall s \in [0, +\infty), \forall x \in \mathbb{R}^d, \forall y, y' \in \mathbb{R}, |yF(s, x, y) - y'F(s, x, y')| \leq K_F |y - y'|$$

we say that $P \in \mathcal{P}(C([0, +\infty), \mathbb{R}^d))$ with time marginals $(P_t)_{t \geq 0}$ absolutely continuous with respect to Lebesgue measure for $t > 0$ solves the nonlinear martingale problem (MP2) starting at $m \in \mathcal{P}(\mathbb{R}^d)$ if $P_0 = m$ and for any $\phi \in C_b^2(\mathbb{R}^d)$

$$\phi(X_t) - \phi(X_0) - \int_0^t \left(\frac{1}{2} \Delta \phi(X_s) + F(s, X_s, p(s, X_s)) \cdot \nabla \phi(X_s) \right) ds \text{ is a } P\text{-martingale}$$

where for any $t > 0$, $p(t, \cdot)$ is a density of P_t .

We prove existence and uniqueness for (MP2). This generalizes a result given by Méléard and Roelly in [7] for $F : \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}^d$ bounded and satisfying a stronger Lipschitz continuity property

$$\forall x, x' \in \mathbb{R}^d, \forall y, y' \in \mathbb{R}, |F(x, y) - F(x', y')| + |yF(x, y) - y'F(x', y')| \leq K_F (|x - x'| + |y - y'|)$$

They obtain existence for the corresponding martingale problem (MP2) as a consequence of a propagation of chaos result for a sequence of moderately interacting particle systems. As for us, we give a direct proof again based on a fixed-point method for the evolution equation satisfied by p .

Thanks to this result, we show how it is possible to associate a probabilistic representation to some classical solutions of Burgers' equation, as it was sketched by Oelschläger in [9]. The initial conditions concerned are bounded probability densities on \mathbb{R} .

In the last section we generalize the previous existence and uniqueness results to similar martingale problems with a Lipschitz continuous, bounded and uniformly elliptic diffusion coefficient.

Notations

Let $\Omega = C([0, +\infty), \mathbb{R}^d)$ endowed with the topology of uniform convergence on compact sets and

with the corresponding Borel σ -field, $\Omega^T = C([0, T], \mathbb{R}^d)$ endowed with the topology of uniform convergence, X be the canonical process. For a Borel space E , $\mathcal{P}(E)$ is the space of probability measures on E endowed with the topology of weak convergence. We also define the metric of total variation on $\mathcal{P}(E)$

$$V(\mu, \mu') = \sup\left\{\int \phi d\mu - \int \phi d\mu'; \|\phi\|_{L^\infty(E)} \leq 1\right\}$$

If Z is a random variable with values in E let $\mathcal{L}(Z) \in \mathcal{P}(E)$ denote its law.

If $P \in \mathcal{P}(\Omega)$, $(P_t)_{t \geq 0}$ is the set of time marginals of P .

$$\tilde{\mathcal{P}}(\Omega) = \{P \in \mathcal{P}(\Omega); \forall t > 0, P_t \text{ is absolutely continuous with respect to Lebesgue measure}\}$$

If $P \in \tilde{\mathcal{P}}(\Omega)$, there is a measurable function $p(s, x)$ on $(0, +\infty) \times \mathbb{R}^d$ such that for any $s > 0$, $p(s, \cdot)$ is a density of P_s with respect to Lebesgue measure. See for example Meyer [8] pages 193-194. Such a function is called a measurable version of the densities.

For $x \in \mathbb{R}^d$, let $|x|$ be the Euclidian norm of x .

For $t > 0$, G_t denotes the heat kernel on \mathbb{R}^d : $G_t(x) = \frac{1}{(2\pi t)^{\frac{d}{2}}} \exp(-\frac{|x|^2}{2t})$.

The following estimate will be very useful :

$$\text{for any } 1 \leq i \leq d, \quad \left\| \frac{\partial G_t}{\partial x_i} \right\|_{L^1} \leq \frac{1}{\sqrt{t}} \quad (0.1)$$

Acknowledgement: I would like to thank Professor Sylvie Méléard for introducing me to this topic and helping me in this work.

1 The mean field martingale problem

1.1 Existence and uniqueness

Let F be a measurable \mathbb{R}^d valued function on $[0, +\infty) \times \mathbb{R}^d \times \mathcal{P}(\mathbb{R}^d)$ bounded by M_F which satisfies the following Lipschitz continuity property

$$\exists K_F \geq 0, \forall s \in [0, +\infty), \forall x \in \mathbb{R}^d, \forall \mu, \mu' \in \mathcal{P}(\mathbb{R}^d), |F(s, x, \mu) - F(s, x, \mu')| \leq K_F V(\mu, \mu')$$

Definition 1.1 Let $m \in \mathcal{P}(\mathbb{R}^d)$. We say that $P \in \mathcal{P}(\Omega)$ with time marginals $(P_t)_{t \geq 0}$ solves the nonlinear martingale problem (MP1) starting at m if $P_0 = m$ and for any $\phi \in C_b^2(\mathbb{R}^d)$

$$\phi(X_t) - \phi(X_0) - \int_0^t \left(\frac{1}{2} \Delta \phi(X_s) + F(s, X_s, P_s) \cdot \nabla \phi(X_s) \right) ds \text{ is a } P\text{-martingale} \quad (1.1)$$

Theorem 1.2 For any $m \in \mathcal{P}(\mathbb{R}^d)$, the nonlinear problem (MP1) starting at m admits a unique solution.

We need the following lemma to prove Theorem 1.2.

Lemma 1.3 *Let $m \in \mathcal{P}(\mathbb{R}^d)$, g be a measurable \mathbb{R}^d valued function on $[0, +\infty) \times \mathbb{R}^d$ bounded by M_g and P be the unique solution of the martingale problem: $P_0 = m$ and*

$$\forall \phi \in C_b^2(\mathbb{R}^d), \phi(X_t) - \phi(X_0) - \int_0^t \left(\frac{1}{2} \Delta \phi(X_s) + g(s, X_s) \cdot \nabla \phi(X_s) \right) ds \text{ is a } P\text{-martingale}$$

Then $P \in \tilde{\mathcal{P}}(\Omega)$. Any measurable version of the densities $p(s, x)$ satisfies the evolution equation,

$$\forall t > 0, p(t, x) = G_t * m(x) - \sum_{i=1}^d \int_0^t \frac{\partial G_{t-s}}{\partial x_i} * ((pg_i)(s, \cdot))(x) ds \text{ almost everywhere} \quad (1.2)$$

Moreover, if q is a measurable function on $(0, +\infty) \times \mathbb{R}^d$ which satisfies (1.2) and

$$\forall T > 0, \sup_{t \in (0, T]} \|q(t, \cdot)\|_{L^1} < +\infty$$

then q is a measurable version of the densities for P .

Proof of Lemma 1.3 : Existence and uniqueness for the martingale problem is a consequence of Girsanov's theorem. Let us prove that the solution P belongs to $\tilde{\mathcal{P}}(\Omega)$.

Under P , by Paul Levy's characterization, $X_t - X_0 - \int_0^t g(s, X_s) ds$ is a Brownian motion. We introduce the exponential martingale

$$Z_s = \exp \left(- \int_0^s g(r, X_r) \cdot dX_r + \frac{1}{2} \int_0^s |g(r, X_r)|^2 dr \right)$$

Let $t > 0$. We set $Q = Z_t \times P$. Then Girsanov's theorem implies that $(\beta_s = X_s - X_0)_{s \in [0, t]}$ is a Brownian motion under Q . Let f be a continuous function with compact support in \mathbb{R}^d .

$$\mathbb{E}(|f(X_t)|) = \mathbb{E}^Q \left(\frac{1}{Z_t} |f(X_t)| \right) \leq \sqrt{\mathbb{E}^Q \left(\frac{1}{Z_t^2} \right)} \sqrt{\mathbb{E}^Q (f^2(X_t))} \quad (1.3)$$

$$\mathbb{E}^Q (f^2(X_t)) = \mathbb{E}^Q (f^2(X_t - X_0 + X_0)) = \int_{\mathbb{R}^d} f^2(x) G_t * m(x) dx \leq \frac{1}{(2\pi t)^{\frac{d}{2}}} \|f\|_{L^2}^2 \quad (1.4)$$

$$\mathbb{E}^Q \left(\frac{1}{Z_t^2} \right) = \mathbb{E}^Q \left(\exp \left(\int_0^t 2g(s, X_s) \cdot d\beta_s - \frac{1}{2} \int_0^t |2g(s, X_s)|^2 ds + \int_0^t |g(s, X_s)|^2 ds \right) \right) \leq \exp(M_g^2 t) \quad (1.5)$$

With equations (1.3), (1.4) and (1.5), we conclude

$$|\mathbb{E}(f(X_t))| \leq \mathbb{E}(|f(X_t)|) \leq \frac{1}{(2\pi t)^{\frac{d}{4}}} \exp \left(\frac{M_g^2 t}{2} \right) \|f\|_{L^2} \quad (1.6)$$

Hence P_t is absolutely continuous with respect to Lebesgue measure and $P \in \tilde{\mathcal{P}}(\Omega)$.

Let $p(s, x)$ be a measurable version of the densities for P , ψ be a C^2 function with compact support in \mathbb{R}^d and $t > 0$. We set $\phi(s, x) = G_{t-s} * \psi(x)$ for $s \in [0, t)$ and $\phi(t, x) = \psi(x)$. The

function ϕ belongs to $C_b^{1,2}([0, t] \times \mathbb{R}^d)$ and satisfies

$$\forall (s, x) \in [0, t] \times \mathbb{R}^d, \quad \frac{\partial \phi}{\partial s}(s, x) + \frac{1}{2} \Delta \phi(s, x) = 0 \quad (1.7)$$

$$\forall (s, x) \in [0, t] \times \mathbb{R}^d, \quad |\nabla \phi(s, x)| = \sqrt{\sum_{i=1}^d \left(\int_{\mathbb{R}^d} \frac{\partial G_{t-s}}{\partial x_i}(x-y) \psi(y) dy \right)^2} \leq \frac{\sqrt{d} \|\psi\|_{L^\infty}}{\sqrt{t-s}} \quad (1.8)$$

Since $X_t - X_0 - \int_0^t g(s, X_s) ds$ is a P -Brownian motion, Itô's formula implies

$$\mathbb{E}(\phi(t, X_t)) = \mathbb{E} \left(\phi(0, X_0) + \int_0^t \left(\frac{\partial \phi}{\partial s}(s, X_s) + \frac{1}{2} \Delta \phi(s, X_s) + g(s, X_s) \cdot \nabla \phi(s, X_s) \right) ds \right)$$

By (1.7), we get rid of $\frac{\partial \phi}{\partial s} + \frac{1}{2} \Delta \phi$. Moreover, (1.8) allows to apply Fubini's theorem to obtain

$$\begin{aligned} \int_{\mathbb{R}^d} \psi(x) p(t, x) dx &= \int_{\mathbb{R}^d} G_t * \psi(x) m(dx) + \int_{(0,t] \times \mathbb{R}^d} \sum_{i=1}^d \int_{\mathbb{R}^d} \frac{\partial G_{t-s}}{\partial x_i}(x-y) \psi(y) dy (g_i p)(s, x) dx ds \\ &= \int_{\mathbb{R}^d} \psi(x) G_t * m(x) dx - \int_{\mathbb{R}^d} \sum_{i=1}^d \int_0^t \frac{\partial G_{t-s}}{\partial x_i} * ((g_i p)(s, \cdot))(y) ds \psi(y) dy \end{aligned}$$

Hence p satisfies (1.2).

To conclude the proof, we consider q a measurable function on $(0, +\infty) \times \mathbb{R}^d$ which satisfies (1.2) and $\forall T > 0, \sup_{t \in (0, T]} \|q(t, \cdot)\|_{L^1} < +\infty$. As p also satisfies (1.2),

$$\begin{aligned} \|p(t, \cdot) - q(t, \cdot)\|_{L^1} &\leq \sum_{i=1}^d \int_0^t \left\| \frac{\partial G_{t-s}}{\partial x_i} \right\|_{L^1} \|g_i(s, \cdot)(p(s, \cdot) - q(s, \cdot))\|_{L^1} ds \\ &\leq M_g \sqrt{d} \int_0^t \frac{\|p(s, \cdot) - q(s, \cdot)\|_{L^1}}{\sqrt{t-s}} ds \end{aligned}$$

After an iteration, we get

$$\begin{aligned} \|p(t, \cdot) - q(t, \cdot)\|_{L^1} &\leq M_g^2 d \int_0^t \frac{1}{\sqrt{t-s}} \int_0^s \frac{\|p(r, \cdot) - q(r, \cdot)\|_{L^1}}{\sqrt{s-r}} dr ds \\ &\leq M_g^2 d \int_0^t \|p(r, \cdot) - q(r, \cdot)\|_{L^1} \int_r^t \frac{1}{\sqrt{t-s} \sqrt{s-r}} ds dr \\ &\leq \pi M_g^2 d \int_0^t \|p(r, \cdot) - q(r, \cdot)\|_{L^1} dr \end{aligned}$$

Gronwall's lemma implies $\forall t > 0, \|p(t, \cdot) - q(t, \cdot)\|_{L^1} = 0$ which proves that q is a measurable version of the densities for P . \blacksquare

Proof of Theorem 1.2 : The proof is based on the following idea. If $(Q(t))_{t \geq 0}$ belongs to $C([0, +\infty), \mathcal{P}(\mathbb{R}^d))$, by Girsanov's theorem, the martingale problem in which the nonlinearity P_s in (1.1) is replaced by $Q(s)$ admits a unique solution P^Q . We consider the correspondance between $(Q(t))_{t \geq 0}$ and the time marginals $(P_t^Q)_{t \geq 0}$ of the solution. If P solves the nonlinear problem (MP1), then $(P_t)_{t \geq 0}$ is a fixed-point of this map. Conversely, if $(Q(t))_{t \geq 0}$ is a fixed-point, then P^Q solves the nonlinear problem (MP1).

Let $T > 0$. We define

$$A_{m,T} = \{Q \in C([0, T], \mathcal{P}(\mathbb{R}^d)); Q(0) = m \text{ and } \forall t \in (0, T], Q(t) \text{ is absolutely continuous} \\ \text{with respect to Lebesgue measure}\}$$

If $Q \in A_{m,T}$, let $\lambda(Q)$ denote a measurable version of the densities for Q . $A_{m,T}$ is complete for the metric $D(Q, Q') = \sup_{t \in (0, T]} V(Q(t), Q'(t)) = \sup_{t \in (0, T]} \|\lambda(Q)(t) - \lambda(Q')(t)\|_{L^1}$.

Let $t_0 \geq 0$. For $Q \in A_{m,T}$ we define $(\psi_{t_0, m}(Q)(t))_{t \in [0, T]}$ as the time marginals of the unique solution of the martingale problem :

$$\left\{ \begin{array}{l} P \in \mathcal{P}(\Omega^T) \text{ and } P_0 = m \\ \forall \phi \in C_b^2(\mathbb{R}^d), \phi(X_t) - \phi(X_0) - \int_0^t \left(\frac{1}{2} \Delta \phi(X_s) + F(t_0 + s, X_s, Q_s) \cdot \nabla \phi(X_s) \right) ds \text{ is a } P\text{-martingale} \end{array} \right.$$

Lemma 1.3 implies that for any $t \in (0, T]$, $\psi_{t_0, m}(Q)(t)$ is absolutely continuous with respect to Lebesgue measure. Hence $\psi_{t_0, m}(Q) \in A_{m,T}$. We are going to prove that if T is small enough, $\psi_{t_0, m}$ is a contraction on $A_{m,T}$. Using equation (1.2) given by Lemma 1.3, we obtain for $Q, Q' \in A_{m,T}$ and $t \in (0, T]$,

$$\begin{aligned} & \|\lambda(\psi_{t_0, m}(Q))(t) - \lambda(\psi_{t_0, m}(Q'))(t)\|_{L^1} \\ & \leq \sum_{i=1}^d \int_0^t \left\| \frac{\partial G_{t-s}}{\partial x_i} \right\|_{L^1} \|\lambda(\psi_{t_0, m}(Q))(s) F_i(t_0 + s, \cdot, Q(s)) - \lambda(\psi_{t_0, m}(Q'))(s) F_i(t_0 + s, \cdot, Q'(s))\|_{L^1} ds \\ & \leq \int_0^t \frac{1}{\sqrt{t-s}} \left(\|\lambda(\psi_{t_0, m}(Q))(s)\|_{L^1} \left\| \sum_{i=1}^d |F_i(t_0 + s, \cdot, Q(s)) - F_i(t_0 + s, \cdot, Q'(s))| \right\|_{L^\infty} \right. \\ & \quad \left. + \|\lambda(\psi_{t_0, m}(Q))(s) - \lambda(\psi_{t_0, m}(Q'))(s)\|_{L^1} \left\| \sum_{i=1}^d |F_i(t_0 + s, \cdot, Q'(s))| \right\|_{L^\infty} \right) ds \\ & \leq \sqrt{d} \int_0^t \frac{1}{\sqrt{t-s}} \left(K_F V(Q(s), Q'(s)) + M_F \|\lambda(\psi_{t_0, m}(Q))(s) - \lambda(\psi_{t_0, m}(Q'))(s)\|_{L^1} \right) ds \\ & \leq 2\sqrt{dT} (K_F D(Q, Q') + M_F D(\psi_{t_0, m}(Q), \psi_{t_0, m}(Q'))) \end{aligned}$$

Hence

$$(1 - 2\sqrt{dT} M_F) D(\psi_{t_0, m}(Q), \psi_{t_0, m}(Q')) \leq 2\sqrt{dT} K_F D(Q, Q')$$

We set $T = \frac{1}{4d(M_F + 2K_F)^2}$. Then $D(\psi_{t_0, m}(Q), \psi_{t_0, m}(Q')) \leq \frac{1}{2} D(Q, Q')$. Picard's fixed-point theorem implies that $\psi_{t_0, m}$ admits a unique fixed-point in $A_{m,T}$.

existence for the martingale problem (MP1)

Let Q^0 denote the fixed-point of $\psi_{0, m}$ in $A_{m,T}$. If Q^n is constructed, let Q^{n+1} be the fixed-point of $\psi_{(n+1)T, Q^n(T)}$ in $A_{Q^n(T), T}$.

We set $Q(t) = Q^n(t - nT)$ if $t \in [nT, (n+1)T]$. Let P be the solution of the martingale problem in which the nonlinearity in (1.1) is replaced by $Q(s)$. For any $\phi \in C_b^2(\mathbb{R}^d)$,

$$\phi(X_{nT+t}) - \phi(X_{nT}) - \int_0^t \left(\frac{1}{2} \Delta \phi(X_{nT+s}) + F(nT + s, X_{nT+s}, Q^n(s)) \cdot \nabla \phi(X_{nT+s}) \right) ds, \quad t \in [0, T]$$

is a P -martingale.

Hence, by induction, for any $n \in \mathbb{N}$, for any $t \in [0, T]$, $P_{nT+t} = Q^n(t) = Q(nT + t)$. And P solves the problem (MP1).

uniqueness for the martingale problem (MP1)

If P is a solution, Lemma 1.3 implies that for any $t > 0$, P_t is absolutely continuous with respect to Lebesgue measure. For any $n \in \mathbb{N}$, $(P_{nT+t})_{t \in [0, T]}$ is the fixed-point of $\psi_{nT, P_{nT}}$ in $A_{P_{nT}, T}$. By induction, uniqueness for the fixed-points implies uniqueness for the time marginals $(P_t)_{t \geq 0}$. Since the nonlinearity in the definition of (MP1) is limited to the dependence of the drift coefficient on the time marginals, uniqueness for this problem follows immediately. \blacksquare

1.2 Application

Theorem 1.2 implies existence and uniqueness for martingale problems associated with a class of partial differential equations which includes Burgers' equation.

We set $q \geq 1$, $m \in \mathcal{P}(\mathbb{R})$. Let $H(x) = 1_{\{x \geq 0\}}$ and $f : (x, \mu) \in \mathbb{R} \times \mathcal{P}(\mathbb{R}) \rightarrow (\int_{\mathbb{R}} H(x-y)\mu(dy))^q$. As f is the pointwise limit of the continuous functions $(x, \mu) \rightarrow (\int_{\mathbb{R}} H_n(x-y)\mu(dy))^q$ where $H_n(x) = n(x + 1/n)1_{\{-1/n \leq x \leq 0\}} + 1_{\{x > 0\}}$, this function is measurable. Moreover, since f takes its values in $[0, 1]$,

$$|f(x, \mu) - f(x, \mu')| \leq q \left| \int_{\mathbb{R}} H(x-y)\mu(dy) - \int_{\mathbb{R}} H(x-y)\mu'(dy) \right| \leq qV(\mu, \mu')$$

By Theorem 1.2, the martingale problem (MP1) corresponding to the choice $F(s, x, \mu) = f(x, \mu)$ admits a unique solution P starting at m . Let $V(t, x)$ and $v(x)$ be the distribution functions of P_t and m .

In [1], Bossy and Talay deal with the case $q = 1$. They prove that V is a weak solution of Burgers' equation

$$\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2} - \frac{1}{2} \frac{\partial (u^2)}{\partial x}$$

with initial condition v and obtain P as the propagation of chaos limit of a sequence of weakly interacting particle systems. Indeed they define $(X^{1,n}, \dots, X^{n,n})$ as the unique weak solution of the stochastic differential equation

$$X_t^{i,n} = X_0^{i,n} + B_t^{i,n} + \int_0^t \frac{1}{n} \sum_{j=1}^n H(X_s^{i,n} - X_s^{j,n}) ds, \quad 1 \leq i \leq n$$

where $(B^{1,n}, \dots, B^{n,n})$ is a \mathbb{R}^n -valued Brownian motion and $\mathcal{L}((X_0^{1,n}, \dots, X_0^{n,n})) = m^{\otimes n}$. They prove that for any $k \in \mathbb{N}^*$, $\mathcal{L}((X^{1,n}, \dots, X^{k,n}))$ converges weakly to $P^{\otimes k}$ when $n \rightarrow +\infty$.

We generalize their results to any $q \geq 1$ in Proposition 1.4. In fact, we follow the idea of Méléard and Roelly in [7] and prove a trajectorial propagation of chaos result. To obtain this result, we introduce a coupling between the particle systems and the limit processes with law P that we define on the same probability space. Let $B^i, i \in \mathbb{N}^*$ be a sequence of independent \mathbb{R} -valued Brownian motions and $X_0^i, i \in \mathbb{N}^*$ be a sequence of random variables IID with law m independent of the Brownian motions. Proposition 5.17 p.341 [6] implies that the one-dimensional stochastic differential equation

$$Y_t^i = X_0^i + B_t^i + \int_0^t (H * P_s(Y_s^i))^q ds$$

admits a unique strong solution. Moreover, considering the linear martingale problem associated with this equation, by the existence part of the proof of theorem 1.2, we obtain that the law of

the solution is P . The process Y_i is nonlinear in the following sense : the drift coefficient of the stochastic differential equation that it satisfies depends on the time marginals of its law.

Unlike in the one-dimensional case, to obtain a strong solution for a n -dimensional stochastic differential equation with $n > 1$, it is necessary to assume that the coefficients are locally Lipschitz continuous. That is why we replace H by H_n ($H_n(x) = n(x + 1/n)1_{\{-1/n \leq x \leq 0\}} + 1_{\{x > 0\}}$) and define the weakly interacting particle system as the unique strong solution of the stochastic differential equation

$$X_t^{i,n} = X_0^i + B_t^i + \int_0^t \left(\frac{1}{n} \sum_{j=1}^n H_n(X_s^{i,n} - X_s^{j,n}) \right)^q ds, \quad 1 \leq i \leq n$$

Proposition 1.4 *For any $q \geq 1$,*

(i) *The function V is a weak solution of the generalized Burgers' equation*

$$\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2} - \frac{1}{q+1} \frac{\partial(u^{q+1})}{\partial x} \quad \text{with initial condition } v.$$

(ii) *If \hat{P} denotes the image of P by the mapping $X \in \Omega \rightarrow (X, X) \in \Omega^2$ (the support of \hat{P} is enclosed in the diagonal of Ω^2), for any $k \in \mathbb{N}^*$, $\mathcal{L}(((X^{1,n}, Y^1), \dots, (X^{k,n}, Y^k)))$ converges weakly to $\hat{P}^{\otimes k}$ as $n \rightarrow +\infty$.*

To understand the trajectorial nature of the propagation of chaos result (ii) remark for instance that, unlike the classical result : $\forall k \in \mathbb{N}^*$, $\mathcal{L}((X^{1,n}, \dots, X^{k,n}))$ converges weakly to $P^{\otimes k}$, it implies : $\forall T > 0$, $\lim_{n \rightarrow +\infty} \mathbb{E}(1 \wedge \sup_{0 \leq t \leq T} |(X_t^{1,n}, \dots, X_t^{k,n}) - (Y_t^1, \dots, Y_t^k)|) = 0$.

Proof : (i) Our proof is a generalization of the one given by Bossy and Talay in [1]. Under P , by Paul Levy's characterization, $X_t - X_0 - \int_0^t V^q(s, X_s) ds$ is a Brownian motion. Let p be a measurable version of the densities for P and $\phi \in \mathcal{D}((0, +\infty) \times \mathbb{R})$. Applying Itô's formula and taking expectations, we get

$$\int_0^{+\infty} \int_{\mathbb{R}} p(t, x) \left(\frac{\partial \phi}{\partial t}(t, x) + \frac{1}{2} \frac{\partial^2 \phi}{\partial x^2}(t, x) + \frac{\partial \phi}{\partial x}(t, x) V^q(t, x) \right) dx dt = 0$$

Hence p is a solution in $\mathcal{D}'((0, +\infty) \times \mathbb{R})$ of the equation $\frac{\partial p}{\partial t} = \frac{1}{2} \frac{\partial^2 p}{\partial x^2} - \frac{\partial}{\partial x}(pV^q)$. Clearly, $\frac{\partial V}{\partial x} = p$ in $\mathcal{D}'((0, +\infty) \times \mathbb{R})$. Moreover, approximating $p(t, \cdot)$ in $L^1(\mathbb{R})$ by continuous functions, we obtain that the distribution function of the bounded measure $p(t, x) V^q(t, x) dx$ is $\frac{1}{q+1} V^{q+1}(t, x)$ which implies $pV^q = \frac{1}{q+1} \frac{\partial(V^{q+1})}{\partial x}$. Hence

$$\frac{\partial}{\partial x} \left(\frac{\partial V}{\partial t} - \frac{1}{2} \frac{\partial^2 V}{\partial x^2} + \frac{1}{q+1} \frac{\partial(V^{q+1})}{\partial x} \right) = 0$$

The spatial derivative of the distribution $\frac{\partial V}{\partial t} - \frac{1}{2} \frac{\partial^2 V}{\partial x^2} + \frac{1}{q+1} \frac{\partial(V^{q+1})}{\partial x}$ is zero. This implies that the distribution is invariant by spatial translation. If $\phi \in \mathcal{D}'((0, +\infty) \times \mathbb{R})$ and $z \rightarrow +\infty$,

$$\int_{(0, +\infty) \times \mathbb{R}} V(t, x - z) \left(\frac{\partial \phi}{\partial t}(t, x) + \frac{1}{2} \frac{\partial^2 \phi}{\partial x^2}(t, x) + \frac{V^q(t, x - z)}{q+1} \frac{\partial \phi}{\partial x}(t, x) \right) dx dt$$

goes to 0 by Lebesgue's theorem. Therefore for any $\phi \in \mathcal{D}((0, +\infty) \times \mathbb{R})$,

$$\int_{(0, +\infty) \times \mathbb{R}} V(t, x) \left(\frac{\partial \phi}{\partial t}(t, x) + \frac{1}{2} \frac{\partial^2 \phi}{\partial x^2}(t, x) + \frac{V^q(t, x)}{q+1} \frac{\partial \phi}{\partial x}(t, x) \right) dx dt = 0 \quad (1.9)$$

We conclude by proving that the initial condition is v . By density, equation (1.9) still holds if ϕ is $C^{1,2}$ with compact support in $(0, +\infty) \times \mathbb{R}$.

Let ψ be $C^{1,2}$ with compact support in $[0, +\infty) \times \mathbb{R}$. For $n \in \mathbb{N}^*$, we introduce the C^1 functions

$$g_n(s) = \begin{cases} 0 & \text{if } s \in [0, \frac{1}{2n}] \\ 12n^2(s - \frac{1}{2n})^2 - 16n^3(s - \frac{1}{2n})^3 & \text{if } s \in [\frac{1}{2n}, \frac{1}{n}] \\ 1 & \text{if } s \geq \frac{1}{n} \end{cases}$$

The function $\phi_n = g_n \psi$ is $C^{1,2}$ with compact support in $(0, +\infty) \times \mathbb{R}$. Using (1.9) for ϕ_n we get

$$\begin{aligned} & \int_{(0, +\infty) \times \mathbb{R}} \left(\frac{\partial \psi}{\partial t}(t, x) + \frac{1}{2} \frac{\partial^2 \psi}{\partial x^2}(t, x) + \frac{V^q(t, x)}{q+1} \frac{\partial \psi}{\partial x}(t, x) \right) V(t, x) dt dx \\ &= \int_{(0, \frac{1}{n}] \times \mathbb{R}^d} (1 - g_n(t)) \left(\frac{\partial \psi}{\partial t}(t, x) + \frac{1}{2} \frac{\partial^2 \psi}{\partial x^2}(t, x) + \frac{V^q(t, x)}{q+1} \frac{\partial \psi}{\partial x}(t, x) \right) V(t, x) dt dx \\ & - \int_{(0, \frac{1}{n}] \times \mathbb{R}^d} \frac{dg_n}{dt}(t) \psi(t, x) V(t, x) dt dx \end{aligned} \quad (1.10)$$

Since $P \in \mathcal{P}(\Omega)$, the map $t \rightarrow P_t$ is continuous and $\lim_{t \rightarrow 0} V(t, x) = v(x)$ for any x such that v is continuous at x . Hence by Lebesgue's theorem, $\lim_{t \rightarrow 0} \int_{\mathbb{R}} \psi(t, x) V(t, x) dx = \int_{\mathbb{R}} \psi(0, x) v(x) dx$. When $n \rightarrow +\infty$ in (1.10), we get

$$\int_{(0, +\infty) \times \mathbb{R}} \left(\frac{\partial \psi}{\partial t}(t, x) + \frac{1}{2} \frac{\partial^2 \psi}{\partial x^2}(t, x) + \frac{V^q(t, x)}{q+1} \frac{\partial \psi}{\partial x}(t, x) \right) V(t, x) dt dx = - \int_{\mathbb{R}} \psi(0, x) v(x) dx$$

Thus V is a weak solution of the generalized Burgers' equation with initial condition v .

(ii) We now prove the propagation of chaos result. In the sequel, ν and (X, Y) denote the canonical variables on $\mathcal{P}(\Omega^2)$ and Ω^2 . We set $\bar{\nu}_r = \nu \circ X_r^{-1}$.

The couples $(X^{i,n}, Y^i)$, $1 \leq i \leq n$ are exchangeable. Therefore the propagation of chaos result is equivalent to the convergence in distribution of the empirical measures $\mu_n = \frac{1}{n} \sum_{i=1}^n \delta_{(X^{i,n}, Y^i)}$ considered as $\mathcal{P}(\Omega^2)$ -valued random variables to $\delta_{\bar{\nu}}$ (see for example [10] and the references cited in it). Let π_n denote the law of μ_n .

By exchangeability of the variables $(X^{i,n}, Y^i)$, the tightness of the sequence $(\pi_n)_n$ is equivalent to the tightness of $(\mathcal{L}(X^{1,n}, Y^1))_n$ (see [10]) which is equivalent to the tightness of $(\mathcal{L}(X^{1,n}))_n$. These probability measures are tight since for any $T > 0$ their images by the canonical restriction from Ω to Ω^T are tight (the drift coefficient is bounded by 1 uniformly in t and n).

Let π_∞ denote the limit of a convergent subsequence of $(\pi_n)_n$ that we still index by n for simplicity. To prove that $\pi_\infty = \delta_{\bar{\nu}}$, we set $p \in \mathbb{N}^*$, $0 \leq s_1 \leq s_2 \leq \dots \leq s_p \leq s \leq t$, $\phi \in C_b^2(\mathbb{R}^2)$, $g \in C_b(\mathbb{R}^{2p})$, and define $G(\nu)$ to be equal to

$$\begin{aligned} & \langle \nu, \left(\phi(X_t, Y_t) - \phi(X_s, Y_s) - \int_s^t \left(\frac{1}{2} \left(\frac{\partial^2 \phi}{\partial x^2} + 2 \frac{\partial^2 \phi}{\partial x \partial y} + \frac{\partial^2 \phi}{\partial y^2} \right) (X_r, Y_r) + \frac{\partial \phi}{\partial x} (X_r, Y_r) (H * \bar{\nu}_r(X_r))^q \right. \right. \\ & \quad \left. \left. + \frac{\partial \phi}{\partial y} (X_r, Y_r) (H * P_r(Y_r))^q \right) dr \right) g(X_{s_1}, Y_{s_1}, \dots, X_{s_p}, Y_{s_p}) \rangle \end{aligned}$$

For $k \in \mathbb{N}^*$, we define $G_k(\nu)$ like G with H_k replacing H in $(H * \bar{\nu}_r(X_r))^q$ but not in $(H * P_r(Y_r))^q$. If $\nu^n \rightarrow \nu$, the weak convergence of $\bar{\nu}_r^n$ to $\bar{\nu}_r$ implies that $H_k * \bar{\nu}_r^n(x)$ converges to $H_k * \bar{\nu}_r(x)$ uniformly for $x \in \mathbb{R}$. Moreover, for any $r > 0$, P_r is absolutely continuous with respect to Lebesgue measure and $y \rightarrow H * P_r(y)$ is continuous. Hence G_k is continuous.

We are going to prove that $\mathbb{E}^{\pi_\infty}(G^2(\nu)) = 0$. By the continuity and boundedness of G_k , we have

$$\begin{aligned} \mathbb{E}^{\pi_\infty}(G^2(\nu)) &\leq 2\mathbb{E}^{\pi_\infty}((G - G_k)^2(\nu)) + 2 \lim_{n \rightarrow +\infty} \mathbb{E}(G_k^2(\mu^n)) \\ &\leq 2 \limsup_{k \rightarrow +\infty} \mathbb{E}^{\pi_\infty}((G - G_k)^2(\nu)) + 4 \limsup_{n \rightarrow +\infty} \mathbb{E}(G_n^2(\mu^n)) + 4 \limsup_{k \rightarrow +\infty} \limsup_{n \rightarrow +\infty} \mathbb{E}((G_k - G_n)^2(\mu^n)) \end{aligned} \quad (1.11)$$

Let us show that each term of the right-hand-side of (1.11) is equal to 0.

For the first term, it is a consequence of the convergence of $|H - H_k| * \bar{\nu}_r(x)$ to 0 for any $\nu \in \mathcal{P}(\Omega^2)$, $x \in \mathbb{R}$ and $r \geq 0$ as $k \rightarrow +\infty$. Indeed, by the boundedness of G , G_k , g and $\frac{\partial \phi}{\partial x}$ and the Lipschitz continuity of $x \rightarrow x^q$ for $0 \leq x \leq 1$, we have

$$\mathbb{E}^{\pi_\infty}((G - G_k)^2(\nu)) \leq C\mathbb{E}^{\pi_\infty} |G(\nu) - G_k(\nu)| \leq C\mathbb{E}^{\pi_\infty} \left(\langle \nu, \int_s^t |H - H_k| * \bar{\nu}_r(X_r) dr \rangle \right)$$

The second term is easy to deal with. Applying Itô's formula, we get

$$\mathbb{E}(G_n^2(\mu^n)) = \mathbb{E} \left(\left(\frac{1}{n} \sum_{i=1}^n g(X_{s_1}^{i,n}, Y_{s_1}^i, \dots, X_{s_p}^{i,n}, Y_{s_p}^i) \int_s^t \left(\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \right) (X_r^{i,n}, Y_r^i) dB_r^i \right)^2 \right) \leq \frac{C}{n}$$

and we conclude $\lim_{n \rightarrow +\infty} \mathbb{E}(G_n^2(\mu^n)) = 0$.

The third term is the most ticklish. By a calculation similar to the one carried out for the first term, we get

$$\mathbb{E}((G_k - G_n)^2(\mu^n)) \leq C\mathbb{E} \left(\langle \mu^n, \int_s^t |H_n - H_k| * \bar{\mu}_r^n(X_r) dr \rangle \right)$$

Hence if (X, Y, Z, W) denotes the canonical variable on Ω^4 ,

$$\mathbb{E}((G_k - G_n)^2(\mu^n)) \leq C\mathbb{E} \left(\langle \mu^n \otimes \mu^n, \int_s^t 1_{\{|X_r - Z_r| \leq \frac{1}{n\lambda k}\}} dr \rangle \right) \quad (1.12)$$

By the exchangeability of the couples $(X^{i,n}, Y^i)$, $1 \leq i \leq n$,

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \mathbb{E} \left(\langle \mu^n \otimes \mu^n, \int_s^t 1_{\{|X_r - Z_r| \leq \frac{1}{n\lambda k}\}} dr \rangle \right) &= \limsup_{n \rightarrow +\infty} \mathbb{E} \left(\int_s^t 1_{\{|X_r^{1,n} - X_r^{2,n}| \leq \frac{1}{n\lambda k}\}} dr \right) \\ &\leq \limsup_{n \rightarrow +\infty} \mathbb{E} \left(\int_s^t 1_{\{|X_r^{1,n} - X_r^{2,n}| \leq \frac{1}{k}\}} 1_{\{|X_r^{1,n}| \leq \sqrt{k}\}} dr \right) \\ &\quad + \limsup_{n \rightarrow +\infty} \int_s^t P(|X_r^{1,n}| \geq \sqrt{k}) dr \end{aligned} \quad (1.13)$$

Since $P(|X_r^{1,n}| \geq \sqrt{k}) \leq P(|B_r^1| \geq \frac{\sqrt{k}-r}{2}) + P(|X_0^1| \geq \frac{\sqrt{k}-r}{2})$, the second term of the right-hand-side of (1.13) has a limit equal to 0 when $k \rightarrow +\infty$. To prove that the same is true for the first term, we bound the L^2 norm of the density of $\mathcal{L}((X_r^{1,n}, X_r^{2,n}))$ ($r > 0$) uniformly in n . Like in the beginning of the proof of Lemma 1.3, we obtain an estimate similar to (1.6) :

$$\forall f \in L^2(\mathbb{R}^2), \forall n \geq 2, \forall r > 0, \mathbb{E}(f(X_r^{1,n}, X_r^{2,n})) \leq \frac{1}{\sqrt{2\pi r}} \exp(r) \|f\|_{L^2}$$

Hence $\forall n \geq 2$, $\mathbb{E} \left(\int_s^t \mathbf{1}_{\{|X_r^{1,n} - X_r^{2,n}| \leq \frac{1}{k}\}} \mathbf{1}_{\{|X_r^{1,n}| \leq \sqrt{k}\}} dr \right) \leq \frac{C}{k^{\frac{1}{4}}}$ which implies

$$\lim_{k \rightarrow +\infty} \limsup_{n \rightarrow +\infty} \mathbb{E} \left(\int_s^t \mathbf{1}_{\{|X_r^{1,n} - X_r^{2,n}| \leq \frac{1}{k}\}} \mathbf{1}_{\{|X_r^{1,n}| \leq \sqrt{k}\}} dr \right) = 0$$

With (1.12) and (1.13) we conclude $\lim_{k \rightarrow +\infty} \limsup_{n \rightarrow +\infty} \mathbb{E}((G_k - G_n)^2(\mu^n)) = 0$.

As we have proved that each term of the right-hand-side of (1.11) is equal to 0, $\mathbb{E}^{\pi_\infty}(G^2(\nu)) = 0$. Restricting $\phi, g, s_1, \dots, s_p, s, t$ to countable subsets then taking limits by Lebesgue's theorem, we obtain that π_∞ a.s., ν solves the martingale problem

$$\left\{ \begin{array}{l} \nu_0 = m \otimes m \quad \text{and} \quad \forall \phi \in C_b^2(\mathbb{R}^2), \\ \phi(X_t, Y_t) - \phi(X_0, Y_0) - \int_0^t \left(\frac{1}{2} \left(\frac{\partial^2 \phi}{\partial x^2} + 2 \frac{\partial^2 \phi}{\partial x \partial y} + \frac{\partial^2 \phi}{\partial y^2} \right) (X_s, Y_s) + \frac{\partial \phi}{\partial x} (X_s, Y_s) (H * \bar{\nu}_s(X_s))^q \right. \\ \left. + \frac{\partial \phi}{\partial y} (X_s, Y_s) (H * P_s(Y_s))^q \right) ds \quad \text{is a } \nu\text{-martingale} \end{array} \right.$$

Let us now suppose that ν is solution of this problem.

Choosing $\phi(x, y) = \psi(x)$ with $\psi \in C_b^2(\mathbb{R})$, we obtain that $\nu \circ X^{-1}$ solves the nonlinear martingale problem starting at m . By uniqueness for this problem $\nu \circ X^{-1} = P$ and $\bar{\nu}_s = P_s$. Moreover, it is easy to see that $\beta_t^1 = X_t - X_0 - \int_0^t (H * \bar{\nu}_s(X_s))^q ds = X_t - X_0 - \int_0^t (H * P_s(X_s))^q ds$ and $\beta_t^2 = Y_t - Y_0 - \int_0^t (H * P_s(Y_s))^q ds$ are ν -Brownian motions and next that $\beta^1 = \beta^2$. As ν a.s., $Y_0 = X_0$, by trajectorial uniqueness for the stochastic differential equation satisfied by both X and Y , ν a.s., $X = Y$. Hence $\nu = \hat{P}$.

We conclude that $\pi_\infty = \delta_{\hat{P}}$ which puts an end to the proof. ■

2 The moderate martingale problem

2.1 Existence and uniqueness

Let F be a measurable \mathbb{R}^d valued function on $[0, +\infty) \times \mathbb{R}^d \times \mathbb{R}$ bounded by M_F which satisfies the following Lipschitz continuity property

$$\exists K_F \geq 0, \forall s \in [0, +\infty), \forall x \in \mathbb{R}^d, \forall y, y' \in \mathbb{R}, |yF(s, x, y) - y'F(s, x, y')| \leq K_F |y - y'|$$

Definition 2.1 *Let $m \in \mathcal{P}(\mathbb{R}^d)$. We say that $P \in \tilde{\mathcal{P}}(\Omega)$ solves the nonlinear martingale problem (MP2) starting at m if $P_0 = m$ and for any $\phi \in C_b^2(\mathbb{R}^d)$*

$$\phi(X_t) - \phi(X_0) - \int_0^t \left(\frac{1}{2} \Delta \phi(X_s) + F(s, X_s, p(s, X_s)) \cdot \nabla \phi(X_s) \right) ds \text{ is a } P\text{-martingale} \quad (2.1)$$

where $p(s, x)$ is a measurable version of the densities for P .

This definition does not depend on the choice of the measurable version. Indeed, if $p'(s, x)$ is another such version then

$$P \text{ a.s.}, \forall t \geq 0, \int_0^t F(s, X_s, p(s, X_s)) \cdot \nabla \phi(X_s) ds = \int_0^t F(s, X_s, p'(s, X_s)) \cdot \nabla \phi(X_s) ds$$

Theorem 2.2 *For any $m \in \mathcal{P}(\mathbb{R}^d)$, the nonlinear problem (MP2) admits a unique solution P starting at m .*

Proof :

Uniqueness

It is an easy consequence of the Lipschitz continuity assumption made on F . The proof was given by Méléard and Roelly in [7].

Let P and Q be two solutions of (MP2) starting at m and $p(s, x)$, $q(s, x)$ denote measurable versions of the densities for P and Q . Using equation (1.2) given by Lemma 1.3, we get

$$\begin{aligned} \forall t > 0, \|p(t, \cdot) - q(t, \cdot)\|_{L^1} &\leq \sum_{i=1}^d \int_0^t \left\| \frac{\partial G_{t-s}}{\partial x_i} \right\|_{L^1} \|p(s, \cdot) F_i(s, \cdot, p(s, \cdot)) - q(s, \cdot) F_i(s, \cdot, q(s, \cdot))\|_{L^1} ds \\ &\leq \sqrt{d} K_F \int_0^t \frac{\|p(s, \cdot) - q(s, \cdot)\|_{L^1}}{\sqrt{t-s}} ds \end{aligned} \quad (2.2)$$

By Gronwall's lemma, we conclude that for any $t > 0$, $\|p(t, \cdot) - q(t, \cdot)\|_{L^1} = 0$. Hence both P and Q solve the martingale problem in which the nonlinearity in (2.1) is replaced by $q(s, X_s)$. By uniqueness for this problem, $P = Q$.

Existence

In the sequel, if I is a real interval and $v \in C(I, L^1(\mathbb{R}^d))$ let $v(t, x)$ denote a measurable function on $I \times \mathbb{R}^d$ such that for any $t \in I$ the class of $v(t, \cdot)$ in $L^1(\mathbb{R}^d)$ is $v(t)$.

Let $T > 0$. We define $A_T = \{v \in C((0, T], L^1(\mathbb{R}^d)); \sup_{t \in (0, T]} \|v(t)\|_{L^1} < +\infty\}$. For the metric $D(v, v') = \sup_{t \in (0, T]} \|v(t) - v'(t)\|_{L^1}$, A_T is complete.

Let $m \in \mathcal{P}(\mathbb{R}^d)$. For $v \in A_T$, we set

$$\forall t \in (0, T], \psi_m(v)(t) = G_t * m - \sum_{i=1}^d \int_0^t \frac{\partial G_{t-s}}{\partial x_i} * (v(s, \cdot) F_i(s, \cdot, v(s, \cdot))) ds$$

By the continuity of the map $t \rightarrow G_t \in L^1(\mathbb{R}^d)$, $t \rightarrow G_t * m \in L^1(\mathbb{R}^d)$ is continuous for $t > 0$. Since $\sup_{s \in (0, T]} \|v(s, \cdot) F_i(s, \cdot, v(s, \cdot))\|_{L^1} \leq M_F \sup_{s \in (0, T]} \|v(s, \cdot)\|_{L^1} < +\infty$, it is quite easy to deduce that $\psi_m(v) \in A_T$. Let us find T such that ψ_m is a contraction. For $v, v' \in A_T$ and $t \in (0, T]$, we get an estimate similar to (2.2)

$$\|\psi_m(v)(t) - \psi_m(v')(t)\|_{L^1} \leq \sqrt{d} K_F \int_0^t \frac{\|v(s) - v'(s)\|_{L^1}}{\sqrt{t-s}} ds \leq 2K_F \sqrt{dt} D(v, v')$$

Hence $D(\psi_m(v), \psi_m(v')) \leq 2K_F \sqrt{dT} D(v, v')$. From now on, $T = \frac{1}{16dK_F^2}$. By Picard's fixed-point theorem, ψ_m admits a unique fixed-point in A_T .

Let $t_0 \geq 0$ and $f \in L^1(\mathbb{R}^d)$. For $v \in C([0, T], L^1(\mathbb{R}^d))$ we define

$$\tilde{\psi}_{t_0, f}(v)(t) = G_t * f - \sum_{i=1}^d \int_0^t \frac{\partial G_{t-s}}{\partial x_i} * (v(s, \cdot) F_i(t_0 + s, \cdot, v(s, \cdot))) ds$$

The same estimates as above imply that $\tilde{\psi}_{t_0, f}$ admits a unique fixed-point in $C([0, T], L^1(\mathbb{R}^d))$.

Let v^0 denote the fixed-point of ψ_m in A_T . If v^n is constructed, let v^{n+1} be the fixed-point of $\tilde{\psi}_{(n+1)T, v^n(T)}$ in $C([0, T], L^1(\mathbb{R}^d))$. We set $v(t) = v^n(t - nT)$ if $t \in (nT, (n+1)T]$. The map v belongs to $C((0, +\infty), L^1(\mathbb{R}^d))$ and satisfies

$$\forall t_0 > 0, \quad \sup_{t \in (0, t_0]} \|v(t)\|_{L^1} < +\infty \quad (2.3)$$

Let $t \in (0, T]$. We compute $v(T+t)$ thanks to Fubini's theorem.

$$\begin{aligned} v(T+t) &= G_t * v(T) - \sum_{i=1}^d \int_0^t \frac{\partial G_{t-s}}{\partial x_i} * (v(T+s, \cdot) F_i(T+s, \cdot, v(T+s, \cdot))) ds \\ &= G_t * \left(G_T * m - \sum_{i=1}^d \int_0^T \frac{\partial G_{T-s}}{\partial x_i} * (v(s, \cdot) F_i(s, \cdot, v(s, \cdot))) ds \right) \\ &\quad - \sum_{i=1}^d \int_0^t \frac{\partial G_{t-s}}{\partial x_i} * (v(T+s, \cdot) F_i(T+s, \cdot, v(T+s, \cdot))) ds \\ &= G_t * G_T * m - \sum_{i=1}^d \int_0^T G_t * \left(\frac{\partial G_{T-s}}{\partial x_i} * (v(s, \cdot) F_i(s, \cdot, v(s, \cdot))) \right) ds \\ &\quad - \sum_{i=1}^d \int_0^t \frac{\partial G_{t-s}}{\partial x_i} * (v(T+s, \cdot) F_i(T+s, \cdot, v(T+s, \cdot))) ds \\ &= G_{T+t} * m - \sum_{i=1}^d \int_0^{T+t} \frac{\partial G_{T+t-s}}{\partial x_i} * (v(s, \cdot) F_i(s, \cdot, v(s, \cdot))) ds \end{aligned}$$

By induction, we conclude

$$\forall t > 0, \quad v(t, x) = G_t * m(x) - \sum_{i=1}^d \int_0^t \frac{\partial G_{t-s}}{\partial x_i} * (v(s, \cdot) F_i(s, \cdot, v(s, \cdot)))(x) ds \quad \text{almost surely} \quad (2.4)$$

Let P be the solution of the martingale problem in which the nonlinearity in (2.1) is replaced by $v(s, X_s)$. Equations (2.3), (2.4) and Lemma 1.3 imply that $v(s, x)$ is a measurable version of the densities for P . Hence P solves (MP2). \blacksquare

2.2 Application

Theorem 2.1 allows us to associate a probabilistic representation with some classical solutions of Burgers' equation. The initial conditions concerned are not distribution functions like in Proposition 1.4 but bounded probability densities.

We take up Oelschläger's approach in [9] (pages 306-307). Let u_0 be a probability density on \mathbb{R} bounded by M . The Cole-Hopf transformation (see Cole [2] and Hopf [5])

$$u(0, x) = u_0(x) \quad \text{and} \quad \forall t > 0, \quad u(t, x) = \frac{\int_{\mathbb{R}} G_t(x-y) \exp\left(-\int_{-\infty}^y u_0(z) dz\right) u_0(y) dy}{\int_{\mathbb{R}} G_t(x-y) \exp\left(-\int_{-\infty}^y u_0(z) dz\right) dy}$$

provides a classical solution of Burgers' equation. Indeed, $u \in C^{1,2}((0, +\infty) \times \mathbb{R})$ and

$$\forall t > 0, \forall x \in \mathbb{R}, \frac{\partial u}{\partial t}(t, x) = \frac{1}{2} \frac{\partial^2 u}{\partial x^2}(t, x) - u(t, x) \frac{\partial u}{\partial x}(t, x) \quad (2.5)$$

It is easy to check that $\forall t > 0, \forall x \in \mathbb{R}, |u(t, x)| \leq M$. This boundedness property is essential for the sequel. We set $f(y) = \frac{1}{2}(0 \vee y \wedge M)$. The functions f and $y \rightarrow yf(y)$ are respectively bounded and Lipschitz continuous.

By theorem 2.2, the martingale problem (MP2) corresponding to the choice $F(s, x, y) = f(y)$ admits a unique solution P starting at $u_0(x)dx$. Let us prove that u is a measurable version of the densities for P . Since clearly $\forall t \geq 0, \|u(t, \cdot)\|_{L^1} \leq e$, according to the proof of uniqueness for (MP2) (theorem 2.2), it is enough to establish

$$\forall t > 0, \forall x \in \mathbb{R}, u(t, x) = G_t * u_0(x) - \int_0^t \frac{\partial G_{t-s}}{\partial x} * (u(s, \cdot) f(u(s, \cdot)))(x) ds \quad (2.6)$$

Let $t > 0, \phi$ be a $C^{1,2}$ function with compact support in $[0, t] \times \mathbb{R}$ and $\epsilon \in (0, t)$. As $\frac{\partial u}{\partial s}, \frac{\partial^2 u}{\partial x^2}$ and $\frac{\partial}{\partial x}(uf(u)) = u \frac{\partial u}{\partial x}$ are bounded on the intersection of the support of ϕ with $[\epsilon, t] \times \mathbb{R}$, using the integration by parts formula, Fubini's theorem and (2.5) we get

$$\begin{aligned} \int_{\mathbb{R}} u(t, x) \phi(t, x) dx &= \int_{\mathbb{R}} u(\epsilon, x) \phi(\epsilon, x) dx \\ &+ \int_{(\epsilon, t] \times \mathbb{R}} u(s, x) \left(\frac{\partial \phi}{\partial s}(s, x) + \frac{1}{2} \frac{\partial^2 \phi}{\partial x^2}(s, x) + f(u(s, x)) \frac{\partial \phi}{\partial x}(s, x) \right) dx ds \end{aligned} \quad (2.7)$$

We have $\lim_{s \rightarrow 0} \|u(s, \cdot) - u_0\|_{L^1} = 0$. Indeed for $U(x) = \exp\left(-\int_{-\infty}^x u_0(z) dz\right)$,

$$\begin{aligned} \|u(s, \cdot) - u_0\|_{L^1} &\leq \left\| \frac{1}{G_s * U} \right\|_{L^\infty} \|G_s * (U u_0) - (G_s * U) u_0\|_{L^1} \\ &\leq e \|G_s * (U u_0) - U u_0\|_{L^1} + e \|(G_s * U - U) u_0\|_{L^1} \end{aligned}$$

Since $U u_0 \in L^1(\mathbb{R})$, the first term of the right hand side converges to 0 when $s \rightarrow 0$. The continuity and the boundedness of U imply that $G_s * U$ is bounded uniformly in s and converges pointwise to U . Hence, by Lebesgue's theorem, the second term also goes to 0.

Thus $\lim_{s \rightarrow 0} \int_{\mathbb{R}} u(s, x) \phi(s, x) dx = \int_{\mathbb{R}} u_0(x) \phi(0, x) dx$ and taking the limit $\epsilon \rightarrow 0$ in (2.7), we get

$$\begin{aligned} \int_{\mathbb{R}} u(t, x) \phi(t, x) dx &= \int_{\mathbb{R}} u_0(x) \phi(0, x) dx \\ &+ \int_{(0, t] \times \mathbb{R}} u(s, x) \left(\frac{\partial \phi}{\partial s}(s, x) + \frac{1}{2} \frac{\partial^2 \phi}{\partial x^2}(s, x) + f(u(s, x)) \frac{\partial \phi}{\partial x}(s, x) \right) dx ds \end{aligned}$$

By spatial truncation, this equation still holds if $\phi \in C_b^{1,2}([0, t] \times \mathbb{R})$. For the particular choice $\phi(s, x) = G_{t-s} * \psi(x)$ with ψ C^2 with compact support in \mathbb{R} , we conclude like in the proof of Lemma 1.3 that (2.6) holds. Therefore $u(t, x)$ is a measurable version of the densities for P and P provides a probabilistic representation of u .

3 Extension of the results to martingale problems with a non-constant diffusion coefficient

Let a be a Lipschitz continuous map on \mathbb{R}^d with values in the set of symmetric non-negative $d \times d$ matrices satisfying for $M_a \geq m_a > 0, \forall x, y \in \mathbb{R}^d, m_a |y|^2 \leq y^* a(x) y \leq M_a |y|^2$ and L be

the operator $L\phi(x) = \frac{1}{2} \sum_{i,j=1}^d a_{i,j}(x) \frac{\partial^2 \phi}{\partial x_i \partial x_j}(x)$.

Let σ denote the square-root of a . By the assumptions made on a , the map $x \rightarrow \sigma(x)$ is bounded and Lipschitz continuous. According to Friedman [3] pages 139-150, there is a transition density $\Gamma_s(x, y)$, $s > 0$, $x, y \in \mathbb{R}^d$ associated with the time-homogeneous stochastic differential equation $dX_t = \sigma(X_t)dB_t$.

Moreover, for any $t > 0$ and any continuous function ψ with compact support in \mathbb{R}^d , the function $\phi(s, x) = \int_{\mathbb{R}^d} \Gamma_{t-s}(x, y)\psi(y)dy$ defined on $[0, t) \times \mathbb{R}^d$ satisfies

$$\forall s \in [0, t), \forall x \in \mathbb{R}^d, L\phi(s, x) + \frac{\partial \phi}{\partial s}(s, x) = 0 \text{ and } \forall x \in \mathbb{R}, \lim_{s \rightarrow t} \phi(s, x) = \psi(x) \quad (3.1)$$

and $\forall M > M_a$ there is a constant $C(t)$ such that,

$$\forall s \in (0, t], \forall x, y \in \mathbb{R}^d, \Gamma_s(x, y) \leq \frac{C(t)}{s^{\frac{d}{2}}} \exp\left(-\frac{|x-y|^2}{2Ms}\right) \quad (3.2)$$

$$\forall s \in (0, t], \forall x, y \in \mathbb{R}^d, \forall 1 \leq i \leq d, \left| \frac{\partial \Gamma_s(x, y)}{\partial x_i} \right| \leq \frac{C(t)}{s^{\frac{d+1}{2}}} \exp\left(-\frac{|x-y|^2}{2Ms}\right) \quad (3.3)$$

By (3.3), we obtain the following estimate

$$\forall t > 0, \exists K(t) > 0, \forall s \in (0, t], \forall x \in \mathbb{R}^d, \forall 1 \leq i \leq d, \left\| \frac{\partial \Gamma_s(x, y)}{\partial x_i} \right\|_{L_y^1} \leq \frac{K(t)}{\sqrt{s}} \quad (3.4)$$

Theorem 3.1 $\forall m \in \mathcal{P}(\mathbb{R}^d)$, the martingale problem (MP1) (resp (MP2)) in which $\frac{1}{2}\Delta\phi(X_s)$ is replaced by $L\phi(X_s)$ in (1.1) (resp in (2.1)) admits a unique solution starting at m .

The proofs of Theorem 1.2 and Theorem 2.2 are based on Lemma 1.3. Therefore we explain how to adapt the conclusions and the proof of this lemma. As σ is Lipschitz continuous and bounded, for any $m \in \mathcal{P}(\mathbb{R}^d)$, the martingale problem : $P_0 = m$ and

$$\forall \phi \in C_b^2(\mathbb{R}^d), \phi(X_t) - \phi(X_0) - \int_0^t L\phi(X_s)ds \text{ is a } P\text{-martingale}$$

admits a unique solution P . Moreover, by the existence of Γ , for any $t > 0$, P_t has a density equal to $\int_{\mathbb{R}^d} \Gamma_t(x, y)m(dx)$. For g like in Lemma 1.3, by Girsanov's theorem, as $\sigma^{-1}g$ is bounded, the martingale problem with $L\phi(X_s) + g(s, X_s) \cdot \nabla\phi(X_s)$ replacing $L\phi(X_s)$ admits a unique solution and this solution belongs to $\tilde{\mathcal{P}}(\Omega)$. Let $p(s, x)$ be a measurable version of the densities for the solution. If ψ is a continuous function with compact support on \mathbb{R}^d and $\phi(s, x) = \int_{\mathbb{R}^d} \Gamma_{t-s}(x, y)\psi(y)dy$, by the uniform continuity of ψ and (3.2), the convergence of $\phi(s, x)$ to $\psi(x)$ in (3.1) is uniform in $x \in \mathbb{R}^d$. By (3.4), we upper-bound $\nabla\phi(s, x)$. These two remarks allow to transpose the proof of (1.2) and obtain

$$\forall t > 0, p(t, y) = \int_{\mathbb{R}^d} \Gamma_t(x, y)m(dx) + \int_0^t \int_{\mathbb{R}^d} \nabla_x \Gamma_{t-s}(x, y) \cdot g(s, x) p(s, x) dx ds \text{ almost surely}$$

With this equation and (3.4) instead of (1.2) and (0.1), we easily adapt the proofs of Theorem 1.2 and Theorem 2.2.

References

- [1] M. Bossy and D. Talay. Convergence rate for the approximation of the limit law of weakly interacting particles: Application to the burgers equation. *Annals of applied prob.*, To appear.
- [2] J.D. Cole. On a quasi-linear parabolic equation occuring in aerodynamics. *Quart. Appl. Math.*, 9(3):225–236, 1951.
- [3] A. Friedman. *Stochastic Differential Equations and Applications*. Academic Press, 1975.
- [4] C. Graham. Nonlinear diffusions with jumps. *Ann. Inst. Henri Poincaré*, 28(3):393–402, 1992.
- [5] E. Hopf. The partial differential equation $u_t + uu_x = \mu u_{xx}$. *Comm. Pure Appl. Math.*, 3:201–230, 1950.
- [6] I. Karatzas and S.E. Shreve. *Brownian Motion and Stochastic Calculus*. Springer-Verlag, 1988.
- [7] S. Méléard and S. Roelly-Coppoletta. A propagation of chaos result for a system of particles with moderate interaction. *Stochastic Processes and their Application*, 26:317–332, 1987.
- [8] P.A. Meyer. *Probabilités et Potentiel*. Hermann, 1966.
- [9] K. Oelschläger. A law of large numbers for moderately interacting diffusion processes. *Z. Wahrsch. Verw. Geb.*, 69:279–322, 1985.
- [10] A.S. Sznitman. Topics in propagation of chaos. In *Ecole d'été de probabilités de Saint-Flour XIX - 1989, Lect. Notes in Math. 1464*. Springer-Verlag, 1991.