

Multiscale Finite Element approach for "weakly" random problems and related issues

Claude Le Bris, Frédéric Legoll, Florian Thomines

École Nationale des Ponts et Chaussées, 6 et 8 avenue Blaise Pascal, 77455 Marne-La-Vallée Cedex 2
and INRIA Rocquencourt, MICMAC team-project, Domaine de Voluceau, B.P. 105,
78153 Le Chesnay Cedex, France
lebris@cermics.enpc.fr, {legoll,thominef}@lami.enpc.fr

November 8, 2011

Abstract

We address multiscale elliptic problems with random coefficients that are a perturbation of multiscale deterministic problems. Our approach consists in taking benefit of the perturbative context to suitably modify the classical Finite Element basis into a deterministic multiscale Finite Element basis. The latter essentially shares the same approximation properties as a multiscale Finite Element basis directly generated on the random problem. The specific reference method that we use is the Multiscale Finite Element Method. Using numerical experiments, we demonstrate the efficiency of our approach and the computational speed-up with respect to a more standard approach. We provide a complete analysis of the approach, extending that available for the deterministic setting.

1 Overview of our approach and results

The Multiscale Finite Element Method (henceforth abbreviated as MsFEM) is a popular numerical approach for multiscale problems (see [37, 38, 30, 32, 31, 3, 39, 20, 18, 11]). It consists in a Galerkin approximation of the original problem over a finite dimensional space generated by basis functions that are specifically *adapted* to the problem under consideration.

This approach is popular for a twofold reason. First, its use is not restricted to multiscale problems that converge to a homogenized problem in the limit of vanishing ratio between the small scale and the macroscopic scale. It may be applied to much more general situations. Second, when the problem does converge to a homogenization problem, the MsFEM approach is meant to approximate the solution of the problem with the small scale ε at its actual small value and not "only" in the asymptotic regime $\varepsilon \rightarrow 0$, which is the regime addressed by homogenization theory.

To fix the ideas, consider the problem of finding u^ε solving

$$-\operatorname{div}[A^\varepsilon \nabla u^\varepsilon] = f \quad \text{in } \mathcal{D}, \quad u^\varepsilon = 0 \quad \text{on } \partial\mathcal{D}, \quad (1)$$

on a bounded domain $\mathcal{D} \subset \mathbb{R}^d$, with $f \in L^2(\mathcal{D})$, and where A^ε is a uniformly bounded, coercive matrix that varies at scale ε . A standard Finite Element Method (FEM) would require a space discretization of the domain at the scale ε in order to capture the oscillations of u^ε at scale ε . This is prohibitively expensive. The MsFEM aims at accurately approximating u^ε using a limited number of degrees of freedom. It does not require the matrix A^ε to be periodic (namely $A^\varepsilon(x) = A_{per}(x/\varepsilon)$ for a fixed periodic matrix A_{per}) or stationary.

We now briefly describe the approach and present the aim of this article. Starting from a coarse mesh \mathcal{T}_h with a standard (say \mathbb{P}_1) Finite Element basis set of functions $\{\phi_i^0\}_{i=1}^L$, generating the associated space

$$\mathcal{V}_h := \operatorname{span}(\phi_i^0, i = 1, \dots, L),$$

we first numerically build the MsFEM basis functions ϕ_i^ε . Several definitions of these basis functions have been proposed in the literature (yielding different numerical methods), and we detail this in the sequel (see e.g. (10)-(11)-(12)). For

the moment, it is sufficient to know that, to each ϕ_i^0 , which varies at the macroscopic scale, is associated a function ϕ_i^ε , with variations at the scale ε . In practice, ϕ_i^ε is *numerically* computed (in fact, *pre-computed*), using the specificities of the problem addressed. These highly oscillatory functions ϕ_i^ε generate the finite dimensional space

$$\mathcal{W}_h := \text{span}(\phi_i^\varepsilon, i = 1, \dots, L).$$

Note that \mathcal{W}_h and \mathcal{V}_h share the same dimension.

We next define the MsFEM solution u_M using a Galerkin approximation of (1) on \mathcal{W}_h , instead of \mathcal{V}_h . Again, details will be given below. The MsFEM solution u_M provided by the approach reads

$$u_M(x) = \sum_{i=1}^L (U_M)_i \phi_i^\varepsilon(x),$$

for some coefficients $\{(U_M)_i\}_{i=1}^L$. Of course, these coefficients depend on ε , but this dependency is kept implicit in the sequel.

We now turn our attention to the stochastic problem

$$-\text{div}[A^\varepsilon(\cdot, \omega) \nabla u^\varepsilon(\cdot, \omega)] = f \text{ in } \mathcal{D}, \quad u^\varepsilon(\cdot, \omega) = 0 \text{ on } \partial\mathcal{D}, \quad (2)$$

and a typical quantity of interest $\mathbb{E}[u^\varepsilon(x, \cdot)]$, which is traditionally approximated using a Monte Carlo method. Introducing a set of \mathcal{M} realizations of the stochastic matrix $\{A^{\varepsilon, m}\}_{1 \leq m \leq \mathcal{M}}$, a direct, naive application of the MsFEM paradigm would consist in first computing for each realization m the stochastic MsFEM basis functions $\phi_i^{\varepsilon, m}(x, \omega)$, next performing a Galerkin approximation of (2) using this MsFEM basis set to compute $\{u_M^m(x, \omega)\}_{1 \leq m \leq \mathcal{M}}$, and eventually approximating $\mathbb{E}[u^\varepsilon(x, \cdot)]$ by

$$\mathbb{E}[u^\varepsilon(x, \cdot)] \approx \frac{1}{\mathcal{M}} \sum_{m=1}^{\mathcal{M}} u_M^m(x, \omega).$$

Such an approach is unpractical because of the prohibitively expensive computational load.

To reduce the computational cost and make the MsFEM approach practical in such a stochastic context, a natural idea we investigate in this article is to consider a less generic setting, for which a dedicated, more computationally affordable approach, can be designed. One possibility is to consider matrices $A^\varepsilon(x, \omega) \equiv A^\varepsilon(x) + B(x, \omega)$ in (2) that are not highly oscillatory in their stochastic part. In such cases, dedicated approaches have been proposed, we refer to [35] for more details. Another approach is to reduce the number of Monte-Carlo simulations used for the computation of the multiscale basis functions. In [1, 26], the authors assume that their coefficient can be written as a Karhunen-Loève type expansion, and apply a collocation method to *a priori* choose some sparse realizations for which they compute the multiscale basis functions.

In this article, we consider one of the many alternate variants of problem (2). We suppose that $A^\varepsilon(x, \omega)$ is highly oscillatory in both its deterministic and stochastic components, but that it is a *perturbation* of a deterministic matrix. More precisely, we assume that

$$A^\varepsilon(x, \omega) \equiv A_\eta^\varepsilon(x, \omega) = A_0^\varepsilon(x) + \eta A_1^\varepsilon(x, \omega), \quad (3)$$

where A_0^ε is a deterministic matrix and η is a small deterministic parameter. This model may be well suited for heterogeneous materials (or, more generally, media) that, although not periodic, are *not fully* stochastic, in the sense that they may be considered as a *perturbation* of a deterministic material. We call this setting the *weakly stochastic setting*. Note that many practical situations, involving actual materials or media, can be considered, at a good level of approximation, as perturbations of a deterministic (often periodic) setting (see e.g. [41]).

In a series of recent works (see [14, 15, 25] and [6, 7, 8]; see also [5] for a unified presentation), we have considered such a setting, in the context of homogenization theory (the matrix $A_\eta^\varepsilon(x, \omega)$ in (2)-(3) reads $A_\eta^\varepsilon(x, \omega) = A_\eta(x/\varepsilon, \omega)$ for a stationary matrix $A_\eta(x, \omega)$, which is, in a sense to be made precise, a perturbation of a periodic matrix). We have shown there that, in such a case, the workload for computing the homogenized solution is significantly lighter than for *generic* stochastic homogenization, and actually comparable to the workload for periodic homogenization. We will show in the sequel that the MsFEM can be adapted to this weakly stochastic setting, providing an approximation of the solution u_η^ε to (2)-(3), for fixed ε , at a much smaller computational cost than the direct approach.

The main idea of our proposed approach is to compute a set of *deterministic* MsFEM basis functions using A_0^ε , the deterministic part of A_η^ε in the expansion (3), and then to perform Monte Carlo realizations at the macroscale level using a set of \mathcal{M} realizations of the random matrix $\{A_\eta^{\varepsilon,m}(x, \omega)\}_{1 \leq m \leq \mathcal{M}}$ (see Section 2 for a detailed presentation). Note that, for each of these realizations, we solve the *original* problem, with the *complete* matrix A_η^ε , and not only its deterministic part. Only the basis set is taken deterministic. By construction, the approach provides an approximation

$$u_S(x, \omega) = \sum_{i=1}^L (U_S(\omega))_i \phi_i^\varepsilon(x)$$

of $u_\eta^\varepsilon(x, \omega)$, where the basis functions ϕ_i^ε are *deterministic*. These basis functions are computed only once, hence the cost to compute $\{u_S^m(x, \omega)\}_{1 \leq m \leq \mathcal{M}}$ is much smaller than the cost to compute $\{u_M^m(x, \omega)\}_{1 \leq m \leq \mathcal{M}}$. This is especially true if (2) has to be solved for many right-hand sides f . We expect that this approximation u_S is as accurate as u_M for small η . We show below that this is indeed the case, when A_η^ε is a perturbation of A_0^ε (see Section 3 for numerical tests).

We would like to note that the MsFEM is not the only multiscale technique based on finite elements. The bottom line of our approach, consisting of generating suitable multiscale functions for the discretization of a weakly stochastic problem, using for this purpose the deterministic reference problem, can in principle be applied to other multiscale techniques. Another popular technique is the HMM method [27, 28, 29], for which our approach could in principle be easily adapted.

In the numerical tests reported on in Section 3, we compare, in the H^1 norm, u_η^ε (the exact solution to (2) with the matrix $A^\varepsilon \equiv A_\eta^\varepsilon$ given by (3)) with u_S (the solution provided by our approach) and u_M (the solution provided by the ideal, expensive approach). The quantity $\|u_\eta^\varepsilon - u_M\|_{H^1(\mathcal{D})}$ somewhat represents the best possible accuracy that we can achieve, in the sense that our approach inherits the limitations of the MsFEM approach. We thus cannot expect our approximation u_S to be more accurate than u_M . We can only hope to compute an approximation of comparable quality with a much reduced workload. The numerical results we obtain confirm that, for small η in (3), the quantity $\|u_S - u_\eta^\varepsilon\|_{H^1(\mathcal{D})}$ is of the same order of magnitude as $\|u_M - u_\eta^\varepsilon\|_{H^1(\mathcal{D})}$, although, we repeat it, the computational cost to compute u_S is much smaller than that to compute u_M .

We next derive error bounds for our approach in Section 4. We recall that, in the deterministic setting, a classical context for proving convergence of the MsFEM approach is the case when, in the reference problem (1), the matrix reads $A^\varepsilon(x) = A_{per}\left(\frac{x}{\varepsilon}\right)$ for a fixed periodic matrix A_{per} . Likewise, to be able to perform our theoretical analysis in the stochastic setting, we assume in Section 4 that $A_\eta^\varepsilon(x, \omega) = A_\eta\left(\frac{x}{\varepsilon}, \omega\right)$ for a fixed stationary random matrix A_η . The problem (2)-(3) then admits a homogenized limit when ε vanishes.

Our proof follows the same lines as that in the deterministic setting, which we now briefly review (see the introduction of Section 4 for more details on the structure of the proof). The MsFEM is a Galerkin approximation, that we assume momentarily, for the sake of clarity, to be a *conforming* approximation (this is indeed the case when, for defining the highly oscillatory basis functions ϕ_i^ε , oversampling is *not* used). The error is then estimated using the C ea lemma:

$$\|u^\varepsilon - u_M\|_{H^1(\mathcal{D})} \leq C \inf_{v_h \in \mathcal{W}_h} \|u^\varepsilon - v_h\|_{H^1(\mathcal{D})},$$

where u^ε is the solution to the reference deterministic highly oscillatory problem (1), u_M is the MsFEM solution and C is a constant independent of ε and h . Taking advantage of the homogenization setting, we introduce the two-scale expansion

$$v^\varepsilon = u^* + \varepsilon \sum_{i=1}^d w_{e_i}^0 \left(\frac{\cdot}{\varepsilon}\right) \partial_i u^*$$

of u^ε , where u^* is the homogenized solution, $w_{e_i}^0$ is the periodic corrector associated to $e_i \in \mathbb{R}^d$, and $\partial_i u^*$ denotes the partial derivative $\frac{\partial u^*}{\partial x_i}$. We next write

$$\|u^\varepsilon - u_M\|_{H^1(\mathcal{D})} \leq C \left(\|u^\varepsilon - v^\varepsilon\|_{H^1(\mathcal{D})} + \inf_{v_h \in \mathcal{W}_h} \|v^\varepsilon - v_h\|_{H^1(\mathcal{D})} \right).$$

The first term in the above right-hand side is estimated using standard homogenization results on the *rate* of convergence of $v^\varepsilon - u^\varepsilon$. To estimate the second term, one considers a suitably chosen element $v_h \in \mathcal{W}_h$, for which $\|v^\varepsilon - v_h\|_{H^1(\mathcal{D})}$ can be directly bounded.

Following the same strategy in our stochastic setting, we estimate the distance between the solution u_η^ε to the reference stochastic problem (2)-(3) and the weakly stochastic MsFEM solution u_S as

$$\|u_\eta^\varepsilon(\cdot, \omega) - u_S(\cdot, \omega)\|_{H^1(\mathcal{D})} \leq C \left(\|u_\eta^\varepsilon(\cdot, \omega) - v_\eta^\varepsilon(\cdot, \omega)\|_{H^1(\mathcal{D})} + \inf_{v_h \in \mathcal{W}_h} \|v_\eta^\varepsilon(\cdot, \omega) - v_h\|_{H^1(\mathcal{D})} \right).$$

We observe that a key ingredient for the proof is the rate of convergence of the difference between the reference solution u_η^ε and its two-scale expansion v_η^ε . Such a result is classical in periodic homogenization, but, to the best of our knowledge, open in the *general* stationary case (in dimensions higher than one). One only knows that $u_\eta^\varepsilon - v_\eta^\varepsilon$ vanishes (in some appropriate norm) when $\varepsilon \rightarrow 0$. However, in the particular case when A_η^ε is only *weakly* stochastic, it is possible to obtain such a result, as we have shown in [42]. Hence, exploiting the specificity of our weakly stochastic setting, we are able to obtain (see our main result, Theorem 10 and estimate (63)):

$$\sqrt{\mathbb{E} \left[\|u_\eta^\varepsilon - u_S\|_{H_h^1}^2 \right]} \leq C \left(\sqrt{\varepsilon} + h + \frac{\varepsilon}{h} + \eta \left(\frac{\varepsilon}{h} \right)^{d/2} \ln(N(h)) + \eta + \eta^2 \mathcal{C}(\eta) \right),$$

where $\|\cdot\|_{H_h^1}$ is a broken H^1 norm, C is a constant independent of ε , h and η , \mathcal{C} is a bounded function as η goes to 0, and $N(h)$ is the number of elements in the mesh (roughly of order h^{-d} in dimension d). As is often the case in the deterministic setting, we use here (both for our numerical tests and in the analysis) the oversampling technique. Consequently, the basis functions ϕ_i^ε do not belong to $H_0^1(\mathcal{D})$, hence the use of a broken H^1 norm in the above estimate. As we point out below, when $\eta = 0$ in (3), our approach reduces to the standard deterministic MsFEM (with oversampling), and the above estimates then agree with those proved in [32].

This article is organized as follows. First, in Section 2, we describe the MsFEM approach. For consistency, we begin by the deterministic setting in Section 2.1, and point out there that the direct adaptation to the general stochastic setting yields a prohibitively expensive approach. The adaptation of the approach to the *weakly stochastic setting* is described in Section 2.2. We next turn to numerical simulations, in Section 3. Some procedures to efficiently implement the approach are first described in Section 3.1. We next consider a one-dimensional test (see Section 3.2), which is useful for several reasons. First, it allows to calibrate some numerical parameters, such as the number M of independent realizations when estimating the exact expectation by an empirical mean. Second, we assess the accuracy of our approach with respect to the magnitude of η . We demonstrate there that η does not have to be extremely small for our method to be very efficient. For instance, on the test case considered in Section 3.2, we show that our approach is as accurate as the expensive, direct approach as soon as η is such that

$$\left\| \frac{\eta a_1^\varepsilon}{a_0^\varepsilon} \right\|_{L^\infty(\mathbb{R} \times \Omega)} \text{ is equal to or smaller than } 0.1,$$

where a_0^ε is the deterministic component of the diffusion coefficient a_η^ε and ηa_1^ε is the stochastic component (see expansion (3)). Lastly, we also assess the accuracy of our approach with respect to the presence of frequencies in the random coefficient a_η^ε that are not taken into account in the MsFEM basis set. We next turn to two test cases in dimension two, where we observe that our approach behaves as well as in the one-dimensional case (see Section 3.3). In particular, in Section 3.3.2, we successfully address a classical test-case of the literature.

Section 4 is devoted to the analysis of the approach, in the homogenization setting. Our main result, Theorem 10, is presented in Section 4.1, and proved in Section 4.2. The proofs of some technical results are collected in Appendix A. In addition, we specifically consider the one dimensional case in Section 4.3.

2 MsFEM-type approaches

For consistency and the convenience of the reader, we present in this section the MsFEM approach to solve the original elliptic problem (1). For clarity, we begin by presenting the approach in a deterministic setting. The reader familiar with the MsFEM may easily skip this section and directly proceed to Section 2.2, where we present our approach in a weakly stochastic setting.

2.1 Description in a classical deterministic setting

Let $u^\varepsilon \in H^1(\mathcal{D})$ be the solution to (1), where the matrix $A^\varepsilon \in (L^\infty(\mathcal{D}))^{d \times d}$ satisfies the standard coercivity condition: there exists two constants $a_+ \geq a_- > 0$ such that

$$\forall \varepsilon, \quad \forall x \in \mathcal{D}, \quad \forall \xi \in \mathbb{R}^d, \quad a_- |\xi|^2 \leq \xi^T A^\varepsilon(x) \xi \quad \text{and} \quad \|A^\varepsilon\|_{L^\infty(\mathcal{D})} \leq a_+.$$

Note that the MsFEM approach is not restricted to the periodic setting. We therefore do not assume that $A^\varepsilon(x) = A_{per}(x/\varepsilon)$ for a fixed periodic matrix A_{per} .

The MsFEM approach consists in performing a variational approximation of (1) where the basis functions are *precomputed* and *encode the fast oscillations* present in (1). In the sequel we argue on the following formulation, equivalent to (1):

$$\text{Find } u^\varepsilon \in H_0^1(\mathcal{D}) \text{ such that, for any } v \in H_0^1(\mathcal{D}), \quad \mathcal{A}_\varepsilon(u^\varepsilon, v) = b(v), \quad (4)$$

where

$$\mathcal{A}_\varepsilon(u, v) = \int_{\mathcal{D}} (\nabla v(x))^T A^\varepsilon(x) \nabla u(x) dx \quad \text{and} \quad b(v) = \int_{\mathcal{D}} f(x) v(x) dx.$$

The MsFEM is a three-step approach:

1. introduce a standard discretization of the domain \mathcal{D} using a coarse mesh as compared to the small scale oscillations of A^ε .
2. for each element \mathbf{K} of the coarse mesh, compute the basis function $\phi_i^{\varepsilon, \mathbf{K}}$ as the solution of an elliptic equation posed in \mathbf{K} (see e.g. (10)-(11)-(12) below).
3. solve the Galerkin approximation of (4), for the set of basis functions defined at Step 2.

The advantage of the approach is that, for the same accuracy of the approximation as that provided by a standard FEM, the macroscale mesh can be chosen sufficiently coarse so that the resulting discretized problem has a limited number of degrees of freedom, and may thus be computationally solved inexpensively. This is observed in practice [37], and proven by a theoretical analysis (see [38, 32]) when the problem (4) admits a homogenized limit. See also [30] and references therein.

To further illustrate this fact, we reproduce here a simple one-dimensional analysis we borrow from A. Lozinsky (see [44, Chap. 6] and [17]). This analysis explains remarkably well the interest of the approach, and, in contrast to [38, 32], is not restricted to a homogenization setting. Consider the one-dimensional domain $\mathcal{D} = (0, 1)$ and the reference problem

$$\mathcal{L}u = f, \quad u(0) = u(1) = 0,$$

for the operator $\mathcal{L}u := -(\nu u)'$, where $f \in L^2(0, 1)$ and $\nu \in L^\infty(0, 1)$ with $\nu(x) \geq \nu_{min} > 0$ almost everywhere on $(0, 1)$. The function ν may have oscillations at a small scale. The associated weak formulation reads

$$\text{Find } u \in H_0^1(0, 1) \text{ such that, for any } v \in H_0^1(0, 1), \quad a(u, v) = b(v), \quad (5)$$

with

$$a(u, v) = \int_0^1 \nu(x) u'(x) v'(x) dx \quad \text{and} \quad b(v) = \int_0^1 f(x) v(x) dx.$$

We now introduce the nodes $0 = x_0 < x_1 < \dots < x_L = 1$ that define the elements $K_i = [x_{i-1}, x_i]$. Let $h = \max |x_i - x_{i-1}|$ be the mesh size. The multiscale finite element space

$$\mathcal{W}_h = \{v_h \in C^0(0, 1) \text{ such that } \mathcal{L}v_h = 0 \text{ on each } K_i\}, \quad (6)$$

defined using the operator \mathcal{L} , is *adapted* to the problem under study. We next proceed with a Galerkin approximation of (5) using the space \mathcal{W}_h :

$$\text{Find } u_h \in \mathcal{W}_h \text{ such that, for any } v_h \in \mathcal{W}_h, \quad a(u_h, v_h) = b(v_h).$$

The solution u_h then satisfies

$$\|u - u_h\|_E \leq \frac{h}{\pi \sqrt{\nu_{min}}} \|f\|_{L^2(0,1)} \quad (7)$$

where $\|\cdot\|_E = \sqrt{a(\cdot, \cdot)}$ is the energy norm. The proof of this estimate goes as follows. By definition of u and u_h , we have $a(u - u_h, v_h) = 0$ for any $v_h \in \mathcal{W}_h$. Hence, u_h is the orthogonal projection of u on \mathcal{W}_h according to the scalar product $a(\cdot, \cdot)$. Since $\|\cdot\|_E$ is the norm associated to that scalar product, we have

$$\|u - u_h\|_E = \inf_{v_h \in \mathcal{W}_h} \|u - v_h\|_E. \quad (8)$$

Choose v_h to be the finite element interpolant of u , which is defined by $v_h(x_i) = u(x_i)$ for any $i = 0, 1, \dots, L$, and consider the interpolation error $e = u - v_h$. On each element K_i , we have, precisely because the space \mathcal{W}_h is defined as (6),

$$\mathcal{L}e = -(\nu e')' = f \quad \text{with} \quad e(x_{i-1}) = e(x_i) = 0.$$

We multiply by e , integrate by part and obtain

$$\int_{x_{i-1}}^{x_i} \nu(x) |e'(x)|^2 dx = \int_{x_{i-1}}^{x_i} f(x) e(x) dx \leq \|f\|_{L^2(K_i)} \|e\|_{L^2(K_i)}. \quad (9)$$

Since e vanishes on the boundary of K_i , the Poincaré inequality with the best constant $(x_i - x_{i-1})/\pi$ yields

$$\|e\|_{L^2(K_i)} \leq \frac{x_i - x_{i-1}}{\pi} \|e'\|_{L^2(K_i)} \leq \frac{h}{\pi \sqrt{\nu_{min}}} \left(\int_{x_{i-1}}^{x_i} \nu(x) |e'(x)|^2 dx \right)^{1/2}.$$

By substitution in (9), we obtain

$$\int_{x_{i-1}}^{x_i} \nu(x) |e'(x)|^2 dx \leq \frac{h^2}{\pi^2 \nu_{min}} \|f\|_{L^2(K_i)}^2.$$

Summing over the elements and using (8) yields (7). Using again that ν is bounded from below, we deduce from (7) that

$$\|u - u_h\|_{H^1(0,1)} \leq \frac{h}{C_{\mathcal{D}} \pi \nu_{min}} \|f\|_{L^2(0,1)},$$

where $C_{\mathcal{D}}$ is the Poincaré constant of the domain $\mathcal{D} = (0, 1)$. As pointed out in [44, Chap. 6], the interest of the above estimate (or of estimate (7)) lies in the fact that the constant in the right-hand side only depends on ν through ν_{min} , and remains the same even if ν oscillates at a small scale. In contrast, for a standard finite element method, the error is also proportional to h , but with a constant that depends on the H^2 norm of the exact solution u . With a standard finite element space \mathcal{W}_h , we indeed classically deduce by Céa's lemma that

$$\|u - u_h\|_{H^1(0,1)} \leq \frac{\|\nu\|_{L^\infty(0,1)}}{C_{\mathcal{D}} \nu_{min}} \inf_{v_h \in \mathcal{W}_h} \|u - v_h\|_{H^1(0,1)} = \frac{\|\nu\|_{L^\infty(0,1)}}{C_{\mathcal{D}} \nu_{min}} \|u - R_h u\|_{H^1(0,1)},$$

where $C_{\mathcal{D}}$ is the Poincaré constant of the domain $\mathcal{D} = (0, 1)$, and $R_h u$ is the projection of u on \mathcal{W}_h according to the H^1 scalar product. We thus obtain that

$$\|u - u_h\|_{H^1(0,1)} \leq Ch \frac{\|\nu\|_{L^\infty(0,1)}}{\nu_{min}} \|D^2 u\|_{L^2(0,1)},$$

where C is independent from the functions ν and u . If ν oscillates at a small scale (e.g. $\nu(x) = \bar{\nu}(x/\varepsilon)$ for a fixed function $\bar{\nu}$), the H^2 norm of u may be large (of the order of ε^{-1}). A FEM approach then requires h to be smaller than ε to reach a good accuracy.

We conclude this illustration by noting that such a general analysis of the MsFEM approach is not available in dimension $d \geq 2$. The analysis presented in [38, 32], which is performed without any restriction on the dimension, additionally assumes that the matrix A^ε in (4) reads $A^\varepsilon(x) = A_{per}(x/\varepsilon)$ for a fixed *periodic* matrix A_{per} .

We now describe the MsFEM in a multidimensional setting.

Definition of the coarse mesh For simplicity (see Remark 1 below), we consider a classical \mathbb{P}_1 discretization of the domain \mathcal{D} . We denote by \mathcal{T}_h the corresponding mesh, with L nodes. Let ϕ_i^0 , $i = 1, \dots, L$, be the basis functions. We introduce the finite element space

$$\mathcal{V}_h := \text{span}(\phi_i^0, i = 1, \dots, L),$$

and define the restriction

$$\phi_i^{0,\mathbf{K}} := \phi_i^0|_{\mathbf{K}}$$

of these functions in each element \mathbf{K} .

Remark 1. We refer to [3] for a presentation of a MsFEM method that uses \mathbb{P}_2 macroscale basis functions.

Definition of the MsFEM basis Several definitions of the MsFEM basis functions have been proposed in the literature (see e.g. [37, 38, 32, 3, 30, 39]). They all follow the same pattern but they give rise to various methods. We present in the following the particular method that we have implemented. It makes use of the oversampling technique introduced in [37] and developed in [36].

For any element \mathbf{K} , we consider a domain $\mathbf{S} \supset \mathbf{K}$ (see Figure 1), obtained from \mathbf{K} by an homothetic transformation of center the centroid of \mathbf{K} , and of ratio larger than 1.




Figure 1: Definition of \mathbf{S} (in 2D for clarity)

Let $x_j^{\mathbf{S}}$ denote the coordinate of the vertex j of the domain \mathbf{S} . For any vertex i of \mathbf{S} , we introduce the affine function $\chi_i^{0,\mathbf{S}}$ (defined on \mathbf{S}) that satisfies the condition $\chi_i^{0,\mathbf{S}}(x_j^{\mathbf{S}}) = \delta_{ij}$ for all j . Let $\chi_i^{\varepsilon,\mathbf{S}} \in H^1(\mathbf{S})$ be the unique solution to the problem

$$-\text{div} \left[A^\varepsilon(x) \nabla \chi_i^{\varepsilon,\mathbf{S}}(x) \right] = 0 \quad \text{in } \mathbf{S}, \quad \chi_i^{\varepsilon,\mathbf{S}} = \chi_i^{0,\mathbf{S}} \quad \text{on } \partial\mathbf{S}, \quad (10)$$

which, in practice, is numerically solved e.g. using a finite element method with a mesh size adapted to the small scale ε . We then define the local basis functions

$$\phi_i^{\varepsilon,\mathbf{K}} = \sum_{j=1}^{d+1} \alpha_{ij} \chi_j^{\varepsilon,\mathbf{S}} \Big|_{\mathbf{K}} \quad (11)$$

as linear combinations of the restrictions of $\chi_i^{\varepsilon,\mathbf{S}}$ on \mathbf{K} , with α_{ij} chosen such that

$$\forall 1 \leq i, j \leq d+1, \quad \phi_i^{0,\mathbf{K}}(x_j^{\mathbf{K}}) = \sum_{j=1}^{d+1} \alpha_{ij} \chi_j^{0,\mathbf{S}}(x_j^{\mathbf{K}}) = \delta_{ij}, \quad (12)$$

where $x_j^{\mathbf{K}}$ denotes the coordinate of the j th vertex of the element \mathbf{K} . Note that the condition (12) is enforced on the function $\phi_i^{0,\mathbf{K}}$, and not on $\phi_i^{\varepsilon,\mathbf{K}}$. The coefficients α_{ij} are consequently independent from ε . As $\phi_i^{0,\mathbf{K}}$ and $\chi_j^{0,\mathbf{S}}|_{\mathbf{K}}$ are both affine on \mathbf{K} , condition (12) implies that

$$\forall 1 \leq i \leq d+1, \quad \forall x \in \mathbf{K}, \quad \phi_i^{0,\mathbf{K}}(x) = \sum_{j=1}^{d+1} \alpha_{ij} \chi_j^{0,\mathbf{S}}(x). \quad (13)$$

We next introduce the functions ϕ_i^ε defined on \mathcal{D} by $\phi_i^\varepsilon|_{\mathbf{K}} = \phi_i^{\varepsilon, \mathbf{K}}$ for all elements \mathbf{K} .

Note that the problems (10), indexed by \mathbf{S} , are all independent from one another. They may be solved in parallel.

Macroscale problem We now introduce the finite dimensional space

$$\mathcal{W}_h := \text{span}(\phi_i^\varepsilon, i = 1, \dots, L),$$

and proceed with the approximation

$$\text{Find } u_M \in \mathcal{W}_h \text{ such that, for any } v \in \mathcal{W}_h, \quad \mathcal{A}_\varepsilon^h(u_M, v) = b(v), \quad (14)$$

of (4), where

$$\mathcal{A}_\varepsilon^h(u, v) = \sum_{\mathbf{K}} \int_{\mathbf{K}} (\nabla v(x))^T A^\varepsilon(x) \nabla u(x) dx \quad \text{and} \quad b(v) = \int_{\mathcal{D}} f(x) v(x) dx.$$

Observe that ϕ_i^ε has jumps across the edges of the triangulation (due to the use of the oversampling technique), hence $\mathcal{W}_h \not\subset H^1(\mathcal{D})$, thus the broken integral used to define $\mathcal{A}_\varepsilon^h(u, v)$. On the other hand, since $\mathcal{W}_h \subset L^2(\mathcal{D})$, the linear form b is well defined for $v \in \mathcal{W}_h$. The formulation (14) is a *non-conforming* Galerkin approximation of (4). This brings additional error terms in the error estimation (see Lemma 12 in Section 4). On another note, remark that the dimension of \mathcal{W}_h is equal to L . The formulation (14) hence requires solving a linear system with only a limited number of degrees of freedom.

We are now in position to substantiate our claim in the introduction, where we briefly mentioned that, in the stochastic setting, a direct application of the MsFEM to approximate the solution to (2) is *unpractical*. It would indeed lead to compute, for *each realization* of $A^\varepsilon(x, \omega)$, first a basis set and second a macroscale solution. This approach has been briefly examined theoretically in [21]. It is prohibitively expensive. We therefore turn to an alternate approach.

2.2 A weakly stochastic setting

We now restrict the general setting and propose a dedicated, practical MsFEM type approach. Following up on previous works (see [5, 13, 24, 41]) and as announced in (3), we assume here that the random matrix $A^\varepsilon(x, \omega)$ in (2) is a *perturbation of a deterministic matrix*, in the sense that

$$A^\varepsilon(x, \omega) \equiv A_\eta^\varepsilon(x, \omega) = A_0^\varepsilon(x) + \eta A_1^\varepsilon(x, \omega), \quad (15)$$

where $\eta \in \mathbb{R}$ is a small deterministic parameter, A_0^ε and A_1^ε are bounded matrices, and A_0^ε is coercive, uniformly in ε . We also assume that the matrix A_η^ε itself satisfies the coercivity and boundedness assumptions, uniformly in η and ε (we refer to [6, 7, 8] and [15, 25] for other perturbative settings).

The principle of the proposed approach is to compute the MsFEM basis set of functions with the *deterministic part* A_0^ε of the matrix A_η^ε , and next to perform Monte-Carlo realizations for the macroscale problem (2)-(15), where we keep the exact matrix A_η^ε (and not only its deterministic part). Following the approach sketched in Section 2.1, we first solve (10), with $A^\varepsilon(x) \equiv A_0^\varepsilon(x)$, and build the *deterministic* finite dimensional space

$$\mathcal{W}_h := \text{span}(\phi_i^\varepsilon, i = 1, \dots, L)$$

following (11)-(12). We next proceed with a standard Galerkin approximation of (2)-(15) using \mathcal{W}_h . For each $m \in \{1, \dots, M\}$, we consider a realization $A_\eta^{\varepsilon, m}(\cdot, \omega)$ and compute $u_S^m(\cdot, \omega) \in \mathcal{W}_h$ such that

$$\forall v \in \mathcal{W}_h, \quad \sum_{\mathbf{K}} \int_{\mathbf{K}} (\nabla v(x))^T A_\eta^{\varepsilon, m}(x, \omega) \nabla u_S^m(x, \omega) dx = \int_{\mathcal{D}} f(x) v(x) dx. \quad (16)$$

Since the MsFEM basis functions are only computed once (rather than for each realization of $A_\eta^\varepsilon(x, \omega)$), a large computational gain is expected, and obtained, in comparison to the direct approach described above.

3 Numerical simulations

This section is devoted to the many numerical simulations we have performed. We first discuss some implementation details. Next, we numerically estimate the performance of our approach on various test cases, and assess its sensitivity with respect to the magnitude of η . We consider in Section 3.2 a test case in dimension one. In Section 3.3, we next study two test cases in dimension two. We also study how the presence in A_1^ε (the random component of the matrix A_η^ε) of high frequencies that are not present in the deterministic component A_0^ε , and that are thus not encoded in the highly oscillatory basis functions, affects the accuracy of our approach.

Let u_η^ε be the reference solution to (2)-(3) obtained using a finite element method with a mesh size adapted to the small scale ε , u_S be the approximation given by our approach (described in Section 2.2) and u_M be the approximation given by the direct approach (in which the MsFEM basis set is recomputed for each realization $A_\eta^{\varepsilon,m}(x, \omega)$, as explained at the end of Section 2.1). Our goal is to compare the error $u_S - u_\eta^\varepsilon$ of our numerical approximation with the error $u_M - u_\eta^\varepsilon$ of the direct and expensive approach. When η is small, we expect the approximation u_S to be essentially as accurate as the approximation u_M , and we show below that this is indeed the case.

In the sequel, we assess the accuracy using the estimators

$$e_{L^2}(u_1, u_2) = \mathbb{E} \left(\frac{\|u_1 - u_2\|_{L^2(\mathcal{D})}}{\|u_2\|_{L^2(\mathcal{D})}} \right) \quad \text{and} \quad e_{H^1}(u_1, u_2) = \mathbb{E} \left(\frac{\|u_1 - u_2\|_{H_h^1}}{\|u_2\|_{H_h^1}} \right), \quad (17)$$

where u_1 and u_2 are the solutions obtained with any two different methods, and

$$\|u\|_{H_h^1} := \left(\sum_{\mathbf{K} \in \mathcal{T}_h} \|u\|_{H^1(\mathbf{K})}^2 \right)^{1/2} \quad (18)$$

is the broken H^1 norm. The expectation is in turn computed using a Monte-Carlo method. Considering M realizations $\{X_m(\omega)\}_{1 \leq m \leq M}$ of a random variable, e.g. $X(\omega) = \frac{\|u_1(\cdot, \omega) - u_2(\cdot, \omega)\|_{H_h^1}}{\|u_2(\cdot, \omega)\|_{H_h^1}}$, we compute the empirical mean μ_M and the empirical standard deviation σ_M as

$$\mu_M(X) = \frac{1}{M} \sum_{m=1}^M X_m(\omega), \quad \sigma_M^2(X) = \frac{1}{M-1} \sum_{m=1}^M (X_m(\omega) - \mu_M(X))^2. \quad (19)$$

As a classical consequence of the Central Limit Theorem, the following estimate is commonly employed:

$$|\mathbb{E}(X) - \mu_M(X)| \leq 1.96 \frac{\sigma_M}{\sqrt{M}}.$$

It provides a practical evaluation of $\mathbb{E}(X)$ from the knowledge of $\mu_M(X)$ and $\sigma_M(X)$. The numerical parameters have been determined by an empirical study of convergence. For instance, for the reference solution, we choose the mesh size h such that the quantity $\frac{\|u_\eta^{\varepsilon,h} - u_\eta^{\varepsilon,h/2}\|_{H^1(\mathcal{D})}}{\|u_\eta^{\varepsilon,h/2}\|_{H^1(\mathcal{D})}}$ is smaller than 0.03 %, thereby formally admitting that the approximation has converged in h . The MsFEM parameters are determined likewise.

All the computations have been performed using FreeFem++ [33], with the MPI tools.

3.1 Implementation details

In the deterministic version of the MsFEM, the same matrix A^ε appears in the definition (10) of the basis functions and in the macroscale variational formulation (14). This can be used to expedite the computation of the stiffness matrix associated with (14). In our approach, described in Section 2.2, the matrix that appears in the definition of the basis functions is A_0^ε , whereas the macroscale variational problem involves $A_\eta^\varepsilon \equiv A_0^\varepsilon + \eta A_1^\varepsilon$. An additional numerical computation is thus needed.

To solve (16), we need to compute, for each element \mathbf{K} and each realization $A_\eta^{\varepsilon, \overline{m}}(x, \omega)$, the integrals

$$\mathcal{K}_{ij}^{\eta, \overline{m}}(\omega) = \int_{\mathbf{K}} \left(\nabla \phi_i^{\varepsilon, \mathbf{K}}(x) \right)^T A_\eta^{\varepsilon, \overline{m}}(x, \omega) \nabla \phi_j^{\varepsilon, \mathbf{K}}(x) dx, \quad (20)$$

where $\phi_i^{\varepsilon, \mathbf{K}}$ are deterministic functions. We recall that $A_\eta^\varepsilon(x, \omega) = A_0^\varepsilon(x) + \eta A_1^\varepsilon(x, \omega)$ (see (15)). To allow for an efficient evaluation of (20), we assume henceforth that A_1^ε is of the form

$$A_1^\varepsilon(x, \omega) = \sum_{k \in \mathbb{Z}^d} \mathbf{1}_{Q+k} \left(\frac{x}{\varepsilon} \right) X_k(\omega) B_\varepsilon^k(x), \quad (21)$$

where $Q = (0, 1)^d$, where $(X_k)_{k \in \mathbb{Z}^d}$ are scalar random variables, and for any $k \in \mathbb{Z}^d$, $x \mapsto B_\varepsilon^k(x) \in \mathbb{R}^{d \times d}$ are some deterministic functions. We comment on this assumption in Remark 2 below. The important consequence of (21) is that we can write the integral (20) as a linear combination of *deterministic integrals* over cells of size ε , with *random* coefficients. To simplify the notation, we assume that the spatial dimension is $d = 2$. We define

$$p = \left\lfloor \min \left(\frac{y_i}{\varepsilon}, \frac{y_j}{\varepsilon}, \frac{y_k}{\varepsilon} \right) \right\rfloor, \quad q = \left\lfloor \max \left(\frac{y_i}{\varepsilon}, \frac{y_j}{\varepsilon}, \frac{y_k}{\varepsilon} \right) \right\rfloor + 1,$$

and likewise, we define the integers l and m (see Fig. 2). We can then write (20) as

$$\mathcal{K}_{ij}^{\eta, \overline{m}}(\omega) = \int_{\mathbf{K}} \left(\nabla \phi_i^{\varepsilon, \mathbf{K}}(x) \right)^T A_\eta^{\varepsilon, \overline{m}}(x, \omega) \nabla \phi_j^{\varepsilon, \mathbf{K}}(x) dx = \mathcal{K}_{ij}^{0, \mathbf{K}} + \eta \sum_{\alpha=p}^{q-1} \sum_{\beta=l}^{m-1} X_{\alpha, \beta}^{\overline{m}}(\omega) \mathcal{K}_{\alpha\beta ij}^{1, \mathbf{K}}, \quad (22)$$

where

$$\mathcal{K}_{ij}^{0, \mathbf{K}} = \int_{\mathbf{K}} \left(\nabla \phi_i^{\varepsilon, \mathbf{K}}(x) \right)^T A_0^\varepsilon(x) \nabla \phi_j^{\varepsilon, \mathbf{K}}(x) dx, \quad (23)$$

$$\mathcal{K}_{\alpha\beta ij}^{1, \mathbf{K}} = \int_{\alpha\varepsilon}^{(\alpha+1)\varepsilon} \int_{\beta\varepsilon}^{(\beta+1)\varepsilon} \mathbf{1}_{\mathbf{K}}(x) \left(\nabla \phi_i^{\varepsilon, \mathbf{K}}(x) \right)^T B_\varepsilon^k(x) \nabla \phi_j^{\varepsilon, \mathbf{K}}(x) dx. \quad (24)$$




Figure 2: To practically compute the integral (20), we write that each element \mathbf{K} (here in dimension $d = 2$) is a subset of a quadrangle (here $[l\varepsilon, m\varepsilon] \times [p\varepsilon, q\varepsilon]$) composed of cells of size ε^d .

We thus compute once the deterministic integrals (23) and (24). Next, for each realization of A_η^ε , we evaluate the stiffness matrix elements $\mathcal{K}_{ij}^{\eta, \overline{m}}(\omega)$ using the right hand side of (22). No numerical quadrature is needed. As a consequence of (21), most of the work for assembling the stiffness matrix is only performed once, independently of the number of Monte Carlo realizations. This significantly contributes to the gain in term of computational cost.

Remark 2. Assumption (21) is quite general, and already covers many interesting cases in practice. As explained above, the point in (21) is that A_1^ε is a direct product (or here, a sum of direct products) of a function depending on x with a random variable that only depends on ω . Otherwise stated, $A_1^\varepsilon(x, \omega)$ depends linearly, in an explicit way, of ω . A similar assumption is made when applying reduced basis methods [45] to a problem of the type

$$\text{Find } u_\lambda \text{ such that, for any } v, a(u_\lambda, v; \lambda) = b(v), \quad (25)$$

where $a(\cdot, \cdot; \lambda)$ is a bilinear form parameterized by λ . Assume this problem has been solved for some values $\{\lambda_i\}_{i=1}^I$ of the parameter, yielding the functions $\{u_{\lambda_i}\}_{i=1}^I$. Under the assumption that $a(\cdot, \cdot; \lambda) = a_0(\cdot, \cdot) + \lambda a_1(\cdot, \cdot)$ (namely, $a(\cdot, \cdot; \lambda)$ depends linearly on λ), one can precompute the stiffness matrix elements $a_0(u_{\lambda_i}, u_{\lambda_j})$ and $a_1(u_{\lambda_i}, u_{\lambda_j})$ for any $1 \leq i, j \leq I$. This allows to next perform a very efficient Galerkin approximation of the problem (25) (for any λ) on the space $\text{Span}(u_{\lambda_i}, i = 1, \dots, I)$.

3.2 One-dimensional test-case

The purpose of this section is threefold. We first calibrate the number M of realizations considered for the Monte-Carlo method for the two-dimensional numerical experiments that we consider in the sequel. We next investigate how the accuracy of our approach depends on η and on the presence of frequencies in the random coefficient a_η^ε that are not taken into account in the MsFEM basis set functions. The low computational costs that we face in this one-dimensional situation allow us to test our approach more comprehensively than in the two-dimensional test-cases described below.

Let $(X_k)_{k \in \mathbb{Z}}$ denote a sequence of independent, identically distributed scalar random variables uniformly distributed in $[0, 1]$. We consider the random coefficient

$$a_\eta^\varepsilon(x, \omega) = \sum_{k \in \mathbb{Z}} \mathbf{1}_{(k, k+1]} \left(\frac{x}{\varepsilon} \right) \left(5 + 50 \sin^2 \left(\frac{\pi x}{\varepsilon} \right) + \eta X_k(\omega) \kappa \sin^2 \left(\frac{\zeta \pi x}{\varepsilon} \right) \right),$$

which is a particular example of the expansion (15) with

$$a_0^\varepsilon(x) = 5 + 50 \sin^2 \left(\frac{\pi x}{\varepsilon} \right) \quad \text{and} \quad a_1^\varepsilon(x, \omega) = \sum_{k \in \mathbb{Z}} \mathbf{1}_{(k, k+1]} \left(\frac{x}{\varepsilon} \right) X_k(\omega) \kappa \sin^2 \left(\frac{\zeta \pi x}{\varepsilon} \right),$$

and that satisfies the structural assumption (21). We set $\varepsilon = 0.025$ and choose κ such that the quantity

$$\mathcal{K}(\kappa, \zeta) = \left\| \frac{a_1^\varepsilon}{a_0^\varepsilon} \right\|_{L^\infty(\mathcal{D} \times \Omega)} = \text{SupEss}_{\omega \in \Omega} \left\| \frac{a_1^\varepsilon(\cdot, \omega)}{a_0^\varepsilon(\cdot, \omega)} \right\|_{L^\infty(\mathcal{D})} \quad (26)$$

has the same value $\mathcal{K} = 1$ for the three different values of $\zeta = \{1, 3, 7\}$ we consider below. We analytically compute the reference function u_η^ε , solution to

$$-\frac{d}{dx} \left(a_\eta^\varepsilon(x, \omega) \frac{du_\eta^\varepsilon}{dx}(x, \omega) \right) = 1 \quad \text{in } (0, 1), \quad u_\eta^\varepsilon(0, \omega) = u_\eta^\varepsilon(1, \omega) = 0,$$

as well as the MsFEM basis functions for both approaches. Let u_M and u_S be the approximation of u_η^ε by the two MsFEM approaches described above, where the coarse mesh size is $h = 1/30$.

We first calibrate the number of independent realizations to accurately approximate the exact expectation in (17) by the empirical mean (19). To this aim, we present on Fig. 3 the mean and the confidence interval computed using (19) for an increasing number M of realizations (we compute up to 1000 independent realizations). We check that this indicator reaches a plateau for $M \geq 30$, and thus converges fast. On this example, considering 30 realizations is hence sufficient to accurately compute the error (17). Based on this observation, we will only consider $M = 30$ realizations in the two dimensional examples of Section 3.3.

Remark 3. There is no reason to think that the calibration of our parameters that we perform in the one-dimensional situation provides an adequate adaptation of these parameters for the higher dimensional setting. We however see no other manner to proceed and the approach has indeed provided us with good results.

Note also that the MsFEM approach is much more accurate in the one-dimensional setting than in the two-dimensional setting (compare Tables 1, 2 and 3 with Tables 9 and 10 below). This is due to the specificity of the one dimensional setting. However, one-dimensional examples remain relevant for e.g. assessing how the MsFEM accuracy depends on η .




Figure 3: Convergence of the indicator $e_{H^1}(u_M, u_\eta^\varepsilon)$ (see (17)), for $\eta = 1$, $\zeta = 1$ and $\kappa = 55$. For each value of M , we plot the empirical mean along with its confidence interval, computed from the first M realizations. We only plot the results for the first 50 realizations.

We now check how the accuracy of our approach depends on η . In Tables 1, 2, 3, 4, 5 and 6, we report the estimators (17), along with their confidence intervals, for various choices of (κ, ζ) that all correspond to $\mathcal{K}(\kappa, \zeta) = 1$. For $\eta \leq 0.1$, we observe that $\|u_S - u_\eta^\varepsilon\|$ and $\|u_M - u_\eta^\varepsilon\|$ are of the same order of magnitude, and are both larger than $\|u_M - u_S\|$ (both in L^2 and broken H^1 norms). We thus obtain the same accuracy with the direct and the weak stochastic MsFEM approaches, whereas the weak stochastic MsFEM is computationally (much) less expensive. For $\eta = 1$, as expected, the accuracy of the approximation u_S deteriorates. The accuracy of u_M is independent of η .

Remark 4. In Section 4, we estimate in the H^1 (broken) norm the error between the reference solution u_η^ε and the weak MsFEM solution u_S . For information, we also include in Tables 1–6 the numerical comparison in the L^2 norm.

We now turn to a different question. In the example considered here, some frequencies present in a_1^ε do not appear in a_0^ε , and are thus not captured in the highly oscillatory basis functions ϕ_i^ε . We now show that our approach can still handle this case, provided the amplitude of these modes remains small.

We first consider the case when the amplitude κ associated to the frequency ζ is kept constant, and compare the performance of our approach in the case $\zeta = 1$ and $\zeta = 3$. In the latter case, a relevant high frequency is not taken into account in the basis set functions. Comparing Tables 1 and 4 (corresponding to $\zeta = 1$) with Tables 7 and 8 (corresponding to $\zeta = 3$) for a given value of η , we see that the accuracy of our approach deteriorates. This is not unexpected, of course. Otherwise stated, to achieve a given accuracy (say an error of 15 % in the broken H^1 norm), we need to take smaller values of η (namely $\eta \leq 0.01$) when $\zeta = 3$ than when $\zeta = 1$ (in which case $\eta = 0.1$ is already a sufficiently small value).

We now run the comparison differently. As we increase the gap between the frequency present in a_1^ε and that present in a_0^ε (i.e., as we increase ζ), we simultaneously decrease the amplitude κ of that mode. In practice, we do this by keeping constant the parameter $\mathcal{K}(\kappa, \zeta)$ defined by (26). Then the accuracy of our approach remains constant, and is independent of ζ . See indeed the numerical results of Tables 1–6, that all correspond to the choice $\mathcal{K}(\kappa, \zeta) = 1$, for three different values of ζ . We observe that, at fixed η , errors are comparable, and independent of the value of (κ, ζ) .

In conclusion, the accuracy of our approach depends *both* on the amplitude κ and the value ζ of the high frequency not taken into account in the MsFEM basis set functions. If ζ and κ are scaled so that $\mathcal{K}(\kappa, \zeta)$ remains constant (which implies that κ decreases if ζ increases), then the accuracy of our approach remains constant.

Table 1: $H^1(0, 1)$ error (17) (in %) for $\kappa = 55$ and $\zeta = 1$

η	$e_{H^1}(u_M, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_M)$
1	0.14644 ± 0.00036	2.62550 ± 0.02696	2.44359 ± 0.02696
0.1	0.16001 ± 0.00006	0.15021 ± 0.00051	0.07036 ± 0.00044
0.01	0.16258 ± 0.00000	0.10837 ± 0.00002	0.04825 ± 0.00025

Table 2: $H^1(0, 1)$ error (17) (in %) for $\kappa = 14.38$ and $\zeta = 3$

η	$e_{H^1}(u_M, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_M)$
1	0.18269 ± 0.00030	2.38950 ± 0.02277	2.23869 ± 0.02230
0.1	0.16529 ± 0.00003	0.14959 ± 0.00055	0.08082 ± 0.00041
0.01	0.16314 ± 0.00000	0.10840 ± 0.00000	0.04954 ± 0.00001

Table 3: $H^1(0, 1)$ error (17) (in %) for $\kappa = 8.39$ and $\zeta = 7$

η	$e_{H^1}(u_M, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_M)$
1	0.17436 ± 0.00026	$2,34495 \pm 0.02105$	$2,27358 \pm 0.02089$
0.1	0.16465 ± 0.00004	0.15748 ± 0.00067	0.09803 ± 0.00053
0.01	0.16308 ± 0.00000	0.10846 ± 0.00000	0.05054 ± 0.00001

Table 4: $L^2(0, 1)$ error (17) (in %) for $\kappa = 55$ and $\zeta = 1$

η	$e_{L^2}(u_M, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_M)$
1	0.00018 ± 0.00000	0.07286 ± 0.00317	0.06861 ± 0.00306
0.1	0.00018 ± 0.00000	0.00045 ± 0.00002	0.00024 ± 0.00001
0.01	0.00018 ± 0.00000	0.00015 ± 0.00000	0.00002 ± 0.00000

Table 5: $L^2(0, 1)$ error (17) (in %) for $\kappa = 14.38$ and $\zeta = 3$

η	$e_{L^2}(u_M, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_M)$
1	0.00019 ± 0.00000	0.06658 ± 0.00270	0.06238 ± 0.00261
0.1	0.00018 ± 0.00000	0.00036 ± 0.00001	0.00019 ± 0.00001
0.01	0.00018 ± 0.00000	0.00015 ± 0.00000	0.00002 ± 0.00000

Table 6: $L^2(0, 1)$ error (17) (in %) for $\kappa = 8.39$ and $\zeta = 7$

η	$e_{L^2}(u_M, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_M)$
1	0.00018 ± 0.00000	0.08903 ± 0.00310	0.08410 ± 0.00261
0.1	0.00018 ± 0.00000	0.00037 ± 0.00002	0.00016 ± 0.00000
0.01	0.00018 ± 0.00000	0.00015 ± 0.00000	0.00003 ± 0.00000

Table 7: $H^1(0, 1)$ error (17) (in %) for $\kappa = 55$ and $\zeta = 3$

η	$e_{H^1}(u_M, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_M)$
1	0.21826 ± 0.00073	12.30047 ± 0.10647	12.01694 ± 0.10617
0.1	0.17142 ± 0.00013	0.59293 ± 0.00519	0.49523 ± 0.00489
0.01	0.16383 ± 0.00001	0.11448 ± 0.00014	0.05247 ± 0.00007

Table 8: $L^2(0, 1)$ error (17) (in %) for $\kappa = 55$ and $\zeta = 3$

η	$e_{L^2}(u_M, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_M)$
1	0.00022 ± 0.00000	1.53780 ± 0.03878	1.51837 ± 0.00385
0.1	0.00019 ± 0.00000	0.00503 ± 0.00027	0.00406 ± 0.00024
0.01	0.00018 ± 0.00000	0.00018 ± 0.00000	0.00005 ± 0.00000

3.3 Two-dimensional test-cases

We now test our approach on two-dimensional test cases. Using the first test case, we show, similarly to the one-dimensional situation, that the weak stochastic MsFEM yields accurate results, provided the parameter η is sufficiently small, and provided that the amplitude associated to frequencies present in A_η^ε but not encoded in the deterministic basis functions is small (see Section 3.3.1). Next, in Section 3.3.2, we consider a test case similar to a classical benchmark test case of the literature. We again observe that our approach is efficient. For both cases, we show that the parameter η does not need to be extremely small for our approach to be highly competitive.

3.3.1 A multi-frequency case

In line with what we observed in the one-dimensional case, we show here that the weak stochastic MsFEM provides interesting results even in the case when not all the frequencies present in A_η^ε are captured in the deterministic basis functions, provided their amplitude is not too large. To this aim, we consider the following numerical example.

Let $(X_{k,l})_{(k,l) \in \mathbb{Z}^2}$ denote a sequence of independent, identically distributed scalar random variables uniformly distributed in the interval $[0, 1]$. We consider the random matrix

$$A_\eta^\varepsilon(x, y, \omega) = a_0^\varepsilon(x, y) \text{Id}_2 + \eta a_1^\varepsilon(x, y, \omega) \text{Id}_2,$$

with

$$\begin{aligned} a_0^\varepsilon(x, y) &= 5 + 50 \sin^2\left(\frac{\pi x}{\varepsilon}\right) \sin^2\left(\frac{\pi y}{\varepsilon}\right), \\ a_1^\varepsilon(x, y, \omega) &= \sum_{(k,l) \in \mathbb{Z}^2} \mathbf{1}_{(k,k+1)}\left(\frac{x}{\varepsilon}\right) \mathbf{1}_{(l,l+1)}\left(\frac{y}{\varepsilon}\right) \left(X_{k,l}(\omega) \kappa \sin^2\left(\frac{\zeta \pi x}{\varepsilon}\right) \sin^2\left(\frac{\zeta \pi y}{\varepsilon}\right) \right). \end{aligned}$$

Again, this choice is a particular example of the expansion (15) satisfying the structural assumption (21). We consider two different values of ζ , namely $\zeta = 1$ and $\zeta = 3$. As in the previous test case, the frequency ζ is not present in the deterministic part of A_η^ε , and thus not encoded in the basis functions. In line with what we observed in Section 3.2, we choose the amplitude κ associated to that frequency such that the quantity (26) has the same value $\mathcal{K} = 1$ for both values of ζ . We compute u_η^ε solution to

$$-\text{div} [A_\eta^\varepsilon(\cdot, \omega) \nabla u_\eta^\varepsilon(\cdot, \omega)] = 1 \quad \text{in } \mathcal{D}, \quad u_\eta^\varepsilon(\cdot, \omega) = 0 \quad \text{on } \partial\mathcal{D},$$

on the domain $\mathcal{D} = (0, 1)^2$ with $\varepsilon = 0.025$. Let u_M and u_S be its approximation by the two MsFEM approaches described above. The numerical parameters for the computation are again determined using an empirical study of convergence. We use for the reference solution u_η^ε a fine mesh of size $h_f = \varepsilon/40$. The MsFEM basis functions are computed in each element \mathbf{K} using a mesh of size $h_M = \varepsilon/80$. The oversampling parameter (i.e. the scale ratio of the homothetic transformation between \mathbf{K} and \mathbf{S} , see Fig. 1) is equal to 3. The coarse mesh size is $h = 1/30$. In view of the results of Section 3.2, we consider $M = 30$ independent realizations, which will prove to again be sufficient to obtain accurate results.

In Tables 9 and 10 (Tables 11 and 12 respectively), we report the estimator (17), along with its confidence interval, for the broken $H^1(\mathcal{D})$ norm and for the $L^2(\mathcal{D})$ norm, respectively. The results obtained here confirm our observations in the one-dimensional setting (Section 3.2):

- for given ζ and κ , we observe that, when η is sufficiently small (here, $\eta \leq 0.1$), the alternative approach provides a solution u_S that is an approximation of u_η^ε as accurate as u_M , for a much smaller computational cost (as the MsFEM basis set has only been computed once rather than for each independent realization of A_η^ε).
- our approach yields accurate results even if the frequency ζ is not encoded in the basis functions ϕ_i^ε , provided the associated amplitude κ is scaled accordingly. Figures in Table 9 (respectively Table 11) are very close to those of Table 10 (respectively Table 12). This confirms that the error made by the weak stochastic MsFEM seems to be independent of κ and ζ , provided these two parameters are scaled so that $\mathcal{K}(\kappa, \zeta)$ remains constant. If ζ becomes different than 1, the frequency present in a_0^ε , then the amplitude κ associated to the frequency ζ has to decrease to keep $\mathcal{K}(\kappa, \zeta)$ (and thus the accuracy of u_S) constant.

These observations again demonstrate the efficiency of the approach.

Table 9: $H^1(\mathcal{D})$ error (17) (in %) for $\kappa = 73.61$ and $\zeta = 1$

η	$e_{H^1}(u_M, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_M)$
1	7.8437 ± 0.1350	19.8818 ± 0.4123	18.8662 ± 0.4216
0.1	6.8053 ± 0.0165	7.3868 ± 0.0276	3.1528 ± 0.0517
0.01	6.7338 ± 0.0017	6.9795 ± 0.0016	1.8763 ± 0.0013

Table 10: $H^1(\mathcal{D})$ error (17) (in %) for $\kappa = 10$ and $\zeta = 3$

η	$e_{H^1}(u_M, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_M)$
1	6.7224 ± 0.0368	12.7292 ± 0.2172	10.8128 ± 0.2442
0.1	6.7154 ± 0.0044	7.1069 ± 0.0128	2.2925 ± 0.0206
0.01	6.1725 ± 0.0004	6.9770 ± 0.0010	1.8504 ± 0.0003

Table 11: $L^2(\mathcal{D})$ error (17) (in %) for $\kappa = 73.61$ and $\zeta = 1$

η	$e_{L^2}(u_M, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_M)$
1	1.4355 ± 0.0795	4.1649 ± 0.1652	2.8468 ± 0.1694
0.1	1.0630 ± 0.0108	1.1369 ± 0.0075	0.1441 ± 0.0354
0.01	1.0211 ± 0.0011	1.1512 ± 0.0007	0.1351 ± 0.0014

Table 12: $L^2(\mathcal{D})$ error (17) (in %) for $\kappa = 10$ and $\zeta = 3$

η	$e_{L^2}(u_M, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_M)$
1	1.0744 ± 0.0127	1.8433 ± 0.0582	0.8426 ± 0.0832
0.1	1.0226 ± 0.0015	1.1249 ± 0.0038	0.1147 ± 0.0073
0.01	1.0170 ± 0.0001	1.1551 ± 0.0004	0.1427 ± 0.0003

Remark 5. In Tables 9-12, we observe that the size of the confidence interval is much smaller than the distance between two different errors. This a posteriori validates the choice of the number M of Monte Carlo realizations according to the calibration we performed in the one-dimensional setting. In the two-dimensional setting studied here, we observe that considering $M = 30$ realizations is again sufficient. The same conclusion holds for results presented in Tables 13-16 below.

3.3.2 A classical test case

We consider in this section a test case similar to a classical test case of the literature (see e.g. [37, 39, 19, 32]). Let $(X_{k,l})_{(k,l) \in \mathbb{Z}^2}$ denote a sequence of independent, identically distributed scalar random variables uniformly distributed in the interval $[0, 1]$. We consider the random matrix

$$A_\eta^\varepsilon(x, y, \omega) = \sum_{(k,l) \in \mathbb{Z}^2} \mathbf{1}_{(k,k+1]} \left(\frac{x}{\varepsilon} \right) \mathbf{1}_{(l,l+1]} \left(\frac{y}{\varepsilon} \right) \left(\frac{2 + P \sin(2\pi x/\varepsilon)}{2 + P \sin(2\pi y/\varepsilon)} + \frac{2 + \sin(2\pi y/\varepsilon)}{2 + P \sin(2\pi x/\varepsilon)} \right) (1 + \eta X_{k,l}(\omega)) \text{Id}_2,$$

with $P = 1.8$ and $\varepsilon = 0.025$. We compute the reference solution u_η^ε and its two approximations u_M and u_S with the same numerical parameters as in Section 3.3.1.

In Tables 13 and 14, we report the estimator (17), along with its confidence interval, for the broken $H^1(\mathcal{D})$ norm and for the $L^2(\mathcal{D})$ norm, respectively. We again see that, when η is sufficiently small, u_S is an approximation of the reference solution u_η^ε as accurate as u_M . In Tables 15 and 16, we report on the accuracy of u_S , for more values of η . Assuming that the accuracy of u_M does not depend on η (which is consistent with the results reported in Tables 13 and 14), we see that our approach is as accurate as the direct, expensive MsFEM approach, as soon as $\eta \leq 0.1$ (if we use the broken H^1 norm to assess accuracy) and $\eta \leq 0.25$ (if we rather use the L^2 norm). The parameter η hence does not need to be extremely small for our approach to be highly competitive.

Table 13: $H^1(\mathcal{D})$ error (17) (in %)

η	$e_{H^1}(u_M, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_\eta^\varepsilon)$	$e_{H^1}(u_S, u_M)$
1	8.1154 ± 0.1913	17.3678 ± 0.7784	15.5113 ± 0.8689
0.1	7.1664 ± 0.0199	7.0524 ± 0.0705	2.5638 ± 0.1006
0.01	7.1453 ± 0.0020	7.2837 ± 0.0067	1.3882 ± 0.0020

Table 14: $L^2(\mathcal{D})$ error (17) (in %)

η	$e_{L^2}(u_M, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_\eta^\varepsilon)$	$e_{L^2}(u_S, u_M)$
1	0.5620 ± 0.0803	1.6855 ± 0.4860	1.4739 ± 0.5048
0.1	0.5354 ± 0.0160	0.5688 ± 0.0630	0.1984 ± 0.0712
0.01	0.5347 ± 0.0012	0.6192 ± 0.0054	0.1072 ± 0.0032

Table 15: $H^1(\mathcal{D})$ error (17) (in %)

η	$e_{H^1}(u_S, u_\eta^\varepsilon)$
1	17.3678 ± 0.7784
0.5	15.9578 ± 0.3461
0.25	10.6130 ± 0.1591
0.1	7.0524 ± 0.0705
0.01	7.2837 ± 0.0067

Table 16: $L^2(\mathcal{D})$ error (17) (in %)

η	$e_{L^2}(u_S, u_\eta^\varepsilon)$
1	1.6855 ± 0.4860
0.5	1.0246 ± 0.4414
0.25	0.5291 ± 0.2285
0.1	0.5688 ± 0.0630
0.01	0.6192 ± 0.0054

4 Analysis

This section is devoted to the analysis of the approach introduced in Section 2.2, and to the derivation of error bounds. As is often the case for the MsFEM (see e.g. [32]), we perform the analysis in a setting where the problem (2)-(3) that we consider admits a homogenized limit as ε vanishes (although, we repeat it, the approach is used in practice for more general cases). The structure of our proof is similar to that for the deterministic setting, which we now overview (we refer to [32] for all the details).

In the case when the oversampling technique is not used, the MsFEM is a conforming Galerkin approximation, and the error is estimated using the Céa lemma:

$$\|u^\varepsilon - u_M\|_{H^1} \leq C \inf_{v_h \in \mathcal{W}_h} \|u^\varepsilon - v_h\|_{H^1},$$

where u^ε is the solution to the reference deterministic highly oscillatory problem (1), u_M is the MsFEM solution, and the constant C is independent from ε and h . In the case when the oversampling technique is used, the MsFEM is a non-conforming Galerkin method. The error is then bounded from above by the sum of the best approximation error (the right-hand side of the above estimate) and the non-conforming error (that we do not detail here):

$$\|u^\varepsilon - u_M\|_{H^1} \leq C \left[\inf_{v_h \in \mathcal{W}_h} \|u^\varepsilon - v_h\|_{H^1} + \text{non-conforming error} \right].$$

Note that, in the non-conforming case, the MsFEM solution u_M does not belong to H^1 , and one should write the above estimate with a broken H^1 norm rather than the H^1 norm. For the sake of clarity, we ignore this distinction in this preliminary discussion.

Taking advantage of the homogenization setting, we introduce the two-scale expansion

$$v^\varepsilon = u^* + \varepsilon \sum_{i=1}^d w_{e_i}^0 \left(\frac{\cdot}{\varepsilon} \right) \partial_i u^*$$

of u^ε , where u^* is the homogenized solution, $w_{e_i}^0$ is the periodic corrector associated to $e_i \in \mathbb{R}^d$, and $\partial_i u^*$ denotes the partial derivative $\frac{\partial u^*}{\partial x_i}$. We next write

$$\|u^\varepsilon - u_M\|_{H^1} \leq C \left[\|u^\varepsilon - v^\varepsilon\|_{H^1} + \inf_{v_h \in \mathcal{W}_h} \|v^\varepsilon - v_h\|_{H^1} + \text{non-conforming error} \right].$$

The first term in the right-hand side is estimated using standard homogenization results. To estimate the second term, one considers a suitably chosen element $v_h \in \mathcal{W}_h$, for which $\|v^\varepsilon - v_h\|_{H^1}$ can be estimated directly. The main idea is that the highly oscillating part of v^ε can be well approached by an element in \mathcal{W}_h , since, by construction, the highly oscillatory basis functions are defined by a problem similar to the corrector problem, and thus encode the same highly oscillatory behavior as that present in the correctors $w_{e_i}^0$. We are thus left with approximating the slowly varying components of v^ε , for which standard FEM estimates are used. Lastly, we again use the fact that our problem admits a homogenized limit to estimate the third term, i.e. the non-conforming error.

In the sequel, we follow the same strategy in our stochastic setting. We hence first write (see (64) below) that

$$\|u_\eta^\varepsilon(\cdot, \omega) - u_S(\cdot, \omega)\|_{H^1} \leq C \left[\inf_{v_h \in \mathcal{W}_h} \|u_\eta^\varepsilon(\cdot, \omega) - v_h(\cdot, \omega)\|_{H^1} + \text{non-conforming error} \right], \quad (27)$$

where u_η^ε is the solution to the reference stochastic problem (2)-(3) and C is a deterministic constant independent from ε , h and η (note that, in (64), we use a *broken* H^1 norm rather than the H^1 norm; as pointed out above, this is due to the fact that our approach is a *non-conforming* Galerkin approximation; we ignore this distinction in the current discussion). To estimate the best approximation error (the first term in the right-hand side of (27) above), we use the triangle inequality, and write (see (84) below) that

$$\inf_{v_h \in \mathcal{W}_h} \|u_\eta^\varepsilon(\cdot, \omega) - v_h(\cdot, \omega)\|_{H^1} \leq \|u_\eta^\varepsilon(\cdot, \omega) - v_\eta^\varepsilon(\cdot, \omega)\|_{H^1} + \inf_{v_h \in \mathcal{W}_h} \|v_\eta^\varepsilon(\cdot, \omega) - v_h(\cdot, \omega)\|_{H^1}, \quad (28)$$

where v_η^ε is the two-scale expansion of the solution u_η^ε truncated at order ε^2 . A first difficulty owes to the fact that, in the general stochastic setting, no estimate is known on $\|u_\eta^\varepsilon(\cdot, \omega) - v_\eta^\varepsilon(\cdot, \omega)\|_{H^1}$. One only knows that its expectation vanishes when $\varepsilon \rightarrow 0$. However, in the present article, we consider a *weakly* stochastic case. In that setting, we have derived such a convergence rate type result in [42], and we can thus bound the first term of (28) (see Section 4.1.2 below for more details). The second term, $\inf_{v_h \in \mathcal{W}_h} \|v_\eta^\varepsilon(\cdot, \omega) - v_h\|_{H^1}$, of (28), is estimated using an explicit construction of a suitable v_h (see (85)), similarly to the deterministic setting. We again use there our specific weakly stochastic setting. Lastly, the non-conforming error (the second term in the right-hand side of (27) above) is estimated following arguments similar to those of the deterministic case, using that our problem admits a homogenized limit and is weakly stochastic.

This section is organized as follows. The error estimation is presented in Section 4.1. We first recall in Section 4.1.1 the formulation of the homogenized problem, and some results specific to the weakly stochastic case. Next, in Section 4.1.2, we establish an error bound between the reference solution u_η^ε and its two-scale expansion v_η^ε (see Theorem 7), which allows to bound the first term in the right-hand side of (28). Our main result, Theorem 10, is given in Section 4.1.3, and proved in Section 4.2. The proof essentially consists in explicitly building a function $v_h \in \mathcal{W}_h$ such that the second term of (28) can be directly estimated. It also makes use of several technical results (Lemmas 13, 14 and 16 below) to bound the non-conforming error, i.e. the second term in the right hand side of (27). The proof of these technical results is postponed until Appendix A. Last, in Section 4.3, we specifically consider the one dimensional case.

Before proceeding further, we recall the setting of stochastic homogenization we work with. The reader familiar with this theory may directly proceed to Section 4.1. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space. For a random variable $X \in L^1(\Omega, d\mathbb{P})$, we denote by $\mathbb{E}(X) = \int_\Omega X(\omega) d\mathbb{P}(\omega)$ its expectation value. We assume that the group $(\mathbb{Z}^d, +)$ acts on Ω . We denote by $(\tau_k)_{k \in \mathbb{Z}^d}$ this action, and assume that it preserves the measure \mathbb{P} , i.e.

$$\forall k \in \mathbb{Z}^d, \quad \forall A \in \mathcal{F}, \quad \mathbb{P}(\tau_k A) = \mathbb{P}(A).$$

We assume that τ is *ergodic*, that is,

$$\forall A \in \mathcal{F}, \quad (\forall k \in \mathbb{Z}^d, \quad \tau_k A = A) \Rightarrow (\mathbb{P}(A) = 0 \quad \text{or} \quad 1).$$

We define the following notion of stationarity: any $F \in L^1_{\text{loc}}(\mathbb{R}^d, L^1(\Omega))$ is said to be *stationary* if

$$\forall k \in \mathbb{Z}^d, \quad F(x+k, \omega) = F(x, \tau_k \omega) \text{ almost everywhere, almost surely.} \quad (29)$$

Note that we have chosen to present the theory in a *discrete* stationary setting, which is more appropriate for our specific purpose, which is to consider a setting close to *periodic* homogenization. Random homogenization is more often presented in the *continuous* stationary setting. This is only a matter of small modifications. We refer to the bibliography for the latter.

For the sake of analysis, we assume in this section that the matrix $A_\eta^\varepsilon(x, \omega)$ in (2)-(3) reads $A_\eta^\varepsilon(x, \omega) = A_\eta\left(\frac{x}{\varepsilon}, \omega\right)$, where the random matrix A_η is stationary in the sense of (29). The problem (2) now reads

$$-\operatorname{div} \left[A_\eta \left(\frac{\cdot}{\varepsilon}, \omega \right) \nabla u_\eta^\varepsilon(\cdot, \omega) \right] = f \text{ in } \mathcal{D}, \quad u_\eta^\varepsilon(\cdot, \omega) = 0 \text{ on } \partial\mathcal{D}, \quad (30)$$

where $A_\eta(\cdot, \omega) \in (L^\infty(\mathbb{R}^d))^{d \times d}$ satisfies the standard coercivity and boundedness conditions: there exists two constants $a_+ \geq a_- > 0$ such that

$$\forall \eta, \forall \xi \in \mathbb{R}^d, \quad a_- |\xi|^2 \leq A_\eta(x, \omega) \xi \cdot \xi \text{ a.e. on } \mathbb{R}^d, \text{ a.s. and } \|A_\eta(\cdot, \omega)\|_{L^\infty(\mathbb{R}^d)} \leq a_+ \text{ a.s.} \quad (31)$$

Due to the stationarity assumption on A_η , the problem (30) admits a homogenized limit when $\varepsilon \rightarrow 0$. Note that, to the best of our knowledge, all analyses of the MsFEM approach in the deterministic setting that have been proposed in the literature are performed under a similar assumption (the matrix A^ε in (1) is assumed to read $A^\varepsilon(x) = A_{\text{per}}\left(\frac{x}{\varepsilon}\right)$ for a fixed periodic matrix A_{per} , see e.g. [38, 32]).

In addition, in line with (3) and (15), we assume that A_η is of the form

$$A_\eta(x, \omega) = A_{\text{per}}(x) + \eta A_1(x, \omega), \quad (32)$$

where $\eta \in \mathbb{R}$ is small parameter (we henceforth assume that $|\eta| \leq 1$), A_{per} is a symmetric bounded Q -periodic matrix ($Q = [0, 1]^d$) satisfying the ellipticity condition almost everywhere on \mathbb{R}^d , and A_1 is a symmetric bounded stationary matrix: $|A_1(x, \omega)| \leq C$ almost everywhere in \mathbb{R}^d , almost surely. Since η is small, our problem is *weakly* stochastic.

In line with (21), we furthermore assume that A_1 is of the form

$$A_1(x, \omega) = \sum_{k \in \mathbb{Z}^d} \mathbf{1}_{Q+k}(x) X_k(\omega) B_{\text{per}}(x), \quad (33)$$

where $(X_k(\omega))_{k \in \mathbb{Z}^d}$ is a sequence of i.i.d. scalar random variables such that

$$\exists C, \forall k \in \mathbb{Z}^d, \quad |X_k(\omega)| \leq C \text{ almost surely,}$$

and $B_{\text{per}} \in (L^\infty(\mathbb{R}^d))^{d \times d}$ is a Q -periodic matrix. Besides being used in Theorem 7 below, this assumption is also used in the proof of Lemma 16, to recognize that some quantity (namely, (126) below) is a normalized sum of *i.i.d. variables*, on which we can use Central Limit Theorem arguments. As mentioned in Section 3.1 above, the form (33) is not essential. The point in (33) is that A_1 is a sum of *direct products* of a function depending on x with a random variable only depending on ω . Assumptions alternative to (33) could be made, that still satisfy this framework.

Finally, we assume that

$$A_{\text{per}} \text{ is Hölder continuous,} \quad (34)$$

$$B_{\text{per}} \text{ is Hölder continuous.} \quad (35)$$

We use these assumptions to obtain a rate of convergence of the two-scale expansion of u_η^ε (see [42] and Theorem 7 below), and hence control the first term in the right-hand side of (28). Such assumptions are standard when proving convergence rates of two-scale expansions (see e.g. [40, p. 28]). In turn, to control the second term in (28) and the non-conforming error (the second term in (27)), we do not need B_{per} to be Hölder continuous, and only use the fact that A_{per} is Hölder continuous (to obtain e.g. Lemmas 9, 13, 14 and 17). The numerical examples that we have considered in Section 3 satisfy assumptions (34)-(35) (remark that assumption (34) is also satisfied in the numerical examples considered in e.g. [30]).

Note that we have assumed A_{per} and B_{per} to be symmetric only for the sake of simplicity. The arguments used below carry over to the non-symmetric case up to slight modifications.

4.1 Error estimation

To bound the error between the reference solution u_η^ε and the MsFEM solution u_S , we use in many instances that we work in a *weakly* stochastic homogenization setting. We first recall in Section 4.1.1 some results specific to weakly stochastic homogenization. This setting also allows to state rates of convergence for the two-scale expansion of u_η^ε , as we explain in Section 4.1.2. Our main result, Theorem 10, is given in Section 4.1.3.

4.1.1 The homogenized equation

Under the conditions recalled above, it is known (see e.g. [12, 40]) that the solution $u_\eta^\varepsilon(\cdot, \omega)$ to (30) a.s. converges weakly in $H_0^1(\mathcal{D})$ as $\varepsilon \rightarrow 0$ to the deterministic solution u_η^* of the homogenized equation

$$-\operatorname{div} [A_\eta^* \nabla u_\eta^*] = f \quad \text{in } \mathcal{D}, \quad u_\eta^* = 0 \quad \text{on } \partial\mathcal{D}. \quad (36)$$

The homogenized matrix is given by

$$(A_\eta^*)_{ij} = \mathbb{E} \left(\int_Q (e_i + \nabla w_{e_i}^\eta(y, \cdot))^T A_\eta(y, \cdot) (e_j + \nabla w_{e_j}^\eta(y, \cdot)) dy \right), \quad (37)$$

where, for any $p \in \mathbb{R}^d$, w_p^η is the unique (up to the addition of a random constant) solution to the corrector problem

$$\begin{cases} -\operatorname{div} [A_\eta(\cdot, \omega) (p + \nabla w_p^\eta(\cdot, \omega))] = 0 & \text{in } \mathbb{R}^d, \\ \nabla w_p^\eta \text{ is stationary in the sense of (29),} \\ \mathbb{E} \left(\int_Q \nabla w_p^\eta(y, \cdot) dy \right) = 0. \end{cases} \quad (38)$$

The variational problem associated with (36) writes: find $u_\eta^* \in H_0^1(\mathcal{D})$ such that

$$\forall v \in H_0^1(\mathcal{D}), \quad \mathcal{A}_\eta^*(u_\eta^*, v) = b(v),$$

where

$$\mathcal{A}_\eta^*(u, v) = \int_{\mathcal{D}} (\nabla v(x))^T A_\eta^* \nabla u(x) dx \quad \text{and} \quad b(v) = \int_{\mathcal{D}} f(x)v(x) dx. \quad (39)$$

As shown in [13, 24], in the weakly stochastic setting, the homogenized matrix A_η^* can be expanded in terms of a series in powers of η :

$$A_\eta^* = A_{per}^* + \eta A_1^* + \eta^2 A_2^*(\eta), \quad (40)$$

where $A_2^*(\eta)$ is a deterministic matrix, that depends on η and is bounded as $\eta \rightarrow 0$, and where, for any $1 \leq i, j \leq d$,

$$(A_{per}^*)_{ij} = \int_Q (e_i + \nabla w_{e_i}^0)^T A_{per} (e_j + \nabla w_{e_j}^0), \quad (41)$$

$$(A_1^*)_{ij} = \int_Q (e_i + \nabla w_{e_i}^0)^T \mathbb{E}(A_1) (e_j + \nabla w_{e_j}^0), \quad (42)$$

where, for any $p \in \mathbb{R}^d$, w_p^0 is the unique (up to the addition of a constant) solution to the deterministic corrector problem associated to the periodic matrix A_{per} :

$$\begin{cases} -\operatorname{div} [A_{per} (p + \nabla w_p^0)] = 0, \\ w_p^0 \text{ is } Q\text{-periodic.} \end{cases} \quad (43)$$

Under the assumption (33), we have $A_1^* = \mathbb{E}(X_0) \bar{B}$, with

$$\forall 1 \leq i, j \leq d, \quad \bar{B}_{ij} = \int_Q (e_i + \nabla w_{e_i}^0)^T B_{per} (e_j + \nabla w_{e_j}^0). \quad (44)$$

Remark 6. In general, when A_{per} is not symmetric, the expression of A_1^* includes additional terms. Indeed, writing $\nabla w_p^\eta = \nabla w_p^0 + \eta \nabla w_p^1 + O(\eta^2)$, we in general need $\mathbb{E}(\nabla w_p^1)$ to compute A_1^* (see e.g. [24, 13]). In the symmetric case, these additional terms vanish, see e.g. [4, Remark 4.2 p. 117]. In the non-symmetric case, the expression (42) of A_1^* needs to be slightly modified, but the expansion (40) remains true. Our arguments hence carry over to the non-symmetric case.

Using the expansion (40) of A_η^\star with respect to η , it is easy to see that the solution u_η^\star to (36) can also be expanded in a series in powers of η . We have

$$u_\eta^\star = u_0^\star + \eta \mathbb{E}(X_0) \bar{u}_1^\star + \eta^2 r_\eta \quad \text{with} \quad \|r_\eta\|_{H^1(\mathcal{D})} \leq C, \quad (45)$$

where C is a constant independent of η , and where u_0^\star solves

$$-\operatorname{div} [A_{per}^\star \nabla u_0^\star] = f \quad \text{in } \mathcal{D}, \quad u_0^\star = 0 \quad \text{on } \partial\mathcal{D}, \quad (46)$$

and \bar{u}_1^\star solves

$$-\operatorname{div} [A_{per}^\star \nabla \bar{u}_1^\star] = \operatorname{div} [\bar{B} \nabla u_0^\star] \quad \text{in } \mathcal{D}, \quad \bar{u}_1^\star = 0 \quad \text{on } \partial\mathcal{D}. \quad (47)$$

The expansion (45) will be useful in the sequel. We will also need a bound on u_η^\star and r_η in the H^2 norm. Recall that u_η^\star is the solution to (36), whereas r_η is solution to

$$-\operatorname{div} [A_\eta^\star \nabla r_\eta] = \operatorname{div} [A_2^\star(\eta) (\nabla u_0^\star + \eta \mathbb{E}(X_0) \nabla \bar{u}_1^\star) + \mathbb{E}(X_0) A_1^\star \nabla \bar{u}_1^\star] \quad \text{in } \mathcal{D}, \quad r_\eta = 0 \quad \text{on } \partial\mathcal{D}. \quad (48)$$

In view of (31), we have, almost surely and almost everywhere, $a_- \operatorname{Id} \leq A_\eta \leq a_+ \operatorname{Id}$ in the sense of symmetric matrices. Recalling that homogenization preserves the order of symmetric matrices (see e.g. [48, page 12]), we deduce that

$$\forall \eta, \quad \forall \xi \in \mathbb{R}^d, \quad a_- |\xi|^2 \leq A_\eta^\star \xi \cdot \xi \leq a_+ |\xi|^2.$$

In addition, the right-hand sides of (36) and (48) are bounded uniformly in η in the L^2 norm. Using [34, Theorems 9.15 and 9.14], we obtain that there exists C such that

$$\forall \eta, \quad \|u_\eta^\star\|_{H^2(\mathcal{D})} \leq C \quad \text{and} \quad \|r_\eta\|_{H^2(\mathcal{D})} \leq C. \quad (49)$$

4.1.2 Two scale expansion of the reference solution u_η^ε

As recalled above, the standard error analysis for the MsFEM in the deterministic setting is performed in the case when the matrix A^ε in (1) reads $A^\varepsilon(x) \equiv A_{per}(x/\varepsilon)$ for a fixed periodic matrix A_{per} . The problem (1) then admits a homogenized limit. To obtain bounds on the MsFEM error, one step of the proof is to approximate the oscillatory solution u^ε by its two-scale expansion $u^\star + \varepsilon \sum_{i=1}^d w_{e_i}^0 \left(\frac{\cdot}{\varepsilon}\right) \partial_i u^\star$, where u^\star is the homogenized solution, w_p^0 is the periodic corrector associated to $p \in \mathbb{R}^d$, and $\partial_i u^\star = \frac{\partial u^\star}{\partial x_i}$. In the deterministic case, it is known (see e.g. [12, 23, 40]) that, under some regularity assumptions on A_{per} and u^\star ,

$$\left\| u^\varepsilon - \left[u^\star + \varepsilon \sum_{i=1}^d w_{e_i}^0 \left(\frac{\cdot}{\varepsilon}\right) \partial_i u^\star \right] \right\|_{H^1(\mathcal{D})} \leq C \sqrt{\varepsilon} \quad (50)$$

for a constant C independent of ε .

In the stochastic case, it is known that $\mathbb{E} \left[\left\| u^\varepsilon - \left[u^\star + \varepsilon \sum_{i=1}^d w_{e_i} \left(\frac{\cdot}{\varepsilon}, \omega\right) \partial_i u^\star \right] \right\|_{H^1(\mathcal{D})}^2 \right]$ converges to 0 as $\varepsilon \rightarrow 0$ (see [47, Theorem 3]), but no rate of convergence is known (except in some one-dimensional situations, see e.g. [10, 16, 43]). However, in the present article, and as announced above, we consider a *weakly* stochastic case. In this setting, we have derived in [42] a result similar to (50). We now state this result, which will be useful for our analysis.

Theorem 7 (from [42], Theorem 2). *Assume $d > 1$. Let v_η^ε be the solution to (30), and assume that A_η satisfies (32)-(33)-(34)-(35). Let A_{per}^\star , w_p^0 , u_0^\star and \bar{u}_1^\star be defined by (41), (43), (46) and (47). The two-scale expansion v_η^ε of u_η^ε reads*

$$\begin{aligned} v_\eta^\varepsilon(\cdot, \omega) = & u_0^\star + \eta \mathbb{E}(X_0) \bar{u}_1^\star + \varepsilon \sum_{i=1}^d \left[w_{e_i}^0 \left(\frac{\cdot}{\varepsilon}\right) (\partial_i u_0^\star + \eta \mathbb{E}(X_0) \partial_i \bar{u}_1^\star) \right. \\ & \left. + \eta \mathbb{E}(X_0) \psi_{e_i} \left(\frac{\cdot}{\varepsilon}\right) \partial_i u_0^\star + \eta \sum_{k \in I_\varepsilon} (X_k(\omega) - \mathbb{E}(X_0)) \chi_{e_i} \left(\frac{\cdot}{\varepsilon} - k\right) \partial_i u_0^\star \right], \quad (51) \end{aligned}$$

where

$$I_\varepsilon = \{k \in \mathbb{Z}^d \text{ such that } \varepsilon(Q+k) \cap \mathcal{D} \neq \emptyset\},$$

and where, for any $p \in \mathbb{R}^d$, ψ_p is the solution (unique up to the addition of a constant) to

$$\begin{cases} -\operatorname{div}[A_{\text{per}} \nabla \psi_p] = \operatorname{div}[B_{\text{per}}(p + \nabla w_p^0)], \\ \psi_p \text{ is } Q\text{-periodic,} \end{cases} \quad (52)$$

and χ_p is the unique solution to

$$\begin{cases} -\operatorname{div}[A_{\text{per}} \nabla \chi_p] = \operatorname{div}[\mathbf{1}_Q B_{\text{per}}(p + \nabla w_p^0)] & \text{in } \mathbb{R}^d, \\ \chi_p \in L^2_{\text{loc}}(\mathbb{R}^d), \quad \nabla \chi_p \in (L^2(\mathbb{R}^d))^d, \\ \lim_{|x| \rightarrow \infty} \chi_p(x) = 0. \end{cases} \quad (53)$$

We assume that $u_0^* \in W^{2,\infty}(\mathcal{D})$ and $\bar{u}_1^* \in W^{2,\infty}(\mathcal{D})$. Then

$$\sqrt{\mathbb{E} \left[\|u_\eta^\varepsilon - v_\eta^\varepsilon\|_{H^1(\mathcal{D})}^2 \right]} \leq C \left(\sqrt{\varepsilon} + \eta \sqrt{\varepsilon \ln(1/\varepsilon)} + \eta^2 \right), \quad (54)$$

where C is a constant independent of ε and η .

As pointed out above, and in [42], the assumptions (34)-(35) are standard assumptions when proving convergence rates of two-scale expansions (see e.g. [40, p. 28]). Likewise, the assumption $u_0^* \in W^{2,\infty}(\mathcal{D})$ (and subsequently $\bar{u}_1^* \in W^{2,\infty}(\mathcal{D})$) is a standard assumption (see e.g. [2, Theorem 2.1] and [40, p. 28]). In view of (46), this assumption implies that the right hand side f in (30) belongs to $L^\infty(\mathcal{D})$.

In dimension $d = 1$, the boundary conditions of (53) need to be modified for this problem to have a solution. We have derived in [42] the following result, which is the one-dimensional version of Theorem 7 (note that we need below weaker assumptions than in Theorem 7, as pointed out in [42]: we do not need to assume (34)-(35), and the assumption $f \in L^2(\mathcal{D})$ is enough):

Theorem 8 (from [42], Theorem 3). *Assume that the dimension d is equal to one. Let u_η^ε be the solution to (30) in the domain \mathcal{D} with $f \in L^2(\mathcal{D})$, and assume that A_η satisfies (32)-(33). Let v_η^ε be defined by (51), where the definition (53) is replaced by*

$$\begin{cases} -[A_{\text{per}} \chi']' = [\mathbf{1}_{(0,1)} B_{\text{per}}(1 + (w^0)')] & \text{in } \mathbb{R}, \\ \chi \in L^2_{\text{loc}}(\mathbb{R}), \quad \chi' \in L^2(\mathbb{R}), \end{cases} \quad (55)$$

where w^0 solves (43). Then

$$\sqrt{\mathbb{E} \left[\|u_\eta^\varepsilon - v_\eta^\varepsilon\|_{L^\infty(\mathcal{D})}^2 \right]} + \sqrt{\mathbb{E} \left[\|u_\eta^\varepsilon - v_\eta^\varepsilon\|_{H^1(\mathcal{D})}^2 \right]} \leq C (\varepsilon + \eta \sqrt{\varepsilon} + \eta^2), \quad (56)$$

where C is a constant independent of ε and η .

The following estimate, which is proved in [42, proof of Proposition 11] and useful to demonstrate (54), will also be useful here:

Lemma 9 (from [42], proof of Proposition 11). *We assume (34) and $d > 1$. For any $p \in \mathbb{R}^d$, any $k \in \mathbb{Z}^d$, and any bounded domain $\mathcal{D} \subset \mathbb{R}^d$, the function χ_p defined by (53) satisfies*

$$\left\| \chi_p \left(\frac{\cdot}{\varepsilon} - k \right) \right\|_{L^2(\mathcal{D})}^2 \leq C \varepsilon^d R_{d,\varepsilon}, \quad (57)$$

for a constant C independent of k and ε , where $R_{d,\varepsilon} = 1$ if $d > 2$, and $R_{d,\varepsilon} = 1 + \ln(1/\varepsilon)$ if $d = 2$.

4.1.3 Main result

Before presenting our main result, we need some useful notation. Following the approach presented in Section 2.2, we recall that

$$\mathcal{W}_h := \text{span}(\phi_i^\varepsilon, i = 1, \dots, L),$$

where ϕ_i^ε are the highly oscillatory MsFEM basis functions. By construction, the solution $u_S \in \mathcal{W}_h$ of the weak stochastic MsFEM approach (16) satisfies

$$\forall v_h \in \mathcal{W}_h, \quad \mathcal{A}_{\varepsilon, \eta}^h(u_S, v_h) = b(v_h) \quad \text{a.s.} \quad (58)$$

where, for any u and v in \mathcal{W}_h ,

$$\mathcal{A}_{\varepsilon, \eta}^h(u, v) = \sum_{\mathbf{K} \in \mathcal{T}_h} \int_{\mathbf{K}} (\nabla v(x))^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \nabla u(x) dx \quad \text{and} \quad b(v) = \int_{\mathcal{D}} f(x) v(x) dx. \quad (59)$$

For future use, we also define, on the standard finite element space

$$\mathcal{V}_h := \text{span}(\phi_i^0, i = 1, \dots, L),$$

the forms

$$\tilde{\mathcal{A}}_{\varepsilon, \eta}^h(u, v) = \sum_{\mathbf{K} \in \mathcal{T}_h} \int_{\mathbf{K}} (\nabla (\mathcal{R}_{\mathbf{K}}^\varepsilon(v))(x))^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \nabla (\mathcal{R}_{\mathbf{K}}^\varepsilon(u))(x) dx \quad \text{and} \quad \tilde{b}_h(v) = \sum_{\mathbf{K} \in \mathcal{T}_h} \int_{\mathbf{K}} f(x) \mathcal{R}_{\mathbf{K}}^\varepsilon(v)(x) dx, \quad (60)$$

where the local, linear operators $\mathcal{R}_{\mathbf{K}}^\varepsilon$ are defined on \mathcal{V}_h by

$$\forall 1 \leq i \leq L, \quad \mathcal{R}_{\mathbf{K}}^\varepsilon(\phi_i^0|_{\mathbf{K}}) = \phi_i^\varepsilon|_{\mathbf{K}}. \quad (61)$$

These local operators give rise to the global operator $\mathcal{R}^\varepsilon : \mathcal{V}_h \rightarrow \mathcal{W}_h$ defined by

$$\forall \mathbf{K}, \quad \forall v \in \mathcal{V}_h, \quad \mathcal{R}^\varepsilon(v)|_{\mathbf{K}} = \mathcal{R}_{\mathbf{K}}^\varepsilon(v|_{\mathbf{K}}). \quad (62)$$

As pointed out above, the space \mathcal{W}_h is not a subspace of $H_0^1(\mathcal{D})$, as the basis functions ϕ_i^ε may have jumps at the finite element boundaries (due to the use of the oversampling technique). We will hence work with the broken H^1 -norm introduced in (18), that reads, we recall,

$$\forall v_h \in \mathcal{W}_h, \quad \|v_h\|_{H_h^1} = \left[\sum_{\mathbf{K} \in \mathcal{T}_h} \|v_h\|_{H^1(\mathbf{K})}^2 \right]^{1/2}.$$

We are now in position to present the main result of this article. We introduce the notation $Q_i^\varepsilon = \varepsilon(i + Q)$ for any $i \in \mathbb{Z}^d$, and denote by $N_{\mathbf{K}}$ the number of cells Q_i^ε in the element \mathbf{K} : $N_{\mathbf{K}} = \text{Card}(i; Q_i^\varepsilon \subset \mathbf{K})$. We make in the theorem below a regularity hypothesis on the macroscopic mesh, assuming that the volume of each element is bounded from below by αh^d , for some $\alpha > 0$, and hence that $N_{\mathbf{K}} \geq \alpha (h/\varepsilon)^d$.

Theorem 10. *Assume that A_η satisfies (32)-(33)-(34)-(35). We assume that u_0^* and \bar{u}_1^* respectively defined by (46) and (47) satisfy $u_0^* \in W^{2, \infty}(\mathcal{D})$ and $\bar{u}_1^* \in W^{2, \infty}(\mathcal{D})$. Let u_η^ε be the solution to (30) and u_S be the weakly stochastic MsFEM solution to (58). Suppose that $d > 1$, $\varepsilon \leq h$, and that there exists $\alpha > 0$, independent of \mathbf{K} , h and ε , such that $N_{\mathbf{K}} \geq \alpha \left(\frac{h}{\varepsilon} \right)^d$. We then have*

$$\sqrt{\mathbb{E} \left[\|u_\eta^\varepsilon - u_S\|_{H_h^1}^2 \right]} \leq C \left(\sqrt{\varepsilon} + h + \frac{\varepsilon}{h} + \eta \left(\frac{\varepsilon}{h} \right)^{d/2} \ln(N(h)) + \eta + \eta^2 \mathcal{C}(\eta) \right), \quad (63)$$

where C is a constant independent of ε , h and η , $N(h)$ is the number of elements \mathbf{K} in the domain \mathcal{D} (which is of order h^{-d} in dimension d), and \mathcal{C} is a bounded function as η goes to 0.

The restriction to $d > 1$ comes from the fact that the proof of this result uses the rate of convergence on the two-scale expansion of u_η^ε that we stated in Theorem 7. This rate of convergence is not optimal in dimension one, as can be seen from the comparison of (54) and (56). The one-dimensional version of the above result is stated in Section 4.3 below (see Theorem 18), where we briefly consider the one-dimensional situation.

Remark 11. *In the case $\eta = 0$, our approach reduces to the standard deterministic MsFEM and we obtain the same estimate as in the deterministic case with oversampling (see e.g. [32, Theorem 3.1]).*

4.2 Proof of Theorem 10

The proof of Theorem 10 is the direct consequence of three lemmas. First we recall the second Strang's lemma [22, Theorem 4.2.2, p. 210].

Lemma 12. *Consider a family of Hilbert spaces \mathcal{W}_h with the norm $\|\cdot\|_{H_h^1}$, a family of continuous bilinear forms $\mathcal{A}_{\varepsilon,\eta}^h$ on \mathcal{W}_h that are uniformly \mathcal{W}_h -elliptic, and a continuous linear form b on \mathcal{W}_h . For any $h > 0$, introduce u_S solution to*

$$\forall v_h \in \mathcal{W}_h, \quad \mathcal{A}_{\varepsilon,\eta}^h(u_S, v_h) = b(v_h)$$

and $u_\eta^\varepsilon \in H_0^1(\mathcal{D})$ solution to

$$\forall v \in H_0^1(\mathcal{D}), \quad \mathcal{A}_{\varepsilon,\eta}^h(u_\eta^\varepsilon, v) = b(v).$$

Then there exists a constant C independent of η , h and ε such that

$$\|u_\eta^\varepsilon - u_S\|_{H_h^1} \leq C \left(\inf_{v_h \in \mathcal{W}_h} \|u_\eta^\varepsilon - v_h\|_{H_h^1} + \sup_{w_h \in \mathcal{W}_h} \frac{|\mathcal{A}_{\varepsilon,\eta}^h(u_\eta^\varepsilon, w_h) - b(w_h)|}{\|w_h\|_{H_h^1}} \right). \quad (64)$$

The first term in the right hand side of (64) is the so-called best approximation error. The main part (step 2) of the proof of Theorem 10 is devoted to its estimation, following up on the estimate (54) provided by Theorem 7.

The second term in the right hand side of (64) is the so-called nonconforming error, which vanishes in the case $\mathcal{W}_h \subset H_0^1(\mathcal{D})$ (the method is then conforming, and we are left with the standard Céa lemma). In our case, we use the oversampling technique, hence our approximation is not conforming, and this second term does not vanish. It will be estimated in the step 3 of the proof of Theorem 10, using the following two results, which are proved in Appendix A.

Lemma 13. *Consider the two bilinear forms \mathcal{A}_η^* and $\tilde{\mathcal{A}}_{\varepsilon,\eta}^h$ respectively defined in (39) and (60). Under assumption (34), there exists a deterministic constant C , independent of η , ε and h , such that, for any $v_h \in \mathcal{V}_h$,*

$$\sup_{w_h \in \mathcal{V}_h} \frac{|\tilde{\mathcal{A}}_{\varepsilon,\eta}^h(v_h, w_h) - \mathcal{A}_\eta^*(v_h, w_h)|}{\|w_h\|_{H^1(\mathcal{D})}} \leq C \left(\frac{\varepsilon}{h} + \eta\lambda(\omega, h, \varepsilon) + \eta^2\mathcal{C}(\eta) \right) \|v_h\|_{H^1(\mathcal{D})} \quad a.s., \quad (65)$$

where C is a deterministic function independent of ε and h and bounded when $\eta \rightarrow 0$, and λ is defined by

$$\lambda(\omega, h, \varepsilon) = \max_{\mathbf{K}} \max_{1 \leq p, m \leq d} \left| \frac{1}{|I_{\mathbf{K}}^\varepsilon|} \int_{I_{\mathbf{K}}^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T \left(A_1 \left(\frac{x}{\varepsilon}, \omega \right) - \mathbb{E} \left(A_1 \left(\frac{x}{\varepsilon}, \cdot \right) \right) \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx \right|, \quad (66)$$

where $I_{\mathbf{K}}^\varepsilon$ is the largest domain composed of cells of size ε included in \mathbf{K} :

$$I_{\mathbf{K}}^\varepsilon = \bigcup_{Q_i^\varepsilon \subset \mathbf{K}} Q_i^\varepsilon, \quad Q_i^\varepsilon = \varepsilon(i + Q), \quad i \in \mathbb{Z}^d.$$

Lemma 14. *Consider the two linear forms b and \tilde{b}_h respectively defined in (39) and (60). Under assumption (34), there exists a deterministic constant C independent of η , ε and h such that*

$$\sup_{w_h \in \mathcal{V}_h} \frac{|\tilde{b}_h(w_h) - b(w_h)|}{\|w_h\|_{H^1(\mathcal{D})}} \leq C\varepsilon \|f\|_{L^2(\mathcal{D})}. \quad (67)$$

Before turning to the proof of Theorem 10, we first give some properties of the random variable $\lambda(\omega, h, \varepsilon)$ that appears in the right hand side of (65), and we next detail a two scale expansion of the highly oscillatory basis functions ϕ_i^ε , which will be useful in the sequel.

Remark 15. *We will show in Lemma 16 below that λ defined in (66) is uniformly bounded with respect to h , ε and ω . Since \mathcal{A}_η^* is coercive, we deduce from (65) that $\tilde{\mathcal{A}}_{\varepsilon,\eta}^h$ is also coercive, in the sense that there exists a deterministic constant $\alpha > 0$, independent of h , ε and η , such that*

$$\forall v_h \in \mathcal{V}_h, \quad \alpha \|v_h\|_{H^1(\mathcal{D})}^2 \leq \tilde{\mathcal{A}}_{\varepsilon,\eta}^h(v_h, v_h).$$

4.2.1 Properties of $\lambda(\omega, h, \varepsilon)$

We state here some useful properties of the random variable $\lambda(\omega, h, \varepsilon)$ that appears in (65). They will be proved in Appendix A. As mentioned above, we recall that the assumption $N_{\mathbf{K}} \geq \alpha (h/\varepsilon)^d$ that we make below is a regularity assumption on the macroscopic mesh (the volume of each element \mathbf{K} is bounded from below by αh^d).

Lemma 16. *Let $\lambda(\omega, h, \varepsilon)$ be defined by (66). Then, there exists a deterministic constant C such that, for any h and ε , we have $0 \leq \lambda(\omega, h, \varepsilon) \leq C$ almost surely.*

Assume now that the random matrix A_1 satisfies (33), where the law of $X_k(\omega)$ is absolutely continuous with respect to the Lebesgue measure. Assume furthermore that the number $N_{\mathbf{K}} = \text{Card}(i; Q_i^\varepsilon \subset \mathbf{K})$ of cells in \mathbf{K} satisfies $N_{\mathbf{K}} \geq \alpha \left(\frac{h}{\varepsilon}\right)^d$ for some $\alpha > 0$ independent of the element \mathbf{K} , h and ε . Then

$$\mathbb{E}(\lambda(\cdot, h, \varepsilon)^2) \leq C \frac{\varepsilon^d}{h^d} [\ln(N(h))]^2, \quad (68)$$

where, we recall, $N(h)$ is the number of elements \mathbf{K} in the domain \mathcal{D} (which is of order h^{-d} in dimension d) and C is a deterministic constant independent of h and ε .

Because of the specific form (33) of A_1 , we will see in the proof of that result (see Appendix A below) that

$$\lambda(\omega, h, \varepsilon) = \max_{\mathbf{K}} \max_{1 \leq m, p \leq d} |S_{\mathbf{K}}^{m,p}|, \quad (69)$$

where each random variable $S_{\mathbf{K}}^{m,p}$ is a normalized sum of $(h/\varepsilon)^d$ i.i.d. variables. Applying the Central Limit Theorem, we hence know that $S_{\mathbf{K}}^{m,p}$ converges, when $\varepsilon \rightarrow 0$, to a Gaussian random variable (up to an appropriate renormalization). Likewise, computing the expectation of $(S_{\mathbf{K}}^{m,p})^2$ is not difficult. However, in the above lemma, the difficulty stems from the fact that $\lambda(\omega, h, \varepsilon)$ is the maximum of *many* such random variables $S_{\mathbf{K}}^{m,p}$ (in (69), the number of elements \mathbf{K} is indeed of the order of h^{-d}). Our main task is hence to control how $\mathbb{E}(\lambda(\cdot, h, \varepsilon)^2)$ depends on h . See also Remark 19.

4.2.2 Two-scale expansion of the highly oscillatory basis functions

Following [32], we recall here an expansion of $\phi_i^{\varepsilon, \mathbf{K}}$ that will be useful in the sequel. By definition (see (11) and (12)), we have, for any $1 \leq i \leq d+1$,

$$\phi_i^{\varepsilon, \mathbf{K}} = \sum_{j=1}^{d+1} \alpha_{ij} \chi_j^{\varepsilon, \mathbf{S}} \Big|_{\mathbf{K}}, \quad (70)$$

where α_{ij} is such that

$$\phi_i^{0, \mathbf{K}} = \sum_{j=1}^{d+1} \alpha_{ij} \chi_j^{0, \mathbf{S}} \Big|_{\mathbf{K}}. \quad (71)$$

We thus first turn to $\chi_i^{\varepsilon, \mathbf{S}}$, which, by definition (see (10)), is the unique solution to

$$-\text{div} \left[A_{\text{per}} \left(\frac{x}{\varepsilon} \right) \nabla \chi_i^{\varepsilon, \mathbf{S}}(x) \right] = 0 \quad \text{in } \mathbf{S}, \quad \chi_i^{\varepsilon, \mathbf{S}}(x) = \chi_i^{0, \mathbf{S}}(x) \quad \text{on } \partial \mathbf{S}. \quad (72)$$

We introduce the function

$$\theta_i^{\varepsilon, \mathbf{S}}(x) = \varepsilon^{-1} \left(\chi_i^{0, \mathbf{S}}(x) + \varepsilon \sum_{j=1}^d w_{e_j}^0 \left(\frac{x}{\varepsilon} \right) \partial_j \chi_i^{0, \mathbf{S}}(x) - \chi_i^{\varepsilon, \mathbf{S}}(x) \right), \quad (73)$$

where $w_{e_j}^0$ is solution to the periodic corrector problem (43). By construction, using (73), (43), (72) and the fact that $\nabla \chi_i^{0, \mathbf{S}}$ is constant on \mathbf{S} , we have

$$\begin{aligned} -\text{div} \left[A_{\text{per}} \left(\frac{x}{\varepsilon} \right) \nabla \theta_i^{\varepsilon, \mathbf{S}}(x) \right] &= \frac{1}{\varepsilon} \text{div} \left[A_{\text{per}} \left(\frac{x}{\varepsilon} \right) \nabla \left(\chi_i^{\varepsilon, \mathbf{S}}(x) - \chi_i^{0, \mathbf{S}}(x) - \varepsilon \sum_{j=1}^d w_{e_j}^0 \left(\frac{x}{\varepsilon} \right) \partial_j \chi_i^{0, \mathbf{S}} \right) \right] \\ &= \frac{1}{\varepsilon} \text{div} \left[A_{\text{per}} \left(\frac{x}{\varepsilon} \right) \nabla \chi_i^{\varepsilon, \mathbf{S}}(x) \right] - \frac{1}{\varepsilon} \sum_{j=1}^d \partial_j \chi_i^{0, \mathbf{S}} \text{div} \left[A_{\text{per}} \left(\frac{x}{\varepsilon} \right) \left(e_j + \nabla w_{e_j}^0 \left(\frac{x}{\varepsilon} \right) \right) \right] \\ &= 0, \end{aligned}$$

while, from (73), $\theta_i^{\varepsilon, \mathbf{S}}(x) = \sum_{j=1}^d \partial_j \chi_i^{0, \mathbf{S}}(x) w_{e_j}^0 \left(\frac{x}{\varepsilon} \right)$ on $\partial \mathbf{S}$. So, by linearity, we obtain

$$\theta_i^{\varepsilon, \mathbf{S}}(x) = \sum_{j=1}^d \partial_j \chi_i^{0, \mathbf{S}} \xi_\varepsilon^j(x), \quad (74)$$

where $\xi_\varepsilon^j \in H^1(\mathbf{S})$ is the unique solution to

$$-\operatorname{div} \left[A_{\text{per}} \left(\frac{x}{\varepsilon} \right) \nabla \xi_\varepsilon^j(x) \right] = 0 \quad \text{in } \mathbf{S}, \quad \xi_\varepsilon^j(x) = w_{e_j}^0 \left(\frac{x}{\varepsilon} \right) \quad \text{on } \partial \mathbf{S}. \quad (75)$$

Using (73), we now obtain a useful relation between $\phi_i^{\varepsilon, \mathbf{K}}$ and $\phi_i^{0, \mathbf{K}}$. Indeed, collecting (70), (71), (73) and (74), we obtain the *exact* expression

$$\phi_i^{\varepsilon, \mathbf{K}}(x) = \phi_i^{0, \mathbf{K}}(x) + \varepsilon \sum_{j=1}^d \left(w_{e_j}^0 \left(\frac{x}{\varepsilon} \right) - \xi_\varepsilon^j(x) \Big|_{\mathbf{K}} \right) \partial_j \phi_i^{0, \mathbf{K}}. \quad (76)$$

Recall now that $\phi_i^{\varepsilon, \mathbf{K}}(x) = \mathcal{R}_{\mathbf{K}}^\varepsilon(\phi_i^{0, \mathbf{K}})$, by definition of the local operator $\mathcal{R}_{\mathbf{K}}^\varepsilon$ (see (61)). Correspondingly, the global operator \mathcal{R}^ε , defined on \mathcal{V}_h by (62), equivalently writes

$$\forall u \in \mathcal{V}_h, \quad \mathcal{R}^\varepsilon(u) = u + \varepsilon \sum_{j=1}^d \left(w_{e_j}^0 \left(\frac{\cdot}{\varepsilon} \right) - \overline{\xi_\varepsilon^j} \right) \partial_j u, \quad (77)$$

where $\overline{\xi_\varepsilon^j}$ is locally defined on each element \mathbf{K} by $\overline{\xi_\varepsilon^j} \Big|_{\mathbf{K}} = \xi_\varepsilon^j \Big|_{\mathbf{K}}$. By construction, for each \mathbf{K} , $\overline{\xi_\varepsilon^j} \in H^1(\mathbf{K})$, but it a priori does not belong to $H^1(\mathcal{D})$. The relation (77) allows to extend the operator \mathcal{R}^ε on $H^1(\mathcal{D})$.

We now recall the following bound on the function ξ_ε^j , that appears in (76). In [32], this lemma is stated in dimension $d = 2$, but its proof, which essentially makes use of [9, Lemma 16], carries over to any dimension.

Lemma 17 (see [32], Lemma 2.1). *Let ξ_ε^j be the solution to (75), with A_{per} satisfying (34). Consider $\mathbf{K} \subset \mathbf{S}$, with $\operatorname{diam}(\mathbf{K}) = h$ and $\operatorname{dist}(\mathbf{K}, \partial \mathbf{S}) \geq h$. Then there exists a constant C independent of h and ε such that*

$$\|\nabla \overline{\xi_\varepsilon^j}\|_{L^\infty(\mathbf{K})} \leq \frac{C}{h}. \quad (78)$$

4.2.3 Proof of Theorem 10

The proof is based on the bound (64) in Lemma 12, where the bilinear form $\mathcal{A}_{\varepsilon, \eta}^h$ and the linear form b are defined by (59). In Step 1, we show that the bilinear form $\mathcal{A}_{\varepsilon, \eta}^h$ is coercive for the norm $\|\cdot\|_{H_h^1}$ defined by (18). Step 2 is devoted to appropriately selecting an element $v_h \in \mathcal{W}_h$ such that $\|u_\eta^\varepsilon - v_h\|_{H_h^1}$ can be analytically estimated. This will provide a bound on the first term in the right hand side of (64). In Step 3, we bound from above the second term in the right hand side of (64), using Lemmas 13 and 14. Step 4 collects our estimates and concludes.

Step 1: We first show that the bilinear form $\mathcal{A}_{\varepsilon, \eta}^h$ defined by (59) is coercive for the norm $\|\cdot\|_{H_h^1}$ defined by (18). Consider the bilinear form $\tilde{\mathcal{A}}_{\varepsilon, \eta}^h$ defined by (60). We pointed out above (see Remark 15) that it is coercive on \mathcal{V}_h . Hence, there exists $\alpha > 0$ such that, for all $v_h \in \mathcal{W}_h$,

$$\alpha \|\tilde{v}_h\|_{H^1(\mathcal{D})}^2 \leq \tilde{\mathcal{A}}_{\varepsilon, \eta}^h(\tilde{v}_h, \tilde{v}_h) = \mathcal{A}_{\varepsilon, \eta}^h(v_h, v_h), \quad (79)$$

where $\tilde{v}_h \in \mathcal{V}_h$ is such that $v_h = \mathcal{R}^\varepsilon(\tilde{v}_h)$. Since, in the bilinear form $\mathcal{A}_{\varepsilon, \eta}^h$, the matrix A_η is bounded, we deduce the estimate

$$\|\tilde{v}_h\|_{H^1(\mathcal{D})}^2 \leq C \|v_h\|_{H_h^1}^2, \quad (80)$$

that we will use in the sequel. The sequel of this step is devoted to proving that there exists \tilde{C} independent of h and ε such that, for all $\tilde{v}_h \in \mathcal{V}_h$,

$$\|v_h\|_{H_h^1}^2 \leq \tilde{C} \|\tilde{v}_h\|_{H^1(\mathcal{D})}^2 \quad \text{with } v_h = \mathcal{R}^\varepsilon(\tilde{v}_h). \quad (81)$$

Combined with (79), this shows that $\mathcal{A}_{\varepsilon, \eta}^h$ is coercive for the norm $\|\cdot\|_{H_h^1}$.

To prove (81), we first write that, since $v_h = \mathcal{R}^\varepsilon(\tilde{v}_h)$ and $\tilde{v}_h \in \mathcal{V}_h$, there exist some coefficients $\{\beta_i\}_{i=1}^L$ such that, for any $x \in \mathcal{D}$, $\tilde{v}_h = \sum_{i=1}^L \beta_i \phi_i^0$ and $v_h = \mathcal{R}^\varepsilon(\tilde{v}_h) = \sum_{i=1}^L \beta_i \phi_i^\varepsilon$. Consider now an element \mathbf{K} , and its corresponding oversampling domain \mathbf{S} . We know from (11) and (13) that

$$\forall x \in \mathbf{K}, \quad \tilde{v}_h(x) = \sum_{i=1}^L \sum_{j=1}^{d+1} \beta_i \alpha_{ij} \chi_j^{0, \mathbf{S}}(x), \quad v_h(x) = \sum_{i=1}^L \sum_{j=1}^{d+1} \beta_i \alpha_{ij} \chi_j^{\varepsilon, \mathbf{S}}(x).$$

Consider now the functions

$$\tilde{w}_h^{\mathbf{S}}(x) := \sum_{i=1}^L \sum_{j=1}^{d+1} \beta_i \alpha_{ij} \chi_j^{0, \mathbf{S}}(x), \quad w_h^{\mathbf{S}}(x) := \sum_{i=1}^L \sum_{j=1}^{d+1} \beta_i \alpha_{ij} \chi_j^{\varepsilon, \mathbf{S}}(x),$$

defined on \mathbf{S} , and that satisfy, by construction,

$$\forall x \in \mathbf{K}, \quad \tilde{v}_h(x) = \tilde{w}_h^{\mathbf{S}}(x), \quad v_h(x) = w_h^{\mathbf{S}}(x). \quad (82)$$

In view of (10), we have

$$-\operatorname{div} [A^\varepsilon(x) \nabla w_h^{\mathbf{S}}(x)] = 0 \quad \text{in } \mathbf{S}, \quad w_h^{\mathbf{S}} = \tilde{w}_h^{\mathbf{S}} \quad \text{on } \partial \mathbf{S},$$

which implies that

$$\|w_h^{\mathbf{S}}\|_{H^1(\mathbf{S})} \leq C \|\tilde{w}_h^{\mathbf{S}}\|_{H^1(\mathbf{S})}.$$

We deduce from (82) and the above bound that

$$\|v_h\|_{H^1(\mathbf{K})} = \|w_h^{\mathbf{S}}\|_{H^1(\mathbf{K})} \leq \|w_h^{\mathbf{S}}\|_{H^1(\mathbf{S})} \leq C \|\tilde{w}_h^{\mathbf{S}}\|_{H^1(\mathbf{S})}. \quad (83)$$

We next see that there exists C independent of h such that, for any piecewise-affine function τ on \mathbf{S} , we have $\|\tau\|_{H^1(\mathbf{S})} \leq C \|\tau\|_{H^1(\mathbf{K})}$, provided there exists $0 < c_- \leq c_+$ independent of the element such that $c_- \leq \frac{|\mathbf{S}|}{|\mathbf{K}|} \leq c_+$. Using this bound for $\tau = \tilde{w}_h^{\mathbf{S}}$, we infer from (83) and (82) that

$$\|v_h\|_{H^1(\mathbf{K})} \leq C \|\tilde{w}_h^{\mathbf{S}}\|_{H^1(\mathbf{S})} \leq \bar{C} \|\tilde{w}_h^{\mathbf{S}}\|_{H^1(\mathbf{K})} = \bar{C} \|\tilde{v}_h\|_{H^1(\mathbf{K})}.$$

Summing over all elements \mathbf{K} , we obtain (81), and this concludes this first step.

Step 2: Let $\Pi^h u_\eta^*$ be the H^1 projection of u_η^* , solution to (36), on the standard FEM space \mathcal{V}_h . We have $\mathcal{R}^\varepsilon(\Pi^h u_\eta^*) \in \mathcal{W}_h$ (recall \mathcal{R}^ε is defined by (62), and equivalently writes as in (77)). Our argument is based on the following triangle inequality:

$$\mathbb{E} \left(\inf_{v_h \in \mathcal{W}_h} \|u_\eta^\varepsilon - v_h\|_{H_h^1}^2 \right) \leq 2\mathbb{E} \left(\|u_\eta^\varepsilon - v_\eta^\varepsilon\|_{H^1(\mathcal{D})}^2 \right) + 2\mathbb{E} \left(\inf_{v_h \in \mathcal{W}_h} \|v_\eta^\varepsilon - v_h\|_{H_h^1}^2 \right) \quad (84)$$

$$\leq 2\mathbb{E} \left(\|u_\eta^\varepsilon - v_\eta^\varepsilon\|_{H^1(\mathcal{D})}^2 \right) + 2\mathbb{E} \left(\|v_\eta^\varepsilon - \mathcal{R}^\varepsilon(\Pi^h u_\eta^*)\|_{H_h^1}^2 \right) \quad (85)$$

$$\leq 2\mathbb{E} \left(\|u_\eta^\varepsilon - v_\eta^\varepsilon\|_{H^1(\mathcal{D})}^2 \right) + 4\mathbb{E} \left(\|v_\eta^\varepsilon - \mathcal{R}^\varepsilon(u_\eta^*)\|_{H_h^1}^2 \right) + 4\|\mathcal{R}^\varepsilon(u_\eta^*) - \mathcal{R}^\varepsilon(\Pi^h u_\eta^*)\|_{H_h^1}^2, \quad (86)$$

where $v_\eta^\varepsilon(\cdot, \omega) \in H^1(\mathcal{D})$ is defined by (51). The estimate (54) in Theorem 7 bounds the first term from above. In the following two sub-steps, we bound the other two terms of (86).

Step 2a: bound on $\mathbb{E} \left(\|v_\eta^\varepsilon - \mathcal{R}^\varepsilon(u_\eta^*)\|_{H_h^1}^2 \right)$

Using the expansion (45) of u_η^* in a series in powers of η , and (77), we write

$$\mathcal{R}^\varepsilon(u_\eta^*) = u_0^* + \eta \mathbb{E}(X_0) \bar{u}_1^* + \varepsilon \sum_{p=1}^d \left(w_{e_p}^0 \left(\frac{\cdot}{\varepsilon} \right) - \bar{\xi}_\varepsilon^p \right) (\partial_p u_0^* + \eta \mathbb{E}(X_0) \partial_p \bar{u}_1^*) + \eta^2 g_\eta,$$

where

$$g_\eta = r_\eta + \varepsilon \sum_{p=1}^d \left(w_{e_p}^0 \left(\frac{\cdot}{\varepsilon} \right) - \overline{\xi_\varepsilon^p} \right) \partial_p r_\eta. \quad (87)$$

Using (51), we thus have

$$\begin{aligned} v_\eta^\varepsilon(\cdot, \omega) - \mathcal{R}^\varepsilon(u_\eta^*) &= \eta \varepsilon \sum_{p=1}^d \left(\mathbb{E}(X_0) \psi_{e_p} \left(\frac{\cdot}{\varepsilon} \right) \partial_p u_0^* + \sum_{k \in I_\varepsilon} (X_k(\omega) - \mathbb{E}(X_0)) \chi_{e_p} \left(\frac{\cdot}{\varepsilon} - k \right) \partial_p u_0^* \right) \\ &\quad + \varepsilon \sum_{p=1}^d \overline{\xi_\varepsilon^p} (\partial_p u_\eta^* - \eta^2 \partial_p r_\eta) - \eta^2 g_\eta. \end{aligned} \quad (88)$$

To bound $\mathbb{E} \left[\sum_{\mathbf{K}} \|v_\eta^\varepsilon - \mathcal{R}^\varepsilon(u_\eta^*)\|_{H^1(\mathbf{K})}^2 \right]$, we first establish a few simple results. First, there exists $\delta > 0$ such that, for any $1 \leq p \leq d$, we have

$$w_{e_p}^0 \in C^{1,\delta}(Q), \quad (89)$$

where w_p^0 is the periodic corrector, solution to (43). This is a consequence of the fact that A_{per} is Hölder-continuous (see assumption (34)), in view of [34, Theorem 8.22 and Corollary 8.36]. We infer from (89) and the periodicity of $w_{e_p}^0$ that, for any $1 \leq p \leq d$, we have

$$w_{e_p}^0 \in W^{1,\infty}(\mathbb{R}^d). \quad (90)$$

Second, for any $1 \leq p \leq d$, we have

$$\left\| \chi_{e_p} \left(\frac{\cdot}{\varepsilon} - k \right) \right\|_{L^2(\mathcal{D})}^2 \leq C \varepsilon^d R_{d,\varepsilon} \quad \text{and} \quad \nabla \chi_{e_p} \in (L^2(\mathbb{R}^d))^d, \quad (91)$$

where C is independent from ε and k , $R_{d,\varepsilon} = 1$ if $d > 2$ and $R_{d,\varepsilon} = 1 + \ln(1/\varepsilon)$ if $d = 2$ (see (53) and (57)). Third, we see that, for any $1 \leq p \leq d$,

$$\left\| \overline{\xi_\varepsilon^p} \right\|_{L^\infty(\mathcal{D})} \leq C \quad \text{and} \quad \|\nabla \xi_\varepsilon^p\|_{L^\infty(\mathbf{K})} \leq \frac{C}{h}, \quad (92)$$

where C is independent from ε and h . The second assertion is given by Lemma 17 above, whereas the first assertion comes (75): using again [34, Theorem 8.22 and Corollary 8.36] and (89), we first see that, for any \mathbf{S} , $\xi_\varepsilon^p \in C^{1,\delta}(\mathbf{S})$ for some $\delta > 0$. Using next the maximum principle on (75), we have $\|\xi_\varepsilon^p\|_{L^\infty(\mathbf{S})} \leq \left\| w_{e_p} \left(\frac{\cdot}{\varepsilon} \right) \right\|_{L^\infty(\mathbb{R}^d)} \leq C$. Lastly, using (87), (49), (90) and (92), we obtain that, for any element \mathbf{K} ,

$$\|g_\eta\|_{H^1(\mathbf{K})} \leq C \|r_\eta\|_{H^2(\mathbf{K})} \left(1 + \varepsilon + \frac{\varepsilon}{h} \right) \leq C \|r_\eta\|_{H^2(\mathbf{K})},$$

hence

$$\|g_\eta\|_{L^2(\mathcal{D})}^2 \leq \sum_{\mathbf{K}} \|g_\eta\|_{H^1(\mathbf{K})}^2 \leq C \|r_\eta\|_{H^2(\mathcal{D})}^2 \leq C. \quad (93)$$

We are now in position to estimate (88). Using that $u_0^* \in W^{1,\infty}(\mathcal{D})$, we deduce from (91), (92) and (93) that

$$\begin{aligned} \mathbb{E} \left[\|v_\eta^\varepsilon - \mathcal{R}^\varepsilon(u_\eta^*)\|_{L^2(\mathcal{D})}^2 \right] &\leq C \eta^2 \varepsilon^2 \sum_{p=1}^d \|\partial_p u_0^*\|_{L^\infty}^2 \left(\mathbb{E}(X_0)^2 \left\| \psi_{e_p} \left(\frac{\cdot}{\varepsilon} \right) \right\|_{L^2(\mathcal{D})}^2 + \text{Var}(X_0) \sum_{k \in I_\varepsilon} \left\| \chi_{e_p} \left(\frac{\cdot}{\varepsilon} - k \right) \right\|_{L^2(\mathcal{D})}^2 \right) \\ &\quad + C \varepsilon^2 \sum_{p=1}^d \left\| \overline{\xi_\varepsilon^p} \right\|_{L^\infty(\mathcal{D})}^2 \left(\|\nabla u_\eta^*\|_{L^2(\mathcal{D})}^2 + \eta^4 \|\nabla r_\eta\|_{L^2(\mathcal{D})}^2 \right) + C \eta^4 \|g_\eta\|_{L^2(\mathcal{D})}^2 \\ &\leq C \eta^2 \varepsilon^2 \left(1 + (\text{Card } I_\varepsilon) \varepsilon^d R_{d,\varepsilon} \right) + C \varepsilon^2 + C \eta^4 \\ &\leq C \left[\eta^2 \varepsilon^2 R_{d,\varepsilon} + \varepsilon^2 + \eta^4 \right] \end{aligned} \quad (94)$$

for some constant C independent of ε , η and h , and where, we recall, $R_{d,\varepsilon} = 1 + \ln(1/\varepsilon)$ if $d = 2$ and $R_{d,\varepsilon} = 1$ if $d > 2$.

We thus have a bound on $v_\eta^\varepsilon(\cdot, \omega) - \mathcal{R}^\varepsilon(u_\eta^*)$ in the L^2 norm. To prove a bound in the broken H^1 norm, we consider $\nabla v_\eta^\varepsilon(\cdot, \omega) - \nabla \mathcal{R}^\varepsilon(u_\eta^*)$: we see from (88) that, in each element \mathbf{K} ,

$$\nabla v_\eta^\varepsilon(\cdot, \omega) - \nabla \mathcal{R}^\varepsilon(u_\eta^*) = \eta\varepsilon D_0 + \eta D_1 + D_2 - \eta^2 \nabla g_\eta, \quad (95)$$

where

$$\begin{aligned} D_0 &= \sum_{p=1}^d \left(\mathbb{E}(X_0) \psi_{e_p} \left(\frac{\cdot}{\varepsilon} \right) \nabla \partial_p u_0^* + \sum_{k \in I_\varepsilon} (X_k(\omega) - \mathbb{E}(X_0)) \chi_{e_p} \left(\frac{\cdot}{\varepsilon} - k \right) \nabla \partial_p u_0^* \right), \\ D_1 &= \sum_{p=1}^d \left(\mathbb{E}(X_0) \nabla \psi_{e_p} \left(\frac{\cdot}{\varepsilon} \right) \partial_p u_0^* + \sum_{k \in I_\varepsilon} (X_k(\omega) - \mathbb{E}(X_0)) \nabla \chi_{e_p} \left(\frac{\cdot}{\varepsilon} - k \right) \partial_p u_0^* \right), \\ D_2 &= \varepsilon \sum_{p=1}^d \left(\overline{\xi_\varepsilon^p} \nabla \partial_p (u_\eta^* - \eta^2 r_\eta) + \nabla \overline{\xi_\varepsilon^p} \partial_p (u_\eta^* - \eta^2 r_\eta) \right). \end{aligned}$$

Note that D_0 and D_1 are globally defined on \mathcal{D} , but D_2 is not (as $\overline{\xi_\varepsilon^p}$ may have jumps from one element \mathbf{K} to the other). We now bound these three quantities. Using (91) and the fact that $u_0^* \in W^{2,\infty}(\mathcal{D})$, we have

$$\begin{aligned} \mathbb{E}(\|D_0\|_{L^2(\mathcal{D})}^2) &\leq C \sum_{p=1}^d \|\nabla \partial_p u_0^*\|_{L^\infty}^2 \left(\mathbb{E}(X_0)^2 \left\| \psi_{e_p} \left(\frac{\cdot}{\varepsilon} \right) \right\|_{L^2(\mathcal{D})}^2 + \text{Var}(X_0) \sum_{k \in I_\varepsilon} \left\| \chi_{e_p} \left(\frac{\cdot}{\varepsilon} - k \right) \right\|_{L^2(\mathcal{D})}^2 \right) \\ &\leq CR_{d,\varepsilon}, \end{aligned} \quad (96)$$

where C is a constant independent of ε and h . We now turn to D_1 : using again (91) and that $u_0^* \in W^{1,\infty}(\mathcal{D})$, we obtain

$$\begin{aligned} \mathbb{E}(\|D_1\|_{L^2(\mathcal{D})}^2) &\leq C \sum_{p=1}^d \|\partial_p u_0^*\|_{L^\infty}^2 \left(\mathbb{E}(X_0)^2 \left\| \nabla \psi_{e_p} \left(\frac{\cdot}{\varepsilon} \right) \right\|_{L^2(\mathcal{D})}^2 + \text{Var}(X_0) \sum_{k \in I_\varepsilon} \left\| \nabla \chi_{e_p} \left(\frac{\cdot}{\varepsilon} - k \right) \right\|_{L^2(\mathcal{D})}^2 \right) \\ &\leq C \sum_{p=1}^d \left(1 + \text{Var}(X_0) \sum_{k \in I_\varepsilon} \varepsilon^d \left\| \nabla \chi_{e_p} \right\|_{L^2(\mathbb{R}^d)}^2 \right) \\ &\leq C, \end{aligned} \quad (97)$$

where C is a constant independent of ε and h . Turning to D_2 , using (92), we have in each element \mathbf{K} that

$$\|D_2\|_{L^2(\mathbf{K})}^2 \leq C \varepsilon^2 \|u_\eta^* - \eta^2 r_\eta\|_{H^2(\mathbf{K})}^2 \left(1 + \frac{1}{h^2} \right) \leq C \frac{\varepsilon^2}{h^2} \|u_\eta^* - \eta^2 r_\eta\|_{H^2(\mathbf{K})}^2,$$

hence, using (49),

$$\sum_{\mathbf{K}} \|D_2\|_{L^2(\mathbf{K})}^2 \leq C \frac{\varepsilon^2}{h^2} \|u_\eta^* - \eta^2 r_\eta\|_{H^2(\mathcal{D})}^2 \leq C \frac{\varepsilon^2}{h^2}. \quad (98)$$

Collecting (95), (96), (97), (98) and (93), we obtain that

$$\mathbb{E} \left[\sum_{\mathbf{K}} \|\nabla v_\eta^\varepsilon - \nabla \mathcal{R}^\varepsilon(u_\eta^*)\|_{L^2(\mathbf{K})}^2 \right] \leq C \left(\eta^2 \varepsilon^2 R_{d,\varepsilon} + \eta^2 + \frac{\varepsilon^2}{h^2} + \eta^4 \right).$$

Collecting this bound with (94), and assuming that $|\eta| < 1$ and $\varepsilon^2 R_{d,\varepsilon} \leq 1$, we deduce that

$$\mathbb{E} \left[\|v_\eta^\varepsilon - \mathcal{R}^\varepsilon(u_\eta^*)\|_{H_h^1}^2 \right] = \mathbb{E} \left[\sum_{\mathbf{K}} \|v_\eta^\varepsilon - \mathcal{R}^\varepsilon(u_\eta^*)\|_{H^1(\mathbf{K})}^2 \right] \leq C \left(\eta^2 + \frac{\varepsilon^2}{h^2} \right), \quad (99)$$

where C is independent from ε , h and η .

Step 2b: bound on $\|\mathcal{R}^\varepsilon(u_\eta^*) - \mathcal{R}^\varepsilon(\Pi^h u_\eta^*)\|_{H_h^1}^2$

Recall that $\Pi^h u_\eta^*$ is the H^1 projection of u_η^* (on the standard \mathbb{P}_1 FEM space \mathcal{V}_h), hence $\|\Pi^h u_\eta^*\|_{H^1(\mathcal{D})} \leq \|u_\eta^*\|_{H^1(\mathcal{D})}$. In addition, using (49), we have, using a standard result from the theory of \mathbb{P}_1 finite elements (see [22, Theorem 3.1.6 p. 124])

$$\|u_\eta^* - \Pi^h u_\eta^*\|_{L^2(\mathcal{D})} + h\|u_\eta^* - \Pi^h u_\eta^*\|_{H^1(\mathcal{D})} \leq Ch^2 \|\nabla^2 u_\eta^*\|_{L^2(\mathcal{D})} \leq Ch^2, \quad (100)$$

where C is a constant independent of h and η . In view of (77), we have

$$\mathcal{R}^\varepsilon(u_\eta^*) - \mathcal{R}^\varepsilon(\Pi^h u_\eta^*) = u_\eta^* - \Pi^h u_\eta^* + \varepsilon \sum_{p=1}^d \left(w_{e_p}^0 \left(\frac{\cdot}{\varepsilon} \right) - \overline{\xi_\varepsilon^p} \right) \partial_p (u_\eta^* - \Pi^h u_\eta^*).$$

We deduce from (90), (92) and (100) that

$$\|\mathcal{R}^\varepsilon(u_\eta^*) - \mathcal{R}^\varepsilon(\Pi^h u_\eta^*)\|_{L^2(\mathcal{D})} \leq C(h^2 + \varepsilon h). \quad (101)$$

We now turn to bounding the gradients. Recall that $\nabla(\Pi^h u_\eta^*)$ is constant in each element \mathbf{K} . We thus have, using (90) and (92), that

$$\begin{aligned} \|\nabla \mathcal{R}^\varepsilon(u_\eta^*) - \nabla \mathcal{R}^\varepsilon(\Pi^h u_\eta^*)\|_{L^2(\mathbf{K})} &\leq \|u_\eta^* - \Pi^h u_\eta^*\|_{H^1(\mathbf{K})} \left(1 + \sum_{p=1}^d \left(\left\| \nabla w_{e_p}^0 \left(\frac{\cdot}{\varepsilon} \right) \right\|_{L^\infty(\mathbf{K})} + \varepsilon \|\nabla \xi_\varepsilon^p\|_{L^\infty(\mathbf{K})} \right) \right) \\ &\quad + \varepsilon \sum_{p=1}^d \left\| w_{e_p}^0 \left(\frac{\cdot}{\varepsilon} \right) - \xi_\varepsilon^p \right\|_{L^\infty(\mathbf{K})} \|u_\eta^*\|_{H^2(\mathbf{K})} \\ &\leq C \|u_\eta^* - \Pi^h u_\eta^*\|_{H^1(\mathbf{K})} \left(1 + \frac{\varepsilon}{h} \right) + \varepsilon \|u_\eta^*\|_{H^2(\mathbf{K})}. \end{aligned}$$

We then deduce, using (100) and (49), that

$$\begin{aligned} \sum_{\mathbf{K}} \|\nabla \mathcal{R}^\varepsilon(u_\eta^*) - \nabla \mathcal{R}^\varepsilon(\Pi^h u_\eta^*)\|_{L^2(\mathbf{K})}^2 &\leq C \|u_\eta^* - \Pi^h u_\eta^*\|_{H^1(\mathcal{D})}^2 \left(1 + \frac{\varepsilon}{h} \right)^2 + \varepsilon^2 \|u_\eta^*\|_{H^2(\mathcal{D})}^2 \\ &\leq Ch^2 \left(1 + \frac{\varepsilon}{h} \right)^2 + C\varepsilon^2, \end{aligned}$$

where C is a constant independent of ε , η and h . Collecting this bound and (101), we obtain

$$\|\mathcal{R}^\varepsilon(\Pi^h u_\eta^*) - \mathcal{R}^\varepsilon(u_\eta^*)\|_{H_h^1}^2 = \sum_{\mathbf{K}} \|\mathcal{R}^\varepsilon(u_\eta^*) - \mathcal{R}^\varepsilon(\Pi^h u_\eta^*)\|_{H^1(\mathbf{K})}^2 \leq C(h^2 + \varepsilon^2), \quad (102)$$

where C is a constant independent of ε , η and h .

Step 2c: We are now in position to bound the first term in (64). We infer from (86), (54), (99) and (102) that

$$\begin{aligned} \sqrt{\mathbb{E} \left(\inf_{v_h \in \mathcal{W}_h} \|u_\eta^\varepsilon - v_h\|_{H_h^1}^2 \right)} &\leq C \left(\sqrt{\varepsilon} + \eta \sqrt{\varepsilon \ln(1/\varepsilon)} + \eta + \frac{\varepsilon}{h} + h \right) \\ &\leq C \left(\sqrt{\varepsilon} + \eta + \frac{\varepsilon}{h} + h \right), \end{aligned} \quad (103)$$

where we have assumed that $\varepsilon \ln(1/\varepsilon) \leq 1$.

Step 3: We next turn to estimating the non-conforming error, namely the second term of the right-hand side of (64). For any $w_h \in \mathcal{W}_h$, introduce $\tilde{w}_h \in \mathcal{V}_h$ such that $\mathcal{R}^\varepsilon(\tilde{w}_h) = w_h$ (recall that \mathcal{R}^ε is defined by (62)). We note that $b(w_h) = \tilde{b}_h(\tilde{w}_h)$, where the linear forms b and \tilde{b}_h are defined by (59) and (60). Using the weak form of the homogenized equation (see (39)), we see that $b(\tilde{w}_h) = \mathcal{A}_\eta^*(u_\eta^*, \tilde{w}_h)$. In addition, by definition of $\tilde{\mathcal{A}}_{\varepsilon,\eta}^h$ (see (60)), we have $\tilde{\mathcal{A}}_{\varepsilon,\eta}^h(\Pi^h u_\eta^*, \tilde{w}_h) = \mathcal{A}_{\varepsilon,\eta}^h(\mathcal{R}^\varepsilon(\Pi^h u_\eta^*), w_h)$. For any $w_h \in \mathcal{W}_h$, we have

$$\begin{aligned} |\mathcal{A}_{\varepsilon,\eta}^h(u_\eta^\varepsilon, w_h) - b(w_h)| &\leq |\mathcal{A}_{\varepsilon,\eta}^h(u_\eta^\varepsilon, w_h) - \mathcal{A}_{\varepsilon,\eta}^h(\mathcal{R}^\varepsilon(\Pi^h u_\eta^*), w_h)| + |\mathcal{A}_{\varepsilon,\eta}^h(\mathcal{R}^\varepsilon(\Pi^h u_\eta^*), w_h) - b(\tilde{w}_h)| + |b(\tilde{w}_h) - b(w_h)| \\ &\leq \|A_\eta\|_{L^\infty} \|u_\eta^\varepsilon - \mathcal{R}^\varepsilon(\Pi^h u_\eta^*)\|_{H_h^1} \|w_h\|_{H_h^1} + \left| \tilde{\mathcal{A}}_{\varepsilon,\eta}^h(\Pi^h u_\eta^*, \tilde{w}_h) - \mathcal{A}_\eta^*(u_\eta^*, \tilde{w}_h) \right| + \left| b(\tilde{w}_h) - \tilde{b}_h(\tilde{w}_h) \right| \\ &\leq \|A_\eta\|_{L^\infty} \|u_\eta^\varepsilon - \mathcal{R}^\varepsilon(\Pi^h u_\eta^*)\|_{H_h^1} \|w_h\|_{H_h^1} + \left| \tilde{\mathcal{A}}_{\varepsilon,\eta}^h(\Pi^h u_\eta^*, \tilde{w}_h) - \mathcal{A}_\eta^*(\Pi^h u_\eta^*, \tilde{w}_h) \right| \\ &\quad + \|A_\eta^*\| \|u_\eta^* - \Pi^h u_\eta^*\|_{H^1(\mathcal{D})} \|\tilde{w}_h\|_{H^1(\mathcal{D})} + \left| b(\tilde{w}_h) - \tilde{b}_h(\tilde{w}_h) \right|, \end{aligned}$$

where we have successively used the continuity of the bilinear forms $\mathcal{A}_{\varepsilon,\eta}^h$ and \mathcal{A}_η^* . Using Lemmas 13 and 14 for the second and the fourth terms respectively, we deduce that

$$\begin{aligned} |\mathcal{A}_{\varepsilon,\eta}^h(u_\eta^\varepsilon, w_h) - b(w_h)| &\leq C \|u_\eta^\varepsilon - \mathcal{R}^\varepsilon(\Pi_h u_\eta^*)\|_{H_h^1} \|w_h\|_{H_h^1} + C \left(\frac{\varepsilon}{h} + \eta \lambda(\omega, h, \varepsilon) + \eta^2 \mathcal{C}(\eta) \right) \|\Pi^h u_\eta^*\|_{H^1(\mathcal{D})} \|\tilde{w}_h\|_{H^1(\mathcal{D})} \\ &\quad + C \|u_\eta^* - \Pi^h u_\eta^*\|_{H^1(\mathcal{D})} \|\tilde{w}_h\|_{H^1(\mathcal{D})} + C \varepsilon \|\tilde{w}_h\|_{H^1(\mathcal{D})}, \end{aligned}$$

hence, using (80) and (100),

$$\frac{|\mathcal{A}_{\varepsilon,\eta}^h(u_\eta^\varepsilon, w_h) - b(w_h)|}{\|w_h\|_{H_h^1}} \leq C \|u_\eta^\varepsilon - \mathcal{R}^\varepsilon(\Pi_h u_\eta^*)\|_{H_h^1} + C \left(h + \frac{\varepsilon}{h} + \eta \lambda(\omega, h, \varepsilon) + \varepsilon + \eta^2 \mathcal{C}(\eta) \right). \quad (104)$$

The first term is bounded as in Step 2:

$$\|u_\eta^\varepsilon - \mathcal{R}^\varepsilon(\Pi_h u_\eta^*)\|_{H_h^1} \leq \|u_\eta^\varepsilon - v_\eta^\varepsilon\|_{H^1(\mathcal{D})} + \|v_\eta^\varepsilon - \mathcal{R}^\varepsilon(u_\eta^*)\|_{H_h^1} + \|\mathcal{R}^\varepsilon(u_\eta^*) - \mathcal{R}^\varepsilon(\Pi_h u_\eta^*)\|_{H_h^1},$$

hence, using (54), (99) and (102), and assuming that $\varepsilon \ln(1/\varepsilon) \leq 1$, we have

$$\sqrt{\mathbb{E} \left[\|u_\eta^\varepsilon - \mathcal{R}^\varepsilon(\Pi_h u_\eta^*)\|_{H_h^1}^2 \right]} \leq C \left(\sqrt{\varepsilon} + \eta + \frac{\varepsilon}{h} + h \right). \quad (105)$$

Collecting (104) and (105), we thus obtain

$$\sqrt{\mathbb{E} \left[\left(\sup_{w_h \in \mathcal{W}_h} \frac{|\mathcal{A}_{\varepsilon,\eta}^h(u_\eta^\varepsilon, w_h) - b(w_h)|}{\|w_h\|_{H_h^1}} \right)^2 \right]} \leq C \left(h + \frac{\varepsilon}{h} + \eta \sqrt{\mathbb{E}[\lambda^2(\cdot, h, \varepsilon)]} + \sqrt{\varepsilon} + \eta + \eta^2 \mathcal{C}(\eta) \right). \quad (106)$$

Step 4: Collecting (64), (103) and (106), we get

$$\sqrt{\mathbb{E} \left[\|u_\eta^\varepsilon - u_S\|_{H_h^1}^2 \right]} \leq C \left(\sqrt{\varepsilon} + h + \frac{\varepsilon}{h} + \eta \sqrt{\mathbb{E}[\lambda^2(\cdot, h, \varepsilon)]} + \eta + \eta^2 \mathcal{C}(\eta) \right)$$

where C is a constant independent of ε , h and η , and \mathcal{C} is a bounded function as η goes to 0. Using (68), we deduce that

$$\sqrt{\mathbb{E} \left[\|u_\eta^\varepsilon - u_S\|_{H_h^1}^2 \right]} \leq C \left(\sqrt{\varepsilon} + h + \frac{\varepsilon}{h} + \eta \left(\frac{\varepsilon}{h} \right)^{d/2} \ln(N(h)) + \eta + \eta^2 \mathcal{C}(\eta) \right)$$

where $N(h)$ is the number of elements \mathbf{K} in the domain (which is of order h^{-d} in dimension d). This concludes the proof of Theorem 10.

4.3 The one dimensional case

In this section, we briefly consider the one dimensional situation. As in the multi-dimensional case, we assume here that $a_\eta^\varepsilon(x, \omega) = a_\eta \left(\frac{x}{\varepsilon}, \omega \right)$, where a_η is a stationary random function satisfying, for any $|\eta| \leq 1$, the condition $0 < a_- \leq a_\eta(x, \omega) \leq a_+$ almost everywhere in \mathbb{R} , almost surely. In line with (32), we assume that

$$a_\eta(x, \omega) = a_{per}(x) + \eta a_1(x, \omega), \quad (107)$$

where η is a small parameter ($|\eta| \leq 1$), a_{per} is a 1-periodic function satisfying the condition $0 < a_- \leq a_{per}(x) \leq a_+$ almost everywhere on \mathbb{R} , and a_1 is a bounded stationary random function: $|a_1(x, \omega)| \leq C$ almost everywhere in \mathbb{R} , almost surely. In the vein of (33), we suppose that

$$a_1(x, \omega) = \sum_{k \in \mathbb{Z}} \mathbf{1}_{(k, k+1]}(x) X_k(\omega) b_{per}(x) \quad \text{such that} \quad \exists C, \forall k \in \mathbb{Z}, \quad |X_k(\omega)| \leq C \quad \text{almost surely}, \quad (108)$$

where $(X_k(\omega))_{k \in \mathbb{Z}}$ is a sequence of i.i.d. scalar random variables and $b_{per} \in L^\infty(\mathbb{R})$ is a 1-periodic function.

Note that, in this one-dimensional setting, we do not make any regularity assumption on a_{per} (in the vein of (34)). In the multi-dimensional case, this assumption is useful to e.g. state that the periodic corrector satisfies $w_p^0 \in W^{1,\infty}(\mathbb{R}^d)$ for any $p \in \mathbb{R}^d$. In the one-dimensional case, the corrector problem can be solved analytically, and one can see that the above assumption $a_{per}(x) \geq a_- > 0$ almost everywhere on \mathbb{R} is sufficient to obtain such regularity on the corrector. Similarly, we do not need to assume here, in contrast to Theorem 10, that u_0^* and \bar{u}_1^* defined by (46) and (47) both belong to $W^{2,\infty}(\mathcal{D})$ (an assumption equivalent to $f \in L^\infty(\mathcal{D})$, in the present one-dimensional setting). The assumption $f \in L^2(\mathcal{D})$ is sufficient.

The problem (30) now reads

$$-\frac{d}{dx} \left(a_\eta \left(\frac{x}{\varepsilon}, \omega \right) \frac{d}{dx} u_\eta^\varepsilon(x, \omega) \right) = f(x) \quad \text{in } (0, 1), \quad u_\eta^\varepsilon(0, \omega) = u_\eta^\varepsilon(1, \omega) = 0. \quad (109)$$

We consider a uniform discretization of the interval $(0, 1)$ in the elements $\mathbf{K}_i = (x_i, x_{i+1})$, with $x_{i+1} - x_i = h = 1/L$ for some $L \in \mathbb{N}^*$.

The one-dimensional version of Theorem 10 reads as follows:

Theorem 18. *In the one-dimensional setting, assume that a_η^ε satisfies (107)-(108). Let u_η^ε be the solution to (109) with $f \in L^2(0, 1)$, and u_S be the weakly stochastic MsFEM solution to (58). Suppose that $h/\varepsilon \in \mathbb{N}^*$. We then have*

$$\sqrt{\mathbb{E} \left[\|u_\eta^\varepsilon - u_S\|_{H^1}^2 \right]} \leq C \left(\varepsilon + h + \eta \left(\frac{\varepsilon}{h} \right)^{1/2} \ln(1/h) + \eta + \eta^2 \mathcal{C}(\eta) \right), \quad (110)$$

where C is a constant independent of ε , h and η and \mathcal{C} is a bounded function as η goes to 0.

Proof. The proof of this result follows the same lines as that for the multi-dimensional case. It is based upon the homogenization result contained in Theorem 8 above. \square

As pointed out above, the rate of convergence stated in Theorem 7 (and hence the estimate provided by Theorem 10) is not optimal in dimension one. This hence motivates Theorems 8 and 18, which are their respective one-dimensional variants. On another note, the assumption $h/\varepsilon \in \mathbb{N}^*$ implies that some terms in the error bound vanish. A result similar to (110) holds in the absence of such assumption, with the additional term ε/h in the right-hand side.

A Proofs of Lemmas 13, 14 and 16

Proof of Lemma 13. This result relies on the expansion

$$\phi_j^{\varepsilon, \mathbf{K}}(x) = \phi_j^{0, \mathbf{K}}(x) + \varepsilon \sum_{m=1}^d \left(w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) - \xi_\varepsilon^m(x) |_{\mathbf{K}} \right) \partial_m \phi_j^{0, \mathbf{K}}$$

from (76) and the fact that $\nabla \phi_j^{0, \mathbf{K}}$ is constant on \mathbf{K} .

For any v_h and w_h in \mathcal{V}_h , we write

$$\left| \mathcal{A}_\eta^*(v_h, w_h) - \tilde{\mathcal{A}}_{\varepsilon, \eta}^h(v_h, w_h) \right| = \left| \sum_{\mathbf{K} \in \mathcal{T}_h} \left(\int_{\mathbf{K}} (\nabla w_h(x))^T A_\eta^* \nabla v_h(x) dx - \sum_{i,j=1}^L v_h^j w_h^i \int_{\mathbf{K}} (\nabla \phi_i^{\varepsilon, \mathbf{K}})^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \nabla \phi_j^{\varepsilon, \mathbf{K}} dx \right) \right|,$$

where $v_h = \sum_{j=1}^L v_h^j \phi_j^0$ and likewise for w_h . Using the above expansion of $\phi_j^{\varepsilon, \mathbf{K}}$ and the fact that $\nabla \phi_j^{0, \mathbf{K}}$ is constant on

\mathbf{K} , we have

$$\begin{aligned}
& \sum_{i,j=1}^L v_h^j w_h^i \int_{\mathbf{K}} \left(\nabla \phi_i^{\varepsilon, \mathbf{K}} \right)^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \nabla \phi_j^{\varepsilon, \mathbf{K}} dx \\
&= \sum_{m,p=1}^d \frac{1}{|\mathbf{K}|} \int_{\mathbf{K}} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) - \varepsilon \nabla \xi_\varepsilon^p(x) \right]^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) - \varepsilon \nabla \xi_\varepsilon^m(x) \right] dx \\
&\quad \times \sum_{i,j=1}^L v_h^j w_h^i \int_{\mathbf{K}} \partial_p \phi_i^{0, \mathbf{K}} \partial_m \phi_j^{0, \mathbf{K}} \\
&= \sum_{m,p=1}^d \frac{1}{|\mathbf{K}|} \int_{\mathbf{K}} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) - \varepsilon \nabla \xi_\varepsilon^p(x) \right]^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) - \varepsilon \nabla \xi_\varepsilon^m(x) \right] dx \int_{\mathbf{K}} \partial_m v_h \partial_p w_h.
\end{aligned}$$

We thus obtain

$$\left| \mathcal{A}_\eta^*(v_h, w_h) - \tilde{\mathcal{A}}_{\varepsilon, \eta}^h(v_h, w_h) \right| = \left| \sum_{\mathbf{K}} \sum_{m,p=1}^d \Lambda_{mp}^{\mathbf{K}} \int_{\mathbf{K}} \partial_m v_h \partial_p w_h \right| \leq \sum_{\mathbf{K}} \|v_h\|_{H^1(\mathbf{K})} \|w_h\|_{H^1(\mathbf{K})} \sum_{m,p=1}^d |\Lambda_{mp}^{\mathbf{K}}|, \quad (111)$$

where

$$\Lambda_{mp}^{\mathbf{K}} = [A_\eta^*]_{mp} - \frac{1}{|\mathbf{K}|} \int_{\mathbf{K}} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) - \varepsilon \nabla \xi_\varepsilon^p(x) \right]^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) - \varepsilon \nabla \xi_\varepsilon^m(x) \right] dx,$$

which we write

$$\Lambda_{mp}^{\mathbf{K}} = D_0 + D_1 - D_2, \quad (112)$$

with

$$\begin{aligned}
D_0 &= [A_\eta^*]_{mp} - \frac{1}{|\mathbf{K}|} \int_{\mathbf{K}} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx, \\
D_1 &= \frac{\varepsilon}{|\mathbf{K}|} \left(\int_{\mathbf{K}} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \nabla \xi_\varepsilon^m(x) dx + \int_{\mathbf{K}} (\nabla \xi_\varepsilon^p(x))^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx \right), \\
D_2 &= \frac{\varepsilon^2}{|\mathbf{K}|} \int_{\mathbf{K}} (\nabla \xi_\varepsilon^p(x))^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \nabla \xi_\varepsilon^m(x) dx.
\end{aligned} \quad (113)$$

We are thus left with bounding $|\Lambda_{mp}^{\mathbf{K}}|$ from above. We first bound D_1 and D_2 . Using Lemma 17 (recall that A_{per} satisfies (34), i.e. is Hölder continuous) and the fact that $w_{e_p}^0 \in H^1(Q)$ and is Q -periodic, we obtain

$$|D_2| \leq C \frac{\varepsilon^2}{h^2} \quad (114)$$

and

$$\begin{aligned}
\left| \frac{\varepsilon}{|\mathbf{K}|} \int_{\mathbf{K}} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \nabla \xi_\varepsilon^m(x) dx \right| &\leq \frac{\varepsilon}{|\mathbf{K}|} \|A_\eta\|_{L^\infty} \frac{C}{h} \int_{\mathbf{K}} \left| e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right| dx \\
&\leq C \frac{\varepsilon}{|\mathbf{K}| h} \left(|\mathbf{K}| + \varepsilon^d \int_{\mathbf{K}/\varepsilon} |\nabla w_{e_p}^0(y)| dy \right) \\
&\leq C \frac{\varepsilon}{h}
\end{aligned}$$

hence

$$|D_1| \leq C \frac{\varepsilon}{h}, \quad (115)$$

where C is a deterministic constant independent of h , ε and η . We next turn to D_0 . We introduce the cells $Q_i^\varepsilon = \varepsilon(Q + i)$, $i \in \mathbb{Z}^d$, let $I_{\mathbf{K}}^\varepsilon = \bigcup_{Q_i^\varepsilon \subset \mathbf{K}} Q_i^\varepsilon$, and recast (113) as

$$D_0 = D_0^{\text{bulk}} - D_0^{\text{boundary}} \quad (116)$$

with

$$\begin{aligned} D_0^{\text{bulk}} &= [A_\eta^*]_{mp} - \frac{1}{|\mathbf{K}|} \int_{I_{\bar{\mathbf{K}}}^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx, \\ D_0^{\text{boundary}} &= \frac{1}{|\mathbf{K}|} \int_{\mathbf{K} \setminus I_{\bar{\mathbf{K}}}^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx. \end{aligned} \quad (117)$$

We denote by $J_{\mathbf{K}}^\varepsilon$ the set of cells Q_i^ε that intersect the element \mathbf{K} , i.e.

$$J_{\mathbf{K}}^\varepsilon = \bigcup_{Q_i^\varepsilon \cap \mathbf{K} \neq \emptyset} Q_i^\varepsilon.$$

By construction, $I_{\bar{\mathbf{K}}}^\varepsilon \subset \mathbf{K} \subset J_{\bar{\mathbf{K}}}^\varepsilon$. Using that $w_{e_p}^0 \in H^1(Q)$ and is Q -periodic, we write

$$\begin{aligned} |D_0^{\text{boundary}}| &\leq \|A_\eta\|_{L^\infty} \frac{1}{|\mathbf{K}|} \sqrt{\int_{J_{\bar{\mathbf{K}}}^\varepsilon \setminus I_{\bar{\mathbf{K}}}^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^2 dx} \sqrt{\int_{J_{\bar{\mathbf{K}}}^\varepsilon \setminus I_{\bar{\mathbf{K}}}^\varepsilon} \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right]^2 dx} \\ &\leq C \frac{\varepsilon^d}{|\mathbf{K}| \varepsilon^{d-1}} \sqrt{\int_Q \left[e_p + \nabla w_{e_p}^0(y) \right]^2 dy} \sqrt{\int_Q \left[e_m + \nabla w_{e_m}^0(y) \right]^2 dy} \\ &\leq C \frac{\varepsilon}{h}. \end{aligned} \quad (118)$$

We next consider (117):

$$D_0^{\text{bulk}} = \frac{|\mathbf{K} \setminus I_{\bar{\mathbf{K}}}^\varepsilon|}{|\mathbf{K}|} [A_\eta^*]_{mp} + \frac{|I_{\bar{\mathbf{K}}}^\varepsilon|}{|\mathbf{K}|} \overline{D}_0^{\text{bulk}}, \quad (119)$$

with

$$\overline{D}_0^{\text{bulk}} = [A_\eta^*]_{mp} - \frac{1}{|I_{\bar{\mathbf{K}}}^\varepsilon|} \int_{I_{\bar{\mathbf{K}}}^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T A_\eta \left(\frac{x}{\varepsilon}, \omega \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx.$$

Using the expansion (40) of A_η^* , we write

$$\begin{aligned} \overline{D}_0^{\text{bulk}} &= \int_Q \left[e_p + \nabla w_{e_p}^0(y) \right]^T A_{per}(y) \left[e_m + \nabla w_{e_m}^0(y) \right] dy \\ &\quad - \frac{1}{|I_{\bar{\mathbf{K}}}^\varepsilon|} \int_{I_{\bar{\mathbf{K}}}^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T A_{per} \left(\frac{x}{\varepsilon} \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx \\ &\quad + \eta \left(\int_Q \left[e_p + \nabla w_{e_p}^0(y) \right]^T \mathbb{E}(A_1(y, \cdot)) \left[e_m + \nabla w_{e_m}^0(y) \right] dy \right. \\ &\quad \left. - \frac{1}{|I_{\bar{\mathbf{K}}}^\varepsilon|} \int_{I_{\bar{\mathbf{K}}}^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T A_1 \left(\frac{x}{\varepsilon}, \omega \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx \right) + \eta^2 \mathcal{C}(\eta). \end{aligned} \quad (120)$$

The leading order term in (120) vanishes. We are hence left with

$$\overline{D}_0^{\text{bulk}} = \frac{\eta}{|I_{\bar{\mathbf{K}}}^\varepsilon|} \int_{I_{\bar{\mathbf{K}}}^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T \left(\mathbb{E} \left(A_1 \left(\frac{x}{\varepsilon}, \cdot \right) \right) - A_1 \left(\frac{x}{\varepsilon}, \omega \right) \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx + \eta^2 \mathcal{C}(\eta). \quad (121)$$

Collecting (112), (114), (115), (116), (118), (119) and (121), together with the fact that $\frac{|\mathbf{K} \setminus I_{\bar{\mathbf{K}}}^\varepsilon|}{|\mathbf{K}|} |[A_\eta^*]_{mp}| \leq C \frac{\varepsilon}{h}$, we obtain

$$|\Lambda_{mp}^{\mathbf{K}}| \leq C \frac{\varepsilon}{h} + \frac{\eta}{|I_{\bar{\mathbf{K}}}^\varepsilon|} \left| \int_{I_{\bar{\mathbf{K}}}^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T \left(\mathbb{E} \left(A_1 \left(\frac{x}{\varepsilon}, \cdot \right) \right) - A_1 \left(\frac{x}{\varepsilon}, \omega \right) \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx \right| + \eta^2 \mathcal{C}(\eta).$$

We set

$$\lambda(\omega, h, \varepsilon) = \max_{\mathbf{K} \in \mathcal{T}_h} \max_{1 \leq m, p \leq d} \left| \frac{1}{|I_{\bar{\mathbf{K}}}^\varepsilon|} \int_{I_{\bar{\mathbf{K}}}^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T \left(\mathbb{E} \left(A_1 \left(\frac{x}{\varepsilon}, \cdot \right) \right) - A_1 \left(\frac{x}{\varepsilon}, \omega \right) \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx \right|.$$

We thus have, for any \mathbf{K} , $\sum_{m,p=1}^d |\Lambda_{mp}^{\mathbf{K}}| \leq C \left(\frac{\varepsilon}{h} + \eta\lambda(\omega, h, \varepsilon) + \eta^2\mathcal{C}(\eta) \right)$. Using (111), we thus obtain that

$$\left| \mathcal{A}_\eta^*(v_h, w_h) - \tilde{\mathcal{A}}_{\varepsilon, \eta}^h(v_h, w_h) \right| \leq C \left(\frac{\varepsilon}{h} + \eta\lambda(\omega, h, \varepsilon) + \eta^2\mathcal{C}(\eta) \right) \|v_h\|_{H^1(\mathcal{D})} \|w_h\|_{H^1(\mathcal{D})}.$$

This concludes the proof of Lemma 13. \square

Proof of Lemma 14. Again, as for Lemma 13, this result relies on the expansion (76) of $\phi_i^{\varepsilon, \mathbf{K}}$ and the fact that $\nabla\phi_i^{0, \mathbf{K}}$ is constant on \mathbf{K} .

Setting $w_h(x) = \sum_i w_h^i \phi_i^0(x)$, we observe that

$$\left| \tilde{b}_h(w_h) - b(w_h) \right| = \left| \sum_{\mathbf{K}} \sum_{i=1}^L w_h^i \int_{\mathbf{K}} f(x) \left(\phi_i^{\varepsilon, \mathbf{K}}(x) - \phi_i^{0, \mathbf{K}}(x) \right) dx \right|.$$

Using (76) and the fact that $\nabla\phi_i^{0, \mathbf{K}}$ is constant on \mathbf{K} , we obtain

$$\sum_{i=1}^L w_h^i \int_{\mathbf{K}} f(x) \left(\phi_i^{\varepsilon, \mathbf{K}}(x) - \phi_i^{0, \mathbf{K}}(x) \right) dx = \sum_{p=1}^d \frac{\varepsilon}{|\mathbf{K}|} \int_{\mathbf{K}} f(x) \left(w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) - \xi_\varepsilon^p(x) \right) dx \sum_{i=1}^L w_h^i \int_{\mathbf{K}} \partial_p \phi_i^{0, \mathbf{K}}. \quad (122)$$

We have

$$\left| \sum_{i=1}^L w_h^i \int_{\mathbf{K}} \partial_p \phi_i^{0, \mathbf{K}} \right| = \left| \int_{\mathbf{K}} \partial_p w_h \right| \leq \sqrt{|\mathbf{K}|} \|w_h\|_{H^1(\mathbf{K})} \quad (123)$$

and

$$\begin{aligned} \left| \int_{\mathbf{K}} f(x) \left(w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) - \xi_\varepsilon^p(x) \right) dx \right| &\leq \|f\|_{L^2(\mathbf{K})} \left(\left\| w_{e_p}^0 \left(\frac{\cdot}{\varepsilon} \right) \right\|_{L^2(\mathbf{K})} + \|\xi_\varepsilon^p\|_{L^2(\mathbf{K})} \right) \\ &\leq \|f\|_{L^2(\mathbf{K})} \sqrt{|\mathbf{K}|} \left(\|w_{e_p}^0\|_{L^\infty(\mathbb{R}^d)} + \|\xi_\varepsilon^p\|_{L^\infty(\mathbf{K})} \right). \end{aligned} \quad (124)$$

Recall now that, since A_{per} satisfies (34) (i.e. is Hölder continuous), we know that ξ_ε^p and $w_{e_p}^0$ are both continuous, and that $w_{e_p}^0 \in L^\infty(\mathbb{R}^d)$. Using the maximum principle on (75), we write

$$\|\xi_\varepsilon^p\|_{L^\infty(\mathbf{K})} \leq \|w_{e_p}^0\|_{L^\infty(\partial\mathbf{S})} \leq \|w_{e_p}^0\|_{L^\infty(\mathbb{R}^d)},$$

and we thus deduce from (124) that

$$\left| \int_{\mathbf{K}} f(x) \left(w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) - \xi_\varepsilon^p(x) \right) dx \right| \leq C \|f\|_{L^2(\mathbf{K})} \sqrt{|\mathbf{K}|} \quad (125)$$

for a constant C independent of h and ε . Collecting (122), (123) and (125), we obtain

$$\left| \tilde{b}_h(w_h) - b(w_h) \right| \leq C\varepsilon \|f\|_{L^2(\mathcal{D})} \|w_h\|_{H^1(\mathcal{D})}.$$

This concludes the proof of Lemma 14. \square

Proof of Lemma 16. We first prove the uniform bound on λ . Recall that the field A_1 is bounded almost surely and almost everywhere. This implies that

$$\begin{aligned} &\left| \frac{1}{|I_{\mathbf{K}}^\varepsilon|} \int_{I_{\mathbf{K}}^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T \left(A_1 \left(\frac{x}{\varepsilon}, \omega \right) - \mathbb{E} \left(A_1 \left(\frac{x}{\varepsilon}, \cdot \right) \right) \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx \right| \\ &\leq 2 \|A_1\|_{L^\infty} \frac{1}{|I_{\mathbf{K}}^\varepsilon|} \left\| e_p + \nabla w_{e_p}^0 \left(\frac{\cdot}{\varepsilon} \right) \right\|_{L^2(I_{\mathbf{K}}^\varepsilon)} \left\| e_m + \nabla w_{e_m}^0 \left(\frac{\cdot}{\varepsilon} \right) \right\|_{L^2(I_{\mathbf{K}}^\varepsilon)}. \end{aligned}$$

Then, using the Q -periodicity of $w_{e_p}^0$, we obtain

$$\left\| e_p + \nabla w_{e_p}^0 \left(\frac{\cdot}{\varepsilon} \right) \right\|_{L^2(I_{\mathbf{K}}^\varepsilon)}^2 = \sum_{Q_i^\varepsilon \subset I_{\mathbf{K}}^\varepsilon} \int_{Q_i^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^2 dx = |I_{\mathbf{K}}^\varepsilon| \left\| e_p + \nabla w_{e_p}^0 \right\|_{L^2(Q)}^2.$$

We thus have

$$\lambda(\omega, h, \varepsilon) \leq 2 \|A_1\|_{L^\infty} \max_{1 \leq p, m \leq d} \left[\left\| e_p + \nabla w_{e_p}^0 \right\|_{L^2(Q)} \left\| e_m + \nabla w_{e_m}^0 \right\|_{L^2(Q)} \right],$$

hence $\lambda(\omega, h, \varepsilon)$ is bounded almost surely by a deterministic constant independent of h and ε .

We next turn to (68). Rewrite (66) as

$$\lambda(\omega, h, \varepsilon) = \max_{\mathbf{K}} \max_{1 \leq m, p \leq d} |S_{\mathbf{K}}^{m,p}|,$$

with

$$S_{\mathbf{K}}^{m,p} := \frac{1}{|I_{\mathbf{K}}^\varepsilon|} \int_{I_{\mathbf{K}}^\varepsilon} \left[e_p + \nabla w_{e_p}^0 \left(\frac{x}{\varepsilon} \right) \right]^T \left(A_1 \left(\frac{x}{\varepsilon}, \omega \right) - \mathbb{E} \left(A_1 \left(\frac{x}{\varepsilon}, \cdot \right) \right) \right) \left[e_m + \nabla w_{e_m}^0 \left(\frac{x}{\varepsilon} \right) \right] dx.$$

Using the periodicity of the correctors w_p^0 and the specific form (33) of A_1 , we have

$$S_{\mathbf{K}}^{m,p} = \tau^{m,p} \frac{1}{N_{\mathbf{K}}} \sum_{i; Q_i^\varepsilon \subset I_{\mathbf{K}}^\varepsilon} \frac{X_i - \mathbb{E}(X_0)}{\sqrt{\text{Var}(X_0)}} \quad (126)$$

with

$$\tau^{m,p} = \sqrt{\text{Var}(X_0)} \int_Q \left[e_p + \nabla w_{e_p}^0(y) \right]^T B_{per}(y) \left[e_m + \nabla w_{e_m}^0(y) \right] dy \quad \text{and} \quad N_{\mathbf{K}} = \text{Card}\{i; Q_i^\varepsilon \subset I_{\mathbf{K}}^\varepsilon\}.$$

Thus, $\lambda(\omega, h, \varepsilon)$ reads

$$\lambda(\omega, h, \varepsilon) = \gamma \max_{\mathbf{K}} |S_{\mathbf{K}}^\varepsilon(\omega)|,$$

where $\gamma = \max_{1 \leq m, p \leq d} \tau^{m,p}$ and

$$S_{\mathbf{K}}^\varepsilon(\omega) = \frac{1}{N_{\mathbf{K}}} \sum_{i; Q_i^\varepsilon \subset I_{\mathbf{K}}^\varepsilon} \frac{X_i - \mathbb{E}(X_0)}{\sqrt{\text{Var}(X_0)}}.$$

Introduce the probability density function $\varphi_{N_{\mathbf{K}}}$ of the random variable $\sqrt{N_{\mathbf{K}}} |S_{\mathbf{K}}^\varepsilon(\omega)|$, and $F_{N_{\mathbf{K}}}(x) = \mathbb{P}(\sqrt{N_{\mathbf{K}}} |S_{\mathbf{K}}^\varepsilon| \leq x)$.

Using the assumption that each element \mathbf{K} contains a number $N_{\mathbf{K}}$ of cells of size ε that satisfies $N_{\mathbf{K}} \geq \alpha \left(\frac{h}{\varepsilon} \right)^d$ for some $\alpha > 0$, independent of \mathbf{K} , h and ε , we write

$$\mathbb{E} \left(\alpha \frac{h^d}{\varepsilon^d} \frac{\lambda^2(\cdot, h, \varepsilon)}{\gamma^2} \right) \leq \mathbb{E} \left(N_{\mathbf{K}} \max_{\mathbf{K}} |S_{\mathbf{K}}^\varepsilon|^2 \right) = \int_0^\infty x^2 \frac{d}{dx} \mathbb{P} \left(\sqrt{N_{\mathbf{K}}} \max_{\mathbf{K}} |S_{\mathbf{K}}^\varepsilon| \leq x \right) dx.$$

Since

$$\mathbb{P} \left(\sqrt{N_{\mathbf{K}}} \max_{\mathbf{K}} |S_{\mathbf{K}}^\varepsilon| \leq x \right) = \left[\mathbb{P} \left(\sqrt{N_{\mathbf{K}}} |S_{\mathbf{K}}^\varepsilon| \leq x \right) \right]^{N(h)} = [F_{N_{\mathbf{K}}}(x)]^{N(h)},$$

we deduce that

$$\mathbb{E} \left(\alpha \frac{h^d}{\varepsilon^d} \frac{\lambda^2(\cdot, h, \varepsilon)}{\gamma^2} \right) \leq \int_0^\infty x^2 N(h) F_{N_{\mathbf{K}}}^{N(h)-1}(x) \varphi_{N_{\mathbf{K}}}(x) dx = e_1 + e_2 \quad (127)$$

where

$$e_1 = \int_0^{c_h} x^2 N(h) F_{N_{\mathbf{K}}}^{N(h)-1}(x) \varphi_{N_{\mathbf{K}}}(x) dx \quad \text{and} \quad e_2 = \int_{c_h}^\infty x^2 N(h) F_{N_{\mathbf{K}}}^{N(h)-1}(x) \varphi_{N_{\mathbf{K}}}(x) dx,$$

with $c_h = 2 \ln(N(h))$. We now successively bound from above e_1 and e_2 . First, integrating by part, and using that $0 \leq F_{N_{\mathbf{K}}} \leq 1$, we obtain

$$0 \leq e_1 = \left[x^2 F_{N_{\mathbf{K}}}^{N(h)}(x) \right]_{x=0}^{x=c_h} - \int_0^{c_h} 2x F_{N_{\mathbf{K}}}^{N(h)}(x) dx \leq c_h^2. \quad (128)$$

Second, again using $0 \leq F_{N_{\mathbf{K}}} \leq 1$, we get

$$0 \leq e_2 \leq \int_{c_h}^{\infty} x^2 N(h) \varphi_{N_{\mathbf{K}}}(x) dx = N(h) \mathbb{E} \left(\mathbf{1}_{\{\sqrt{N_{\mathbf{K}}}|S_{\mathbf{K}}^\varepsilon| > c_h\}} N_{\mathbf{K}} |S_{\mathbf{K}}^\varepsilon|^2 \right).$$

Using the Cauchy-Schwartz inequality, we obtain

$$e_2^2 \leq N(h)^2 \mathbb{E} [N_{\mathbf{K}}^2 |S_{\mathbf{K}}^\varepsilon|^4] \mathbb{P} \left(\sqrt{N_{\mathbf{K}}}|S_{\mathbf{K}}^\varepsilon| > c_h \right).$$

Introduce $Y_i = \frac{X_i - \mathbb{E}(X_0)}{\sqrt{\text{Var}(X_0)}}$, so that $S_{\mathbf{K}}^\varepsilon(\omega) = \frac{1}{N_{\mathbf{K}}} \sum_i Y_i(\omega)$. Recall now that $(Y_i)_{i \in \mathbb{Z}^d}$ is a sequence of independent identically distributed variables, with mean zero. Any such variables satisfy the bounds

$$\forall p \in \mathbb{N}^*, \quad \exists C_p > 0, \quad \forall N \in \mathbb{N}^*, \quad \left| \mathbb{E} \left[\left(\frac{1}{N} \sum_{i=1}^N Y_i \right)^{2p} \right] \right| \leq \frac{C_p}{N^p},$$

for a constant C_p that depends on p and the moments of Y_i , up to order $2p$. Recall that all moments of Y_i are well defined, as Y_i is bounded almost surely. Thus

$$e_2^2 \leq C_4 N(h)^2 \mathbb{P} \left(\sqrt{N_{\mathbf{K}}}|S_{\mathbf{K}}^\varepsilon| > c_h \right) \leq C_4 N(h)^2 \left[\mathbb{P} \left(\sqrt{N_{\mathbf{K}}} S_{\mathbf{K}}^\varepsilon > c_h \right) + \mathbb{P} \left(-\sqrt{N_{\mathbf{K}}} S_{\mathbf{K}}^\varepsilon > c_h \right) \right]. \quad (129)$$

We now recall the Markov inequality: for any positive non-decreasing function ψ on \mathbb{R} , and any real-valued random variable Z , we have

$$\forall b \in \mathbb{R}, \quad \mathbb{P}(Z \geq b) \leq \frac{\mathbb{E}(\psi(Z))}{\psi(b)}.$$

We apply this inequality to the random variable $Z(\omega) = \sqrt{N_{\mathbf{K}}} S_{\mathbf{K}}^\varepsilon(\omega)$, with $\psi = \exp(\cdot)$ for some $t \geq 0$, and $b = c_h$. This yields

$$\mathbb{P}(\sqrt{N_{\mathbf{K}}} S_{\mathbf{K}}^\varepsilon \geq c_h) \leq e^{-tc_h} \mathbb{E} \left[\exp \left(t \sqrt{N_{\mathbf{K}}} S_{\mathbf{K}}^\varepsilon \right) \right] \leq e^{-tc_h} \left[\mathbb{E} \left(\exp \left(\frac{t}{\sqrt{N_{\mathbf{K}}}} Y_0 \right) \right) \right]^{N_{\mathbf{K}}}, \quad (130)$$

where we have used the fact that $S_{\mathbf{K}}^\varepsilon$ is a sum of i.i.d. variables. Using a Taylor expansion with respect to t , we see that

$$\mathbb{E} \left[\exp \left(\frac{t}{\sqrt{N_{\mathbf{K}}}} Y_0 \right) \right] = 1 + \frac{t^2}{2N_{\mathbf{K}}} \mathbb{E}(Y_0^2) + \frac{1}{6N_{\mathbf{K}}^{3/2}} \mathbb{E} \left[Y_0^3 \exp(\xi Y_0 / \sqrt{N_{\mathbf{K}}}) \right] \quad \text{for some } \xi \in (0, t).$$

Thus

$$\left[\mathbb{E} \left(\exp \left(\frac{t}{\sqrt{N_{\mathbf{K}}}} Y_0 \right) \right) \right]^{N_{\mathbf{K}}} \leq \exp \left[\frac{t^2}{2} \mathbb{E}(Y_0^2) + \frac{1}{6\sqrt{N_{\mathbf{K}}}} \mathbb{E} \left(Y_0^3 \exp(\xi Y_0 / \sqrt{N_{\mathbf{K}}}) \right) \right].$$

Using (130), taking $t = 1$, and using that $e^{-c_h} = \frac{1}{N(h)^2}$, we obtain

$$\begin{aligned} \mathbb{P}(\sqrt{N_{\mathbf{K}}} S_{\mathbf{K}}^\varepsilon \geq c_h) &\leq \frac{1}{N(h)^2} \exp \left[\frac{1}{2} \mathbb{E}(Y_0^2) + \frac{1}{6\sqrt{N_{\mathbf{K}}}} \mathbb{E} \left(Y_0^3 \exp(\xi Y_0 / \sqrt{N_{\mathbf{K}}}) \right) \right] \quad \text{for some } \xi \in (0, 1), \\ &\leq \frac{1}{N(h)^2} \exp \left[\frac{1}{2} \mathbb{E}(Y_0^2) + \frac{1}{6} \mathbb{E}(|Y_0|^3 \exp(|Y_0|)) \right]. \end{aligned} \quad (131)$$

Likewise, considering $Z(\omega) = -\sqrt{N_{\mathbf{K}}} S_{\mathbf{K}}^\varepsilon(\omega)$, we obtain a similar bound. Collecting (129), (131) and the fact that Y_0 is bounded almost surely, we have

$$e_2^2 \leq C, \quad (132)$$

with C independent of h and ε . Collecting (127), (128) and (132), we get, for a constant C independent of h and ε ,

$$\mathbb{E}(\lambda(\cdot, h, \varepsilon)^2) \leq C \frac{\varepsilon^d}{h^d} [\ln(N(h))]^2.$$

This concludes the proof of Lemma 16. □

Remark 19. *The above proof shows that, when $\varepsilon \rightarrow 0$, the random variable $\left(\frac{h}{\varepsilon}\right)^{d/2} \lambda(\omega, h, \varepsilon)$ converges in law to $\mathcal{G}_h(\omega) = \max_{\mathbf{K}} |G_{\mathbf{K}}(\omega)|$, where $G_{\mathbf{K}}(\omega)$ are i.i.d. Gaussian random variables. Precise results on the behavior of $\mathcal{G}_h(\omega)$ when $h \rightarrow 0$ (i.e., when the number of Gaussian random variables involved diverges) are given in e.g. [46].*

Acknowledgements: Support from EOARD under Grant FA8655-10-C-4002 is gratefully acknowledged.

References

- [1] J. Aarnes and Y. R. Efendiev, *Mixed multiscale finite element methods for stochastic porous media flows*, SIAM Journal on Scientific Computing, 30(5):2319-2339, 2009.
- [2] G. Allaire and M. Amar, *Boundary layer tails in periodic homogenization*, Control, Optimization and Calculus of Variations, 4:209-243, 1999.
- [3] G. Allaire and R. Brizzi, *A multiscale finite element method for numerical homogenization*, SIAM Multiscale Modeling & Simulation, 4(3):790-812, 2005.
- [4] A. Anantharaman, Thèse de l'Université Paris Est, Ecole des Ponts, 2010.
<http://hal.archives-ouvertes.fr/tel-00558618>
- [5] A. Anantharaman, R. Costaouec, C. Le Bris, F. Legoll and F. Thomines, *Introduction to numerical stochastic homogenization and the related computational challenges: some recent developments*, in *Multiscale modeling and analysis for materials simulation*, W. Bao and Q. Du eds., Lecture Notes Series, Institute for Mathematical Sciences, National University of Singapore, vol. 22, 2011.
- [6] A. Anantharaman and C. Le Bris, *Homogénéisation d'un matériau périodique faiblement perturbé aléatoirement [Homogenization of a weakly randomly perturbed periodic material]*, C. R. Acad. Sci. Série I, 348:529-534, 2010.
- [7] A. Anantharaman and C. Le Bris, *A numerical approach related to defect-type theories for some weakly random problems in homogenization*, SIAM Multiscale Modeling & Simulation, 9(2):513-544, 2011.
- [8] A. Anantharaman and C. Le Bris, *Elements of mathematical foundations for a numerical approach for weakly random homogenization problems*, Communications in Computational Physics, in press, preprint available at <http://arxiv.org/abs/1005.3922>.
- [9] M. Avellaneda and F. H. Lin, *Compactness methods in the theory of homogenization*, Comm. Pure Appl. Math., 40:803-847, 1987.
- [10] G. Bal, J. Garnier, S. Motsch and V. Perrier, *Random integrals and correctors in homogenization*, Asymptot. Anal., 59(1-2):1-26, 2008.
- [11] G. Bal and W. Jing, *Corrector theory for MsFEM and HMM in random media*, preprint available at <http://www.columbia.edu/~gb2030/PAPERS/MsFEM-HMM.pdf>.
- [12] A. Bensoussan, J.-L. Lions and G. Papanicolaou, *Asymptotic analysis for periodic structures*, Studies in Mathematics and its Applications, 5. North-Holland Publishing Co., Amsterdam-New York, 1978.
- [13] X. Blanc, R. Costaouec, C. Le Bris and F. Legoll, *Variance reduction in stochastic homogenization using antithetic variables*, Markov Processes and Related Fields, to appear, preprint available at <http://cermics.enpc.fr/~legoll/hdr/FL24.pdf>
- [14] X. Blanc, C. Le Bris and P.-L. Lions, *Une variante de la théorie de l'homogénéisation stochastique des opérateurs elliptiques [A variant of stochastic homogenization theory for elliptic operators]*, C. R. Acad. Sci. Série I, 343:717-724, 2006.

- [15] X. Blanc, C. Le Bris and P.-L. Lions, *Stochastic homogenization and random lattices*, Journal de Mathématiques Pures et Appliquées, 88:34-63, 2007.
- [16] A. Bourgeat and A. Piatnitski, *Estimates in probability of the residual between the random and the homogenized solutions of one-dimensional second-order operator*, Asymptot. Anal., 21:303-315, 1999.
- [17] L. Carballal Perdiz, P. Degond, F. Deluzet, R. Loubère, A. Lozinski and J.-M. Rovarch, *Multiscale finite element method for perforated domains*, in preparation.
- [18] Z. Chen, *Multiscale methods for elliptic homogenization problems*, Numer. Methods Partial Differential Eq., 22(2):317-360, 2006.
- [19] Z. Chen, M. Cui, T. Y. Savchuk and X. Yu, *The multiscale finite element method with nonconforming elements for elliptic homogenization problems*, SIAM Multiscale Modeling & Simulation, 7(2):517-538, 2008.
- [20] Z. Chen and T. Y. Hou, *A mixed multiscale finite element method for elliptic problems with oscillating coefficients*, Math. Comp., 72(242):541-576, 2002.
- [21] Z. Chen and T. Y. Savchuk, *Analysis of the multiscale finite element method for nonlinear and random homogenization problems*, SIAM Journal on Numerical Analysis, 46(1):260-279, 2008.
- [22] P. G. Ciarlet, *The finite element method for elliptic problems*, North-Holland, 1978.
- [23] D. Cioranescu and P. Donato, *An introduction to homogenization*. Oxford Lecture Series in Mathematics and its Applications, 17. The Clarendon Press, Oxford University Press, New York, 1999.
- [24] R. Costaouec, *Asymptotic expansion of the homogenized matrix in two weakly stochastic homogenization settings*, Appl. Math. Res. Express, 2011, available at <http://amrx.oxfordjournals.org/content/early/2011/10/28/amrx.abr011.short?rss=1>
- [25] R. Costaouec, C. Le Bris and F. Legoll, *Approximation numérique d'une classe de problèmes en homogénéisation stochastique [Numerical approximation of a class of problems in stochastic homogenization]*, C. R. Acad. Sci. Série I, 348:99-103, 2010.
- [26] P. Dostert, Y. R. Efendiev and T. Y. Hou, *Multiscale finite element methods for stochastic porous media flow equations and application to uncertainty quantification*, Computer Methods in Applied Mechanics and Engineering, 197(43-44):3445-3455, 2008.
- [27] W. E and B. Engquist, *The heterogeneous multiscale methods*, Comm. Math. Sci., 1(1):87-132, 2003.
- [28] W. E and B. Engquist, *The Heterogeneous Multiscale Method for homogenization problems*, in *Multiscale Methods in Science and Engineering*, pp. 89-110, Lect. Notes in Comput. Sci. Eng., vol. 44, Springer, Berlin, 2005.
- [29] W. E, B. Engquist, X. Li, W. Ren and E. Vanden-Eijnden, *Heterogeneous multiscale methods: a review*, Comm. Comput. Phys., 2(3):367-450, 2007.
- [30] Y. R. Efendiev and T. Y. Hou, *Multiscale finite element methods: theory and applications*, Surveys and tutorials in the applied mathematical sciences, Springer, New York, 2009.
- [31] Y. R. Efendiev, T. Y. Hou and V. Ginting, *Multiscale finite element methods for nonlinear problems and their applications*, Communications in Mathematical Sciences, 2(4):553-589, 2004.
- [32] Y. R. Efendiev, T. Y. Hou and X.-H. Wu, *Convergence of a nonconforming multiscale finite element method*, SIAM Journal on Numerical Analysis, 37(3):888-910, 2000.
- [33] FreeFEM, <http://www.freefem.org>
- [34] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, reprint of the 1998 ed., Classics in Mathematics, Springer, 2001.
- [35] V. Ginting, A. Malqvist and M. Presho, *A novel method for solving multiscale elliptic problems with randomly perturbed data*, SIAM Multiscale Modeling & Simulation, 8(3):977-996, 2010.

- [36] A. Gloria, *An analytical framework for numerical homogenization. Part II: Windowing and oversampling*, SIAM Multiscale Modeling & Simulation, 7(1):274-293, 2008.
- [37] T. Y. Hou and X.-H. Wu, *A multiscale finite element method for elliptic problems in composite materials and porous media*, Journal of Computational Physics, 134(1):169-189, 1997.
- [38] T. Y. Hou, X.-H. Wu and Z. Cai, *Convergence of a multiscale finite element method for elliptic problems with rapidly oscillating coefficients*, Math. Comp., 68(227):913-943, 1999.
- [39] T. Y. Hou, X.-H. Wu and Y. Zhang, *Removing the cell resonance error in the multiscale finite element method via a Petrov-Galerkin formulation*, Communications in Mathematical Sciences, 2(2):185-205, 2004.
- [40] V. V. Jikov, S. M. Kozlov and O. A. Oleinik, *Homogenization of differential operators and integral functionals*, Springer-Verlag, 1994.
- [41] C. Le Bris, *Some numerical approaches for “weakly” random homogenization*, Proceedings of ENUMATH 2009, Springer Lecture Notes in Computational Science and Engineering, G. Kreiss et al. (eds.), Numerical Mathematics and Advanced Applications 2009, p. 29-45, 2010.
- [42] C. Le Bris, F. Legoll and F. Thomines, *Rate of convergence of a two-scale expansion for some weakly stochastic homogenization problems*, submitted to Asymptotic Analysis, preprint available at <http://arxiv.org/abs/1110.5206>
- [43] F. Legoll and F. Thomines, *Convergence of the residual process of a variant of stochastic homogenization in dimension one*, in preparation.
- [44] A. Lozinski, *Habilitation à Diriger des Recherches*, Université Paul Sabatier (Toulouse), 2010.
<http://www.math.univ-toulouse.fr/~lozinski/hdr.pdf>
- [45] Y. Maday, *Reduced basis method for the rapid and reliable solution of partial differential equations*, in International Congress of Mathematicians, vol. III, pp. 1255-1270, Eur. Math. Soc., Zürich, 2006.
- [46] Y. Mittal, *Limiting behavior of maxima in stationary Gaussian sequences*, The Annals of Probability, 2(2):231-242, 1974.
- [47] G. C. Papanicolaou and S. R. S. Varadhan, *Boundary value problems with rapidly oscillating random coefficients*, in Proc. Colloq. on Random Fields: Rigorous Results in Statistical Mechanics and Quantum Field Theory, 1979 (J. Fritz, J. L. Lebaritz and D. Szasz, eds.), Vol. 10 of Colloquia Mathematica Societ. Janos Bolyai, pp. 835-873, North-Holland, 1981.
- [48] L. Tartar, *Estimations of homogenized coefficients*, in *Topics in the mathematical modelling of composite materials*, A. Cherkaev and R. Kohn eds., Progress in nonlinear differential equations and their applications, vol. 31, Birkhäuser, 1987.