

DERIVATION OF OROWAN'S LAW FROM THE PEIERLS-NABARRO MODEL

RÉGIS MONNEAU

Université Paris-Est, CERMICS, Ecole des Ponts ParisTech,
6-8 avenue Blaise Pascal, Cité Descartes, Champs sur Marne,
77455 Marne la Vallée Cedex 2, France

STEFANIA PATRIZI

Instituto Superior Técnico, Dep. de Matemática
Av. Rovisco Pais Lisboa, Portugal

ABSTRACT. In this paper we consider the time dependent Peierls-Nabarro model in dimension one. This model is a semi-linear integro-differential equation associated to the half Laplacian. This model describes the evolution of phase transitions associated to dislocations. At large scale with well separated dislocations, we show that the dislocations move at a velocity proportional to the effective stress. This implies Orowan's law which claims that the plastic strain velocity is proportional to the product of the density of dislocations by the effective stress.

1. INTRODUCTION

1.1. Setting of the problem. In this paper we consider a one-dimensional Peierls-Nabarro model, describing the motion of dislocations in crystals. In this model dislocations can be seen as phase transitions of a function u^ϵ solving the following equation for $\epsilon = 1$

$$(1.1) \quad \begin{cases} \partial_t u^\epsilon = \mathcal{I}_1[u^\epsilon(t, \cdot)] - W' \left(\frac{u^\epsilon}{\epsilon} \right) & \text{in } \mathbb{R}^+ \times \mathbb{R} \\ u^\epsilon(0, x) = u_0(x) & \text{on } \mathbb{R}. \end{cases}$$

Here $\mathcal{I}_1 = -(-\Delta)^{\frac{1}{2}}$ is the half Laplacian whose expression will be made precise later in (1.8) and W is a one periodic potential which describes the misfit of atoms in the crystal created by the presence of dislocations. Equation (1.1) models the dynamics of parallel straight edge dislocation lines in the same slip plane with the same Burgers vector, moving with self-interactions. In other words equation (1.1) simply describes the motion of dislocations by relaxation of the total energy (elastic + misfit). For a physical introduction to the Peierls-Nabarro model, see for instance [9], [14]; we also refer the reader to the paper of Nabarro [13] which presents an historical tour on the Peierls-Nabarro model. The Peierls-Nabarro model has been originally introduced as a variational (stationary) model (see [13]). The model considered in the present paper, i.e. the time evolution Peierls-Nabarro model as a gradient flow dynamics has only been introduced quite recently, see for instance [12] and [4], and [11] where this model is also presented. See also the paper [3] that initiated several other works about jump-diffusion reaction equations.

In [11] we study the limit as $\epsilon \rightarrow 0$ of the viscosity solution u^ϵ of (1.1) in higher dimensions and with additional periodic terms. Under certain assumptions, we show in particular that u^ϵ converges to the solution of the following equation:

$$(1.2) \quad \begin{cases} \partial_t u = \overline{H}(u_x, \mathcal{I}_1[u(t, \cdot)]) & \text{in } \mathbb{R}^+ \times \mathbb{R} \\ u(0, x) = u_0(x) & \text{on } \mathbb{R}. \end{cases}$$

In mechanics, equation (1.2) can be interpreted as a plastic flow rule, which expresses the plastic strain velocity $\partial_t u$ as a function \overline{H} of the dislocation density u_x and the effective stress $\mathcal{I}_1[u]$ created by the density of dislocations. Mathematically the function \overline{H} , usually called *effective Hamiltonian*, is determined by the following auxiliary problem:

$$(1.3) \quad \begin{cases} \partial_\tau v = L + \mathcal{I}_1[v(\tau, \cdot)] - W'(v) & \text{in } \mathbb{R}^+ \times \mathbb{R} \\ v(0, y) = py & \text{on } \mathbb{R}. \end{cases}$$

Here the quantity L appears to be an additional constant stress field. Indeed, we have

Theorem 1.1 (Theorem 1.1, [11]). *Assume that $W \in C^{1,1}(\mathbb{R})$ and W is 1-periodic. For every $L \in \mathbb{R}$ and $p \in \mathbb{R}$, there exists a unique viscosity solution $v \in C(\mathbb{R}^+ \times \mathbb{R})$ of (1.3) and there exists a unique $\lambda \in \mathbb{R}$ such that $v - py - \lambda\tau$ is bounded in $\mathbb{R}^+ \times \mathbb{R}$. The real number λ is denoted by $\overline{H}(p, L)$. The function $\overline{H}(p, L)$ is continuous on \mathbb{R}^2 and non-decreasing in L .*

This is the starting point of this paper. Our goal is to study the behaviour of $\overline{H}(p, L)$ for small p and L , and in this regime to recover Orowan's law, which claims that

$$(1.4) \quad \overline{H}(p, L) \simeq c_0 |p| L$$

for some constant of proportionality $c_0 > 0$.

1.2. Main result. In order to describe our main result, we need the following assumptions on the potential W :

$$(1.5) \quad \begin{cases} W \in C^{4,\beta}(\mathbb{R}) & \text{for some } 0 < \beta < 1 \\ W(v+1) = W(v) & \text{for any } v \in \mathbb{R} \\ W = 0 & \text{on } \mathbb{Z} \\ W > 0 & \text{on } \mathbb{R} \setminus \mathbb{Z} \\ \alpha = W''(0) > 0. \end{cases}$$

Under (1.5), it is in particular known (see Cabré and Solà-Morales [2]) that there exists a unique function ϕ solution of

$$(1.6) \quad \begin{cases} \mathcal{I}_1[\phi] = W'(\phi) & \text{in } \mathbb{R} \\ \phi' > 0 & \text{in } \mathbb{R} \\ \lim_{x \rightarrow -\infty} \phi(x) = 0, \quad \lim_{x \rightarrow +\infty} \phi(x) = 1, \quad \phi(0) = \frac{1}{2}. \end{cases}$$

Our main result is the following:

Theorem 1.2 (Orowan's law). *Assume (1.5) and let $p_0, L_0 \in \mathbb{R}$. Then the function \overline{H} defined in Theorem 1.1 satisfies*

$$(1.7) \quad \frac{\overline{H}(\delta p_0, \delta L_0)}{\delta^2} \rightarrow c_0 |p_0| L_0 \quad \text{as } \delta \rightarrow 0^+ \quad \text{with} \quad c_0 = \left(\int_{\mathbb{R}} (\phi')^2 \right)^{-1}.$$

Theorem 1.2 shows that in the limit of small density of dislocations p and small stress L , the effective Hamiltonian \overline{H} follows Orowan's law (1.4). This implies that in this regime, the plastic strain velocity $\partial_t u$ in (1.2) is proportional to the dislocation density $|u_x|$ times the effective stress $\mathcal{I}_1[u]$, i.e.

$$\partial_t u \simeq c_0 |u_x| \mathcal{I}_1[u(t, \cdot)].$$

Notice that this last equation has been proposed by Head [8] and self-similar solutions have been studied mathematically in [1].

Notice that in homogenization problems the effective Hamiltonian is usually unknown. Explicit formulas for \overline{H} are known only in very special cases, see for instance [10]. The result of Theorem 1.2 provides an other example of explicit expression for a particular homogenization problem.

Finally we give the precise expression (the Lévy-Khintchine formula in Thm 1 of [5]) of the Lévy operator \mathcal{I}_1 of order 1. For bounded C^2 - functions U and for $r > 0$, we set

$$(1.8) \quad \begin{aligned} \mathcal{I}_1[U](x) = \mathcal{I}_1[U, x] &= \int_{|z| \leq r} (U(x+z) - U(x) - \frac{dU(x)}{dx} \cdot z) \mu(dz) \\ &+ \int_{|z| > r} (U(x+z) - U(x)) \mu(dz), \quad \text{with} \quad \mu(dz) = \frac{1}{\pi} \frac{dz}{z^2}. \end{aligned}$$

Notice that this expression is independent on the choice of $r > 0$, because of the antisymmetry of $z\mu(dz)$. More generally, when U is C^2 such that $U - \ell$ is bounded with ℓ a linear function, we simply define

$$\mathcal{I}_1[U](x) = \mathcal{I}_1[U, x] = \lim_{r \rightarrow 0^+} \int_{r < |z| < 1/r} (U(x+z) - U(x)) \mu(dz)$$

1.3. Organization of the article.

In Section 2, we present the main ideas which allow us to prove Orowan's law and give the proof of the main theorem (Theorem 1.2). This proof is based on Proposition 2.1 which claims asymptotics satisfied by a good Ansatz (see (2.4)). The remaining part of the paper is then devoted to the proof of Proposition 2.1. In Section 3, we recall in Lemmata 3.1 and 3.2, useful asymptotics respectively on the transition layer ϕ and some corrector ψ . The main result of this section is some asymptotics on the non linear PDE evaluated on the Ansatz. In Section 4, we do the proof of Proposition 2.1. Finally in an appendix (Section 5), we give the proof of Lemmata 3.1 and 3.2. We also give the proof of five claims used in Section 3 and a technical lemma (Lemma 4.1) used in Section 4.

2. IDEAS AND PROOF OF OROWAN'S LAW (THEOREM 1.2)

2.1. Heuristic for the proof of Orowan's law.

The idea underlying the proof of Orowan's law is related to a fine asymptotics of equation (1.3). It is also known (see [7]) that if v solves (1.3) with $L = \delta L_0$, i.e.

$$(2.1) \quad \partial_\tau v = \delta L_0 + \mathcal{I}_1[v(\tau, \cdot)] - W'(v)$$

for a choice of initial data with a finite number of indices i :

$$v(0, y) = \frac{\delta L_0}{\alpha} + \sum_{x_i^0 \geq 0} \phi\left(y - \frac{x_i^0}{\delta}\right) + \sum_{x_i^0 < 0} \left(\phi\left(y - \frac{x_i^0}{\delta}\right) - 1\right)$$

where $\alpha = W''(0) > 0$ (defined in (1.5)), then

$$v^\delta(t, x) = v\left(\frac{t}{\delta^2}, \frac{x}{\delta}\right) \rightarrow v^0(t, x) = \sum_{x_i^0 \geq 0} H(x - x_i(t)) + \sum_{x_i^0 < 0} (H(x - x_i(t)) - 1) \quad \text{as } \delta \rightarrow 0$$

where H is the Heaviside function and with the dynamics

$$(2.2) \quad \begin{cases} \frac{dx_i}{dt} = c_0 \left(-L_0 + \frac{1}{\pi} \sum_{j \neq i} \frac{1}{x_i - x_j}\right) \\ x_i(0) = x_i^0. \end{cases}$$

Moreover for the choice $p = \delta p_0$ with $p_0 > 0$ and $x_i^0 = i/p_0$ that we extend formally for all $i \in \mathbb{Z}$, we see (at least formally) that

$$|v(0, y) - \delta p_0 y| \leq C_\delta.$$

This suggests also that the infinite sum in (2.2) should vanish (by antisymmetry) and then the mean velocity should be

$$\frac{dx_i}{dt} \simeq -c_0 L_0$$

i.e., after scaling back

$$v(\tau, y) \simeq \delta p_0 (y - c_1 \tau) + \text{bounded}$$

with the velocity

$$c_1 = \frac{d(x_i/\delta)}{d(t/\delta^2)} \simeq -c_0 L_0 \delta$$

i.e.,

$$v(\tau, y) \simeq \delta p_0 y + \lambda \tau + \text{bounded} \quad \text{with } \lambda \simeq \delta^2 c_0 p_0 L_0.$$

We deduce that we should have

$$\frac{v(\tau, y)}{\tau} \rightarrow \lambda \simeq \delta^2 c_0 p_0 L_0 \quad \text{as } \tau \rightarrow +\infty.$$

We see that this $\lambda = \overline{H}(\delta p_0, \delta L_0)$ is exactly the one we expect asymptotically in Theorem 1.2 when $p_0 > 0$.

2.2. The ansatz used in the proofs.

In the spirit of [6], one may expect to find particular solutions v of (2.1) that we can write

$$v(\tau, y) = h(\delta p_0 y + \lambda \tau)$$

for some $\lambda \in \mathbb{R}$ and a function h (called hull function) satisfying

$$|h(z) - z| \leq C.$$

This means that h solves

$$\lambda h' = \delta L_0 + \delta |p_0| \mathcal{I}_1[h] - W'(h).$$

Then it is natural to introduce the non linear operator:

$$(2.3) \quad NL_{L_0}^\lambda[h] := \lambda h' - \delta L_0 - \delta |p_0| \mathcal{I}_1[h] + W'(h)$$

and for the ansatz for λ :

$$\bar{\lambda}_\delta^{L_0} = \delta^2 c_0 |p_0| L_0$$

it is natural to look for an ansatz $h_\delta^{L_0}$ for h . The answer is indicated by the heuristic of subsection 2.1. Indeed we define (see Proposition 2.1)

$$h_\delta^{L_0}(x) = \lim_{n \rightarrow +\infty} s_{\delta,n}^{L_0}(x)$$

where for all $p_0 \neq 0$, $L_0 \in \mathbb{R}$, $\delta > 0$ and $n \in \mathbb{N}$ we define the sequence of functions $\{s_{\delta,n}^{L_0}(x)\}_n$ by

$$(2.4) \quad s_{\delta,n}^{L_0}(x) = \frac{\delta L_0}{\alpha} + \sum_{i=-n}^n \left[\phi \left(\frac{x-i}{\delta |p_0|} \right) + \delta \psi \left(\frac{x-i}{\delta |p_0|} \right) \right] - n$$

where $\alpha = W''(0) > 0$, ϕ is the solution of (1.6) and the corrector ψ is the solution of the following problem

$$(2.5) \quad \begin{cases} \mathcal{I}_1[\psi] = W''(\phi)\psi + \frac{L_0}{W''(0)}(W''(\phi) - W''(0)) + c\phi' & \text{in } \mathbb{R} \\ \lim_{x \rightarrow +\infty} \psi(x) = 0 \\ c = \frac{L_0}{\int_{\mathbb{R}} (\phi')^2}. \end{cases}$$

From [7], it is known that there exists a unique ψ solution of (2.5). Moreover this corrector ψ has been introduced naturally in [7] in order to perform part of the analysis presented in the heuristic (subsection 2.1), and this is then natural to use it here in our ansatz. We will prove later the following result which justifies that the ansatz is indeed a good ansatz as expected.

Proposition 2.1. (Good ansatz)

Assume (1.5). For any $L \in \mathbb{R}$, $\delta > 0$ and $x \in \mathbb{R}$, there exists the finite limit

$$h_\delta^L(x) = \lim_{n \rightarrow +\infty} s_{\delta,n}^L(x).$$

Moreover h_δ^L has the following properties:

- (i) $h_\delta^L \in C^2(\mathbb{R})$ and satisfies

$$NL_L^{\bar{\lambda}_\delta^L}[h_\delta^L](x) = o(\delta),$$

where $\lim_{\delta \rightarrow 0} \frac{o(\delta)}{\delta} = 0$, uniformly for $x \in \mathbb{R}$ and locally uniformly in $L \in \mathbb{R}$; Here

$$\bar{\lambda}_\delta^L = \delta^2 c_0 |p_0| L$$

and NL_L^λ is defined in (2.3).

- (ii) There exists a constant $C > 0$ such that $|h_\delta^L(x) - x| \leq C$ for any $x \in \mathbb{R}$.

2.3. Proof of Theorem 1.2. We will show that Theorem 1.2 follows from Proposition 2.1, and the comparison principle.

Fix $\eta > 0$ and let $L = L_0 - \eta$. By (i) of Proposition 2.1, there exists $\delta_0 = \delta_0(\eta) > 0$ such that for any $\delta \in (0, \delta_0)$ we have

$$(2.6) \quad NL_{L_0}^{\bar{\lambda}_\delta^L}[h_\delta^L] = NL_L^{\bar{\lambda}_\delta^L}[h_\delta^L] - \delta\eta < 0 \quad \text{in } \mathbb{R}.$$

Let us consider the function $\tilde{v}(\tau, y)$, defined by

$$\tilde{v}(\tau, y) = h_\delta^L(\delta p_0 y + \bar{\lambda}_\delta^L \tau).$$

By (ii) of Proposition 2.1, we have

$$(2.7) \quad |\tilde{v}(\tau, y) - \delta p_0 y - \bar{\lambda}_\delta^L \tau| \leq \lceil C \rceil,$$

where $\lceil C \rceil$ is the ceil integer part of C . Moreover, by (2.6) and (2.7), \tilde{v} satisfies

$$\begin{cases} \tilde{v}_\tau \leq \delta L_0 + \mathcal{I}_1[\tilde{v}] - W'(\tilde{v}) & \text{in } \mathbb{R}^+ \times \mathbb{R} \\ \tilde{v}(0, y) \leq \delta p_0 y + \lceil C \rceil & \text{on } \mathbb{R}. \end{cases}$$

Let $v(\tau, y)$ be the solution of (1.3), with $p = \delta p_0$ and $L = \delta L_0$, whose existence is ensured by Theorem 1.1. Then from the comparison principle and the periodicity of W , we deduce that

$$\tilde{v}(\tau, y) \leq v(\tau, y) + \lceil C \rceil.$$

By the previous inequality and (2.7), we get

$$\bar{\lambda}_\delta^L \tau \leq v(\tau, y) - \delta p_0 y + 2\lceil C \rceil,$$

and dividing by τ and letting τ go to $+\infty$, we finally obtain

$$\delta^2 c_0 |p_0| (L_0 - \eta) = \bar{\lambda}_\delta^L \leq \bar{H}(\delta p_0, \delta L_0).$$

Similarly, it is possible to show that

$$\bar{H}(\delta p_0, \delta L_0) \leq \delta^2 c_0 |p_0| (L_0 + \eta).$$

We have proved that for any $\eta > 0$ there exists $\delta_0 = \delta_0(\eta) > 0$ such that for any $\delta \in (0, \delta_0)$ we have

$$\left| \frac{\bar{H}(\delta p_0, \delta L_0)}{\delta^2} - c_0 |p_0| L_0 \right| \leq c_0 |p_0| \eta,$$

i.e. (1.7), as desired. □

3. PRELIMINARY ASYMPTOTICS

The main goal of this section is to show Lemma 3.3 which is a first result in the direction of Proposition 2.1. We start with preliminary results in a first subsection and prove Lemma 3.3 in the second subsection.

3.1. Preliminary results.

On the function W , we assume (1.5). Then there exists a unique solution of (1.6) which is of class $C^{2,\beta}$, as shown by Cabré and Solà-Morales in [2]. Under (1.5), the existence of a solution of class $C^{1,\beta}$ of the problem (2.5) is proved by González and Monneau in [7]. Actually, the regularity of W implies, that $\phi \in C^{4,\beta}(\mathbb{R})$ and $\psi \in C^{3,\beta}(\mathbb{R})$, see Lemma 2.3 in [2].

To prove Proposition 2.1 we need several preliminary results. We first state the following two lemmata about the behavior of the functions ϕ and ψ at infinity. We denote by $H(x)$ the Heaviside function defined by

$$H(x) = \begin{cases} 1 & \text{for } x \geq 0 \\ 0 & \text{for } x < 0. \end{cases}$$

Then we have

Lemma 3.1 (Behavior of ϕ). *Assume (1.5). Let ϕ be the solution of (1.6), then there exist constants $K_0, K_1 > 0$ such that*

$$(3.1) \quad \left| \phi(x) - H(x) + \frac{1}{\alpha\pi x} \right| \leq \frac{K_1}{x^2}, \quad \text{for } |x| \geq 1,$$

and for any $x \in \mathbb{R}$

$$(3.2) \quad 0 < \frac{K_0}{1+x^2} \leq \phi'(x) \leq \frac{K_1}{1+x^2},$$

$$(3.3) \quad -\frac{K_1}{1+x^2} \leq \phi''(x) \leq \frac{K_1}{1+x^2},$$

$$(3.4) \quad -\frac{K_1}{1+x^2} \leq \phi'''(x) \leq \frac{K_1}{1+x^2}.$$

Lemma 3.2 (Behavior of ψ). *Assume (1.5). Let ψ be the solution of (2.5), then for any $L \in \mathbb{R}$ there exist constants K_2 and K_3 , with $K_3 > 0$, depending on L such that*

$$(3.5) \quad \left| \psi(x) - \frac{K_2}{x} \right| \leq \frac{K_3}{x^2}, \quad \text{for } |x| \geq 1,$$

and for any $x \in \mathbb{R}$

$$(3.6) \quad -\frac{K_3}{1+x^2} \leq \psi'(x) \leq \frac{K_3}{1+x^2},$$

$$(3.7) \quad -\frac{K_3}{1+x^2} \leq \psi''(x) \leq \frac{K_3}{1+x^2}.$$

We postpone the proof of the two lemmata in the appendix (Section 5). For simplicity of notation we denote (for the rest of the paper)

$$x_i = \frac{x-i}{\delta|p_0|}, \quad \tilde{\phi}(z) = \phi(z) - H(z), \quad \mathcal{I}_1[\phi, x_i] = \mathcal{I}_1[\phi](x_i).$$

Then we have the following five claims (whose proofs are also postponed in the appendix (Section 5)).

Claim 1: *Let $x = i_0 + \gamma$, with $i_0 \in \mathbb{Z}$ and $\gamma \in (-\frac{1}{2}, \frac{1}{2}]$, then*

$$\sum_{\substack{i=-n \\ i \neq i_0}}^n \frac{1}{x-i} \rightarrow -2\gamma \sum_{i=1}^{+\infty} \frac{1}{i^2 - \gamma^2} \quad \text{as } n \rightarrow +\infty,$$

$$\sum_{i=-n}^{i_0-1} \frac{1}{(x-i)^2} \rightarrow \sum_{i=1}^{+\infty} \frac{1}{(i+\gamma)^2} \quad \text{as } n \rightarrow +\infty,$$

$$\sum_{i=i_0+1}^n \frac{1}{(x-i)^2} \rightarrow \sum_{i=1}^{+\infty} \frac{1}{(i-\gamma)^2} \quad \text{as } n \rightarrow +\infty.$$

Claim 2: *For any $x \in \mathbb{R}$ the sequence $\{s_{\delta,n}^L(x)\}_n$ converges as $n \rightarrow +\infty$.*

Claim 3: *The sequence $\{(s_{\delta,n}^L)'\}_n$ converges on \mathbb{R} as $n \rightarrow +\infty$, uniformly on compact sets.*

Claim 4: *The sequence $\{(s_{\delta,n}^L)''\}_n$ converges on \mathbb{R} as $n \rightarrow +\infty$, uniformly on compact sets.*

Claim 5: *For any $x \in \mathbb{R}$ the sequences $\sum_{i=-n}^n \mathcal{I}_1[\phi, x_i]$ and $\sum_{i=-n}^n \mathcal{I}_1[\psi, x_i]$ converge as $n \rightarrow +\infty$.*

3.2. First asymptotics.

In order to do the proof of Proposition 2.1, we first get the following result:

Lemma 3.3. (First asymptotics) *We have*

$$-C\delta^2 \leq \lim_{n \rightarrow +\infty} NL_L^{\bar{\lambda}_\delta^L} [s_{\delta,n}^L](x) \leq C\delta^2,$$

where C is independent of x .

Proof of Lemma 3.3.

Step 1: First computation

Fix $x \in \mathbb{R}$, let $i_0 \in \mathbb{Z}$ and $\gamma \in (-\frac{1}{2}, \frac{1}{2}]$ be such that $x = i_0 + \gamma$, let $\frac{1}{\delta|p_0|} \geq 2$ and $n > |i_0|$. Then we have

$$\begin{aligned}
A &:= NL_L^{\bar{\lambda}_\delta^L} [s_{\delta,n}^L](x) \\
&= \frac{\bar{\lambda}_\delta^L}{\delta|p_0|} \sum_{i=-n}^n [\phi'(x_i) + \delta\psi'(x_i)] - \sum_{i=-n}^n [\mathcal{I}_1[\phi, x_i] + \delta\mathcal{I}_1[\psi, x_i]] \\
&\quad + W' \left(\frac{L\delta}{\alpha} + \sum_{i=-n}^n [\phi(x_i) + \delta\psi(x_i)] \right) - \delta L
\end{aligned}$$

where we have used the definitions and the periodicity of W . Using the equation (1.6) satisfied by ϕ , we can rewrite it as

$$\begin{aligned}
A &= \frac{\bar{\lambda}_\delta^L}{\delta|p_0|} \left\{ \phi'(x_{i_0}) + \delta\psi'(x_{i_0}) + \sum_{\substack{i=-n \\ i \neq i_0}}^n [\phi'(x_i) + \delta\psi'(x_i)] \right\} - \sum_{\substack{i=-n \\ i \neq i_0}}^n W'(\tilde{\phi}(x_i)) - \delta\mathcal{I}_1[\psi, x_{i_0}] \\
&\quad - \delta \sum_{\substack{i=-n \\ i \neq i_0}}^n \mathcal{I}_1[\psi, x_i] + W' \left(\frac{L\delta}{\alpha} + \sum_{i=-n}^n [\tilde{\phi}(x_i) + \delta\psi(x_i)] \right) - W'(\tilde{\phi}(x_{i_0})) - \delta L
\end{aligned}$$

Using the definition of $\bar{\lambda}_\delta^L$ and a Taylor expansion of W' , we get

$$\begin{aligned}
A &= \delta c_0 L \left\{ \phi'(x_{i_0}) + \delta\psi'(x_{i_0}) + \sum_{\substack{i=-n \\ i \neq i_0}}^n [\phi'(x_i) + \delta\psi'(x_i)] \right\} - W''(0) \sum_{\substack{i=-n \\ i \neq i_0}}^n \tilde{\phi}(x_i) - \delta \sum_{\substack{i=-n \\ i \neq i_0}}^n \mathcal{I}_1[\psi, x_i] \\
&\quad - \delta\mathcal{I}_1[\psi, x_{i_0}] + W''(\phi(x_{i_0})) \left(\frac{L\delta}{\alpha} + \delta\psi(x_{i_0}) + \sum_{\substack{i=-n \\ i \neq i_0}}^n [\tilde{\phi}(x_i) + \delta\psi(x_i)] \right) - \delta L + E
\end{aligned}$$

with the error term

$$E = \sum_{\substack{i=-n \\ i \neq i_0}}^n O(\tilde{\phi}(x_i))^2 + O \left(\frac{L\delta}{\alpha} + \delta\psi(x_{i_0}) + \sum_{\substack{i=-n \\ i \neq i_0}}^n [\tilde{\phi}(x_i) + \delta\psi(x_i)] \right)^2$$

Simply reorganizing the terms, we get with $c = c_0 L$:

$$\begin{aligned}
A = & \delta c_0 L \left\{ \delta \psi'(x_{i_0}) + \sum_{\substack{i=-n \\ i \neq i_0}}^n [\phi'(x_i) + \delta \psi'(x_i)] \right\} - W''(0) \sum_{\substack{i=-n \\ i \neq i_0}}^n \tilde{\phi}(x_i) - \delta \sum_{\substack{i=-n \\ i \neq i_0}}^n \mathcal{I}_1[\psi, x_i] \\
& + W''(\phi(x_{i_0})) \left(\sum_{\substack{i=-n \\ i \neq i_0}}^n [\tilde{\phi}(x_i) + \delta \psi(x_i)] \right) \\
& + \delta \left(-\mathcal{I}_1[\psi, x_{i_0}] + W''(\phi(x_{i_0}))\psi(x_{i_0}) + \frac{L}{\alpha} W''(\phi(x_{i_0})) - L + c\phi'(x_{i_0}) \right) + E
\end{aligned}$$

Using equation (2.5) satisfied by ψ , we get

$$\begin{aligned}
A = & \delta c_0 L \left\{ \delta \psi'(x_{i_0}) + \sum_{\substack{i=-n \\ i \neq i_0}}^n [\phi'(x_i) + \delta \psi'(x_i)] \right\} + (W''(\phi(x_{i_0})) - W''(0)) \sum_{\substack{i=-n \\ i \neq i_0}}^n \tilde{\phi}(x_i) \\
& - \delta \sum_{\substack{i=-n \\ i \neq i_0}}^n \mathcal{I}_1[\psi, x_i] + W''(\phi(x_{i_0}))\delta \sum_{\substack{i=-n \\ i \neq i_0}}^n \psi(x_i) + E
\end{aligned}$$

Step 2: Bound on $\sum_{\substack{i=-n \\ i \neq i_0}}^n [\phi'(x_i) + \delta \psi'(x_i)]$

Let us bound the second term of the last equality, uniformly in x . From (3.2) and (3.6) it follows that

$$-\delta^3 |p_0|^2 K_3 \sum_{\substack{i=-n \\ i \neq i_0}}^n \frac{1}{(x-i)^2} \leq \sum_{\substack{i=-n \\ i \neq i_0}}^n [\phi'(x_i) + \delta \psi'(x_i)] \leq \delta^2 |p_0|^2 (K_1 + \delta K_3) \sum_{\substack{i=-n \\ i \neq i_0}}^n \frac{1}{(x-i)^2},$$

and then by Claim 1 we get

$$(3.8) \quad -C\delta^3 \leq \lim_{n \rightarrow +\infty} \sum_{\substack{i=-n \\ i \neq i_0}}^n [\phi'(x_i) + \delta \psi'(x_i)] \leq C\delta^2.$$

Here and henceforth, C denotes various positive constants independent of x .

Step 3: Bound on $(W''(\phi(x_{i_0})) - W''(0)) \sum_{\substack{i=-n \\ i \neq i_0}}^n \tilde{\phi}(x_i)$

Now, let us prove that

$$(3.9) \quad -C\delta^2 \leq \lim_{n \rightarrow +\infty} (W''(\phi(x_{i_0})) - W''(0)) \sum_{\substack{i=-n \\ i \neq i_0}}^n \tilde{\phi}(x_i) \leq C\delta^2.$$

By (3.1) we have

$$(3.10) \quad \left| \sum_{\substack{i=-n \\ i \neq i_0}}^n \tilde{\phi}(x_i) + \frac{\delta |p_0|}{\alpha \pi} \sum_{\substack{i=-n \\ i \neq i_0}}^n \frac{1}{x-i} \right| \leq K_1 \delta^2 |p_0|^2 \sum_{\substack{i=-n \\ i \neq i_0}}^n \frac{1}{(x-i)^2}.$$

If $|\gamma| \geq \delta|p_0|$, then again from (3.1), $|\tilde{\phi}(x_{i_0}) + \frac{\delta|p_0|}{\alpha\pi\gamma}| \leq K_1 \frac{\delta^2|p_0|^2}{\gamma^2}$ which implies that

$$|W'''(\tilde{\phi}(x_{i_0})) - W'''(0)| \leq |W'''(0)\tilde{\phi}(x_{i_0})| + O(\tilde{\phi}(x_{i_0}))^2 \leq C \frac{\delta}{|\gamma|} + C \frac{\delta^2}{\gamma^2}.$$

By the previous inequality, (3.10) and Claim 1 we deduce that

$$\left| \lim_{n \rightarrow +\infty} (W''(\phi(x_{i_0})) - W''(0)) \sum_{\substack{i=-n \\ i \neq i_0}}^n \tilde{\phi}(x_i) \right| \leq C \left(\frac{\delta}{|\gamma|} + \frac{\delta^2}{\gamma^2} \right) (\delta|\gamma| + \delta^2) \leq C\delta^2,$$

where C is independent of γ .

Finally, if $|\gamma| < \delta|p_0|$, from (3.10) and Claim 1 we conclude that

$$\left| \lim_{n \rightarrow +\infty} (W''(\phi(x_{i_0})) - W''(0)) \sum_{\substack{i=-n \\ i \neq i_0}}^n \tilde{\phi}(x_i) \right| \leq C\delta|\gamma| + C\delta^2 \leq C\delta^2,$$

and (3.9) is proved.

Step 4: Bound on $\delta \sum_{\substack{i=-n \\ i \neq i_0}}^n \mathcal{I}_1[\psi, x_i]$

We have

(3.11)

$$\mathcal{I}_1[\psi] = W'''(\tilde{\phi})\psi + \frac{L}{\alpha}(W'''(\tilde{\phi}) - W'''(0)) + c\phi' = W'''(0)\psi + \frac{L}{\alpha}W'''(0)\tilde{\phi} + O(\tilde{\phi})\psi + O(\tilde{\phi})^2 + c\phi'.$$

Then by (3.11), (3.1), (3.2), (3.5) and Claim 1, we have

$$(3.12) \quad \left| \lim_{n \rightarrow +\infty} \delta \sum_{\substack{i=-n \\ i \neq i_0}}^n \mathcal{I}_1[\psi, x_i] \right| \leq C\delta^2.$$

Step 5: Bound on $W''(\phi(x_{i_0}))\delta \sum_{\substack{i=-n \\ i \neq i_0}}^n \psi(x_i)$

Similarly

$$(3.13) \quad \left| \lim_{n \rightarrow +\infty} W''(\phi(x_{i_0}))\delta \sum_{\substack{i=-n \\ i \neq i_0}}^n \psi(x_i) \right| \leq C\delta^2.$$

Step 6: Bound on the remaining part E

Finally, still from (3.1), (3.5), and Claim 1 it follows that

$$(3.14) \quad \left| \lim_{n \rightarrow +\infty} \sum_{\substack{i=-n \\ i \neq i_0}}^n (\tilde{\phi}(x_i))^2 + O \left(\frac{L\delta}{\alpha} + \delta\psi(x_{i_0}) + \sum_{\substack{i=-n \\ i \neq i_0}}^n [\tilde{\phi}(x_i) + \delta\psi(x_i)] \right) \right|^2 \leq C\delta^2.$$

Step 7: Conclusion

Therefore, from (3.8), (3.9), (3.12), (3.13) and (3.14) we conclude that

$$-C\delta^2 \leq \lim_{n \rightarrow +\infty} NL_L^{\bar{\lambda}_\delta^L} [s_{\delta,n}^L] \leq C\delta^2$$

with C independent of x and Lemma 3.3 is proved.

4. PROOF OF PROPOSITION 2.1

In order to perform the proof of Proposition 2.1, we will use the following technical result whose proof is postponed to the appendix.

Lemma 4.1. (Vanishing far away contribution)

We have

$$(4.1) \quad \lim_{a \rightarrow +\infty} \lim_{n \rightarrow +\infty} \int_{|y| \geq a} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy) = 0.$$

We also need to introduce the notation

$$\mathcal{I}_1^1[f, x] = \int_{|y| < 1} [f(x+y) - f(x) - f'(x)y] \mu(dy)$$

and

$$\mathcal{I}_1^2[f, x] = \int_{|y| \geq 1} [f(x+y) - f(x)] \mu(dy).$$

Proof of Proposition 2.1

Step 1: proof of ii)

Let $x = i_0 + \gamma$ with $i_0 \in \mathbb{Z}$ and $\gamma \in (-\frac{1}{2}, \frac{1}{2}]$. Let $\frac{1}{\delta|p_0|} \geq 2$ and $n > |i_0|$, then by (3.1) and (3.5) we get

$$\begin{aligned} s_{\delta,n}^L(x) - x &= \frac{L\delta}{\alpha} + \phi(x_{i_0}) + \delta\psi(x_{i_0}) - n - i_0 - \gamma + \sum_{\substack{i=-n \\ i \neq i_0}}^n [\phi(x_i) + \delta\psi(x_i)] \\ &= \frac{L\delta}{\alpha} + \phi(x_{i_0}) + \delta\psi(x_{i_0}) - \gamma + \sum_{\substack{i=-n \\ i \neq i_0}}^n [\tilde{\phi}(x_i) + \delta\psi(x_i)] \\ &\leq \frac{L\delta}{\alpha} + \frac{3}{2} + \delta\|\psi\|_\infty + \sum_{\substack{i=-n \\ i \neq i_0}}^n \left[-\left(\frac{1}{\alpha\pi} - \delta K_2\right) \frac{\delta|p_0|}{x-i} + (K_1 + \delta K_3) \frac{\delta^2|p_0|^2}{(x-i)^2} \right]. \end{aligned}$$

Then, by Claim 1

$$h_\delta^L(x) - x = \lim_{n \rightarrow +\infty} s_{\delta,n}^L(x) - x \leq C.$$

Similarly we can prove that

$$h_\delta^L(x) - x \geq -C,$$

which concludes the proof of ii).

Step 2: proof of i)

The function $h_\delta^L(x) = \lim_{n \rightarrow +\infty} s_{\delta,n}^L(x)$ is well defined for any $x \in \mathbb{R}$ by Claim 2. Moreover, by Claim 3 and 4 and classical analysis results, it is of class C^2 on \mathbb{R} with

$$(h_\delta^L)'(x) = \lim_{n \rightarrow +\infty} (s_{\delta,n}^L)'(x) = \lim_{n \rightarrow +\infty} \frac{1}{\delta|p_0|} \sum_{i=-n}^n \left[\phi' \left(\frac{x-i}{\delta|p_0|} \right) + \delta\psi' \left(\frac{x-i}{\delta|p_0|} \right) \right],$$

$$(h_\delta^L)''(x) = \lim_{n \rightarrow +\infty} (s_{\delta,n}^L)''(x) = \lim_{n \rightarrow +\infty} \frac{1}{\delta^2 |p_0|^2} \sum_{i=-n}^n \left[\phi'' \left(\frac{x-i}{\delta |p_0|} \right) + \delta \psi'' \left(\frac{x-i}{\delta |p_0|} \right) \right],$$

and the convergence of $\{s_{\delta,n}^L\}_n$, $\{(s_{\delta,n}^L)'\}_n$ and $\{(s_{\delta,n}^L)''\}_n$ is uniform on compact sets.

Let us show that for any $x \in \mathbb{R}$

$$(4.2) \quad \mathcal{I}_1[h_\delta^L, x] = \lim_{n \rightarrow +\infty} \mathcal{I}_1[s_{\delta,n}^L, x].$$

Step 2.1: term $\mathcal{I}_1^1[h_\delta^L, x]$

First, we prove that

$$(4.3) \quad \mathcal{I}_1^1[h_\delta^L, x] = \lim_{n \rightarrow +\infty} \mathcal{I}_1^1[s_{\delta,n}^L, x].$$

Fix $x \in \mathbb{R}$, we know that for any $y \in [-1, 1]$, $y \neq 0$

$$\frac{s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x) - (s_{\delta,n}^L)'(x)y}{|y|^2} \rightarrow \frac{h_\delta^L(x+y) - h_\delta^L(x) - (h_\delta^L)'(x)y}{|y|^2} \quad \text{as } n \rightarrow +\infty.$$

By the uniform convergence of the sequence $\{(s_{\delta,n}^L)''\}_n$ we have

$$\frac{|s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x) - (s_{\delta,n}^L)'(x)y|}{|y|^2} \leq \sup_{z \in [x-1, x+1]} (s_{\delta,n}^L)''(z) \leq C,$$

where C is independent of n , and (4.3) follows from the dominate convergence Theorem.

Step 2.2: term $\mathcal{I}_1^2[h_\delta^L, x]$

Then, to prove (4.2) it suffices to show that

$$\mathcal{I}_1^2[h_\delta^L, x] = \lim_{n \rightarrow +\infty} \mathcal{I}_1^2[s_{\delta,n}^L, x].$$

From Claim 5 and (4.3), we know that for any $x \in \mathbb{R}$ there exists $\lim_{n \rightarrow +\infty} \mathcal{I}_1^2[s_{\delta,n}^L, x]$. For $a > 1$, we have

$$\mathcal{I}_1^2[s_{\delta,n}^L, x] = \int_{1 \leq |y| \leq a} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy) + \int_{|y| \geq a} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy).$$

By the uniform convergence of $\{s_{\delta,n}^L\}_n$ on compact sets

$$\lim_{n \rightarrow +\infty} \int_{1 \leq |y| \leq a} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy) = \int_{1 \leq |y| \leq a} [h_\delta^L(x+y) - h_\delta^L(x)] \mu(dy),$$

then there exists the limit

$$\lim_{n \rightarrow +\infty} \int_{|y| \geq a} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy).$$

Then, we finally get

$$\begin{aligned}
\lim_{n \rightarrow +\infty} \mathcal{I}_1^2[s_{\delta,n}^L, x] &= \lim_{a \rightarrow +\infty} \lim_{n \rightarrow +\infty} \mathcal{I}_1^2[s_{\delta,n}^L, x] \\
&= \lim_{a \rightarrow +\infty} \lim_{n \rightarrow +\infty} \int_{1 \leq |y| \leq a} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy) \\
&+ \lim_{a \rightarrow +\infty} \lim_{n \rightarrow +\infty} \int_{|y| > a} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy) \\
&= \lim_{a \rightarrow +\infty} \int_{1 \leq |y| \leq a} [h_\delta^L(x+y) - h_\delta^L(x)] \mu(dy) \\
&= \mathcal{I}_1^2[h_\delta^L, x],
\end{aligned}$$

as desired, where we have used Lemma 4.1.

Step 2.3: conclusion

Now we can conclude the proof of (i). Indeed, by Claim 2, Claim 3 and (4.2), for any $x \in \mathbb{R}$

$$NL_L^{\bar{\lambda}_\delta^L}[h_\delta^L](x) = \lim_{n \rightarrow +\infty} NL_L^{\bar{\lambda}_\delta^L}[s_{\delta,n}^L](x),$$

and Lemma 3.3 implies that

$$NL_L^{\bar{\lambda}_\delta^L}[h_\delta^L](x) = o(\delta), \quad \text{as } \delta \rightarrow 0,$$

where $\lim_{\delta \rightarrow 0} \frac{o(\delta)}{\delta} = 0$, uniformly for $x \in \mathbb{R}$.

5. APPENDIX

In this appendix, we prove the following technical results used in the previous section: Lemmata 3.1 and 3.2, the Claims 1-5 and Lemma 4.1.

5.1. Proof of Lemma 3.1. Properties (3.1) and (3.2) are proved in [7].

Let us show (3.3).

For $a > 0$, we denote by $\phi'_a(x) = \phi'\left(\frac{x}{a}\right)$. Remark that ϕ'_a is a solution of

$$\mathcal{I}_1[\phi'_a] = \frac{1}{a} W''(\phi_a) \phi'_a \quad \text{in } \mathbb{R}.$$

Since ϕ'' is bounded and of class $C^{2,\beta}$, $\mathcal{I}_1[\phi'']$ is well defined and by deriving twice the equation in (1.6) we see that ϕ'' is a solution of

$$\mathcal{I}_1[\phi''] = W''(\phi) \phi'' + W'''(\phi) (\phi')^2.$$

Let $\bar{\phi} = \phi'' - C \phi'_a$, with $C > 0$, then $\bar{\phi}$ satisfies

$$\begin{aligned}
\mathcal{I}_1[\bar{\phi}] - W''(\phi) \bar{\phi} &= C \phi'_a \left(W''(\phi) - \frac{1}{a} W''(\phi_a) \right) + W'''(\phi) (\phi')^2 \\
&= C \phi'_a \left(W''(\phi) - \frac{1}{a} W''(\phi_a) \right) + o\left(\frac{1}{1+x^2}\right),
\end{aligned}$$

as $|x| \rightarrow +\infty$, by (3.2). Fix $a > 0$ and $R > 0$ such that

$$(5.1) \quad \begin{cases} W''(\phi) - \frac{1}{a} W''(\phi_a) > \frac{1}{2} W''(0) > 0 & \text{on } \mathbb{R} \setminus [-R, R]; \\ W''(\phi) > 0, & \text{on } \mathbb{R} \setminus [-R, R]. \end{cases}$$

Then from (3.2), for C large enough we get

$$\mathcal{I}_1[\bar{\phi}] - W''(\phi)\bar{\phi} \geq 0 \quad \text{on } \mathbb{R} \setminus [-R, R].$$

Choosing C such that moreover

$$\bar{\phi} < 0 \quad \text{on } [-R, R],$$

we can ensure that $\bar{\phi} \leq 0$ on \mathbb{R} . Indeed, assume by contradiction that there exists $x_0 \in \mathbb{R} \setminus [-R, R]$ such that

$$\bar{\phi}(x_0) = \sup_{\mathbb{R}} \bar{\phi} > 0.$$

Then

$$\begin{cases} \mathcal{I}_1[\bar{\phi}, x_0] \leq 0; \\ \mathcal{I}_1[\bar{\phi}, x_0] - W''(\phi(x_0))\bar{\phi}(x_0) \geq 0; \\ W''(\phi(x_0)) > 0, \end{cases}$$

from which

$$\bar{\phi}(x_0) \leq 0,$$

a contradiction. Therefore $\bar{\phi} \leq 0$ on \mathbb{R} and then, by renaming the constants, from (3.2) we get $\phi'' \leq \frac{K_1}{1+x^2}$.

To prove that $\phi'' \geq -\frac{K_1}{1+x^2}$, we look at the infimum of the function $\phi'' + C\phi'_a$ to get similarly that $\phi'' + C\phi'_a \geq 0$ on \mathbb{R} .

To show (3.4) we proceed as in the proof of (3.3). Indeed, the function ϕ''' which is bounded and of class $C^{1,\beta}$, satisfies

$$\mathcal{I}_1[\phi'''] = W''(\phi)\phi''' + 3W'''(\phi)\phi'\phi'' + W^{IV}(\phi)(\phi')^3 = W''(\phi)\phi''' + o\left(\frac{1}{1+x^2}\right),$$

as $|x| \rightarrow +\infty$, by (3.2) and (3.3). Then, as before, for C and a large enough $\phi''' - C\phi'_a \leq 0$ and $\phi''' + C\phi'_a \geq 0$ on \mathbb{R} , which implies (3.4). \square

5.2. Proof of Lemma 3.2. Let us prove (3.5).

For $a > 0$ we denote by $\phi_a(x) = \phi\left(\frac{x}{a}\right)$, which is solution of

$$\mathcal{I}_1[\phi_a] = \frac{1}{a}W'(\phi_a) \quad \text{in } \mathbb{R}.$$

Let a and b be positive numbers, then making a Taylor expansion of the derivatives of W , we get

$$\begin{aligned} \mathcal{I}_1[\psi - (\phi_a - \phi_b)] &= W''(\phi)\psi + \frac{L}{\alpha}(W''(\phi) - W''(0)) + c\phi' + \left(\frac{1}{b}W'(\phi_b) - \frac{1}{a}W'(\phi_a)\right) \\ &= W''(\phi)(\psi - (\phi_a - \phi_b)) + W''(\tilde{\phi})(\phi_a - \phi_b) + \frac{L}{\alpha}(W''(\tilde{\phi}) - W''(0)) \\ &\quad + c\phi' + \left(\frac{1}{b}W'(\tilde{\phi}_b) - \frac{1}{a}W'(\tilde{\phi}_a)\right) \\ &= W''(\phi)(\psi - (\phi_a - \phi_b)) + W''(0)(\phi_a - \phi_b) + \frac{L}{\alpha}W'''(0)\tilde{\phi} + c\phi' \\ &\quad + W''(0)\left(\frac{1}{b}\tilde{\phi}_b - \frac{1}{a}\tilde{\phi}_a\right) + (\phi_a - \phi_b)O(\tilde{\phi}) + O(\tilde{\phi})^2 + O(\tilde{\phi}_a)^2 + O(\tilde{\phi}_b)^2, \end{aligned}$$

and then the function $\bar{\psi} = \psi - (\phi_a - \phi_b)$ satisfies

$$\begin{aligned} \mathcal{I}_1[\bar{\psi}] - W''(\phi)\bar{\psi} &= \alpha(\phi_a - \phi_b) + \frac{L}{\alpha}W'''(0)\tilde{\phi} + c\phi' + \alpha\left(\frac{1}{b}\tilde{\phi}_b - \frac{1}{a}\tilde{\phi}_a\right) \\ &\quad + (\phi_a - \phi_b)O(\tilde{\phi}) + O(\tilde{\phi})^2 + O(\tilde{\phi}_a)^2 + O(\tilde{\phi}_b)^2. \end{aligned}$$

We want to estimate the right-hand side of the last equality. By Lemma 3.1, for $|x| \geq \max\{1, |a|, |b|\}$ we have

$$\alpha(\phi_a - \phi_b) + \frac{L}{\alpha}W'''(0)\tilde{\phi} \geq -\frac{1}{\pi x} \left[(a-b) + \frac{L}{\alpha^2}W'''(0) \right] - \frac{K_1\alpha}{x^2} \left(a^2 + b^2 + \frac{|L|}{\alpha^2}|W'''(0)| \right).$$

Choose $a, b > 0$ such that $(a-b) + \frac{L}{\alpha^2}W'''(0) = 0$, then

$$\alpha(\phi_a - \phi_b) + \frac{L}{\alpha}W'''(0)\tilde{\phi} \geq -\frac{C}{x^2},$$

for $|x| \geq \max\{1, |a|, |b|\}$. Here and in what follows, as usual C denotes various positive constants. From Lemma 3.1 we also derive that

$$\begin{aligned} \alpha\left(\frac{1}{b}\tilde{\phi}_b - \frac{1}{a}\tilde{\phi}_a\right) &\geq -\frac{C}{x^2}, \\ c\phi' &\geq -\frac{C}{1+x^2}, \end{aligned}$$

and

$$(\phi_a - \phi_b)O(\tilde{\phi}) + O(\tilde{\phi})^2 + O(\tilde{\phi}_a)^2 + O(\tilde{\phi}_b)^2 \geq -\frac{C}{1+x^2},$$

for $|x| \geq \max\{1, |a|, |b|\}$. Then we conclude that there exists $R > 0$ such that for $|x| \geq R$ we have

$$\mathcal{I}_1[\bar{\psi}] - W''(\phi)\bar{\psi} \geq -\frac{C}{1+x^2}.$$

Now, let us consider the function $\phi'_d(x) = \phi'\left(\frac{x}{d}\right)$, $d > 0$, which is solution of

$$\mathcal{I}_1[\phi'_d] = \frac{1}{d}W''(\phi_d)\phi'_d \quad \text{in } \mathbb{R},$$

and denote

$$\bar{\bar{\psi}} = \bar{\psi} - \tilde{C}\phi'_d,$$

with $\tilde{C} > 0$. Then, for $|x| \geq R$ we have

$$\mathcal{I}_1[\bar{\bar{\psi}}] \geq W''(\phi)\bar{\psi} - \frac{\tilde{C}}{d}W''(\phi_d)\phi'_d - \frac{C}{1+x^2} = W''(\phi)\bar{\bar{\psi}} + \tilde{C}\phi'_d \left(W''(\phi) - \frac{1}{d}W''(\phi_d) \right) - \frac{C}{1+x^2}.$$

Let us choose $d > 0$ and $R_2 > R$ such that

$$\begin{cases} W''(\phi) - \frac{1}{d}W''(\phi_d) > \frac{1}{2}W''(0) > 0 & \text{on } \mathbb{R} \setminus [-R_2, R_2]; \\ W''(\phi) > 0 & \text{on } \mathbb{R} \setminus [-R_2, R_2], \end{cases}$$

then from (3.2), for \tilde{C} large enough we get

$$\mathcal{I}_1[\bar{\bar{\psi}}] - W''(\phi)\bar{\bar{\psi}} \geq 0 \quad \text{on } \mathbb{R} \setminus [-R_2, R_2],$$

and

$$\bar{\bar{\psi}} < 0 \quad \text{on } [-R_2, R_2].$$

As in the proof of Lemma 3.1, we deduce that $\overline{\psi} \leq 0$ on \mathbb{R} and then

$$\psi \leq \frac{K_2}{x} + \frac{K_3}{x^2} \quad \text{for } |x| \geq 1,$$

for some $K_2 \in \mathbb{R}$ and $K_3 > 0$.

Looking at the function $\psi - (\phi_a - \phi_b) + \tilde{C}\phi'_a$, we conclude similarly that

$$\psi \geq \frac{K_2}{x} - \frac{K_3}{x^2} \quad \text{for } |x| \geq 1,$$

and (3.5) is proved.

Now let us turn to (3.6). By deriving the first equation in (2.5), we see that the function ψ' which is bounded and of class $C^{2,\beta}$, is a solution of

$$\mathcal{I}_1[\psi'] = W''(\phi)\psi' + W'''(\phi)\phi'\psi + \frac{L}{\alpha}W'''(\phi)\phi' + c\phi'' \quad \text{in } \mathbb{R}.$$

Then the function $\overline{\psi}' = \psi' - C\phi'_a$, satisfies

$$\begin{aligned} \mathcal{I}_1[\overline{\psi}'] - W''(\phi)\overline{\psi}' &= C\phi'_a \left(W''(\phi) - \frac{1}{a}W''(\phi_a) \right) + W'''(\phi)\phi'\psi + \frac{L}{\alpha}W'''(\phi)\phi' + c\phi'' \\ &= C\phi'_a \left(W''(\phi) - \frac{1}{a}W''(\phi_a) \right) + O\left(\frac{1}{1+x^2}\right), \end{aligned}$$

by (3.2), (3.3) and (3.5), and as in the proof of Lemma 3.1, we deduce that for C and a large enough $\overline{\psi}' \leq 0$ on \mathbb{R} , which implies that $\psi' \leq \frac{K_3}{1+x^2}$. The inequality $\psi' \geq -\frac{K_3}{1+x^2}$ is obtained similarly by proving that $\overline{\psi}' + C\phi'_a \geq 0$ on \mathbb{R} .

Finally, with the same proof as before, using (3.2)-(3.6), we can prove the estimate (3.7) for the function ψ'' which is a bounded $C^{1,\beta}$ solution of

$$\begin{aligned} \mathcal{I}_1[\psi''] &= W''(\phi)\psi'' + 2W'''(\phi)\phi'\psi' + W^{IV}(\phi)(\phi')^2\psi + W'''(\phi)\phi''\psi + \frac{L}{\alpha}W'''(\phi)\phi'' \\ &\quad + \frac{L}{\alpha}W^{IV}(\phi)(\phi')^2 + c\phi''' \\ &= W''(\phi)\psi'' + O\left(\frac{1}{1+x^2}\right). \end{aligned}$$

□

5.3. Proof of Claims 1-5.

Proof of Claim 1.

We have for $n > |i_0|$

$$\begin{aligned} \sum_{\substack{i=-n \\ i \neq i_0}}^n \frac{1}{x-i} &= \sum_{i=-n}^{i_0-1} \frac{1}{i_0 + \gamma - i} + \sum_{i=i_0+1}^n \frac{1}{i_0 + \gamma - i} = \sum_{i=1}^{n+i_0} \frac{1}{i + \gamma} - \sum_{i=1}^{n-i_0} \frac{1}{i - \gamma} \\ &= \begin{cases} \sum_{i=1}^n \frac{-2\gamma}{i^2 - \gamma^2}, & \text{if } i_0 = 0 \\ \sum_{i=1}^{n-i_0} \frac{-2\gamma}{i^2 - \gamma^2} + \sum_{i=n-i_0+1}^{n+i_0} \frac{1}{i+\gamma}, & \text{if } i_0 > 0 \\ \sum_{i=1}^{n+i_0} \frac{-2\gamma}{i^2 - \gamma^2} - \sum_{i=n+i_0+1}^{n-i_0} \frac{1}{i-\gamma}, & \text{if } i_0 < 0 \end{cases} \rightarrow -2\gamma \sum_{i=1}^{+\infty} \frac{1}{i^2 - \gamma^2} \quad \text{as } n \rightarrow +\infty. \end{aligned}$$

Let us prove the second limit of the claim.

$$\sum_{i=-n}^{i_0-1} \frac{1}{(x-i)^2} = \sum_{i=1}^{n+i_0} \frac{1}{(i+\gamma)^2} \rightarrow \sum_{i=1}^{+\infty} \frac{1}{(i+\gamma)^2} \quad \text{as } n \rightarrow +\infty.$$

Finally

$$\sum_{i=i_0+1}^n \frac{1}{(x-i)^2} = \sum_{i=1}^{n-i_0} \frac{1}{(i-\gamma)^2} \rightarrow \sum_{i=1}^{+\infty} \frac{1}{(i-\gamma)^2} \quad \text{as } n \rightarrow +\infty,$$

and the claim is proved.

By Claim 1 $\sum_{\substack{i=-n \\ i \neq i_0}}^n \frac{1}{x-i}$, $\sum_{i=-n}^{i_0-1} \frac{1}{(x-i)^2}$ and $\sum_{i=i_0+1}^n \frac{1}{(x-i)^2}$ are Cauchy sequences and then for $k > m > |i_0|$ we have

$$(5.2) \quad \sum_{i=-k}^{-m-1} \frac{1}{x-i} + \sum_{i=m+1}^k \frac{1}{x-i} \rightarrow 0 \quad \text{as } m, k \rightarrow +\infty,$$

$$(5.3) \quad \sum_{i=-k}^{-m-1} \frac{1}{(x-i)^2} \rightarrow 0 \quad \text{as } m, k \rightarrow +\infty,$$

and

$$(5.4) \quad \sum_{i=m+1}^k \frac{1}{(x-i)^2} \rightarrow 0 \quad \text{as } m, k \rightarrow +\infty.$$

Proof of Claim 2.

We show that $\{s_{\delta,n}^L(x)\}_n$ is a Cauchy sequence. Fix $x \in \mathbb{R}$ and let $i_0 \in \mathbb{Z}$ be the closest integer to x such that $x = i_0 + \gamma$, with $\gamma \in (-\frac{1}{2}, \frac{1}{2}]$ and $|x-i| \geq \frac{1}{2}$ for $i \neq i_0$. Let δ be so small that $\frac{1}{\delta|p_0|} \geq 2$, then $\frac{|x-i|}{\delta|p_0|} \geq 1$ for $i \neq i_0$. Let $k > m > |i_0|$, using (3.1) and (3.5) we get

$$\begin{aligned} s_{\delta,k}^L(x) - s_{\delta,m}^L(x) &= -(k-m) + \sum_{i=-k}^{-m-1} [\phi(x_i) + \delta\psi(x_i)] + \sum_{i=m+1}^k [\phi(x_i) + \delta\psi(x_i)] \\ &= \sum_{i=-k}^{-m-1} [(\phi(x_i) - 1) + \delta\psi(x_i)] + \sum_{i=m+1}^k [\phi(x_i) + \delta\psi(x_i)] \\ &\leq -\left(\frac{1}{\alpha\pi} - \delta K_2\right) \delta|p_0| \sum_{i=-k}^{-m-1} \frac{1}{x-i} + (K_1 + \delta K_3) \delta^2 |p_0|^2 \sum_{i=-k}^{-m-1} \frac{1}{(x-i)^2} \\ &\quad - \left(\frac{1}{\alpha\pi} - \delta K_2\right) \delta|p_0| \sum_{i=m+1}^k \frac{1}{x-i} + (K_1 + \delta K_3) \delta^2 |p_0|^2 \sum_{i=m+1}^k \frac{1}{(x-i)^2}, \end{aligned}$$

and

$$\begin{aligned} s_{\delta,k}^L(x) - s_{\delta,m}^L(x) &\geq -\left(\frac{1}{\alpha\pi} - \delta K_2\right) \delta|p_0| \left(\sum_{i=-k}^{-m-1} \frac{1}{x-i} + \sum_{i=m+1}^k \frac{1}{x-i} \right) \\ &\quad - (K_1 + \delta K_3) \delta^2 |p_0|^2 \left(\sum_{i=-k}^{-m-1} \frac{1}{(x-i)^2} + \sum_{i=m+1}^k \frac{1}{(x-i)^2} \right). \end{aligned}$$

Then from (5.2), (5.3), (5.4), we conclude that

$$|s_{\delta,k}^L(x) - s_{\delta,m}^L(x)| \rightarrow 0 \quad \text{as } m, k \rightarrow +\infty,$$

as desired.

Proof of Claim 3.

To prove the uniform convergence, it suffices to show that $\{(s_{\delta,n}^L)'(x)\}_n$ is a Cauchy sequence uniformly on compact sets. Let us consider a bounded interval $[a, b]$ and let $x \in [a, b]$. For $\frac{1}{\delta|p_0|} \geq 2$ and $k > m > 1/2 + \max\{|a|, |b|\}$, by (3.2) and (3.6) we have

$$\begin{aligned} (s_{\delta,k}^L)'(x) - (s_{\delta,m}^L)'(x) &= \frac{1}{\delta|p_0|} \sum_{i=-k}^{-m-1} [\phi'(x_i) + \delta\psi'(x_i)] + \frac{1}{\delta|p_0|} \sum_{i=m+1}^k [\phi'(x_i) + \delta\psi'(x_i)] \\ &\leq (K_1 + \delta K_3)\delta|p_0| \left[\sum_{i=-k}^{-m-1} \frac{1}{(x-i)^2} + \sum_{i=m+1}^k \frac{1}{(x-i)^2} \right] \\ &\leq (K_1 + \delta K_3)\delta|p_0| \left[\sum_{i=-k}^{-m-1} \frac{1}{(a-i)^2} + \sum_{i=m+1}^k \frac{1}{(b-i)^2} \right], \end{aligned}$$

and

$$(s_{\delta,k}^L)'(x) - (s_{\delta,m}^L)'(x) \geq -K_3\delta^2|p_0| \left[\sum_{i=-k}^{-m-1} \frac{1}{(a-i)^2} + \sum_{i=m+1}^k \frac{1}{(b-i)^2} \right].$$

Then by (5.3) and (5.4)

$$\sup_{x \in [a, b]} |(s_{\delta,k}^L)'(x) - (s_{\delta,m}^L)'(x)| \rightarrow 0 \quad \text{as } k, m \rightarrow +\infty,$$

and Claim 3 is proved.

Proof of Claim 4.

Claim 4 can be proved like Claim 3. Indeed

$$(s_{\delta,n}^L)''(x) = \frac{1}{\delta^2|p_0|^2} \sum_{i=-n}^n [\phi''(x_i) + \delta\psi''(x_i)]$$

and using (3.3) and (3.7), it is easy to show that $\{(s_{\delta,n}^L)''\}_n$ is a Cauchy sequence uniformly on compact sets.

Proof of Claim 5.

We have

$$\mathcal{I}_1[\phi] = W'(\phi) = W'(\tilde{\phi}) = W''(0)\tilde{\phi} + O(\tilde{\phi})^2.$$

Let $x = i_0 + \gamma$ with $\gamma \in (-\frac{1}{2}, \frac{1}{2}]$, and $k > m > |i_0|$. From (3.1), (5.2), (5.3) and (5.4) we get

$$\begin{aligned} \sum_{i=-k}^k \mathcal{I}_1[\phi, x_i] - \sum_{i=-m}^m \mathcal{I}_1[\phi, x_i] &= \sum_{i=-k}^{-m-1} [\alpha\tilde{\phi}(x_i) + O(\tilde{\phi}(x_i))^2] + \sum_{i=m+1}^k [\alpha\tilde{\phi}(x_i) + O(\tilde{\phi}(x_i))^2] \\ &\leq -\frac{\delta|p_0|}{\pi} \left[\sum_{i=-k}^{-m-1} \frac{1}{x-i} + \sum_{i=m+1}^k \frac{1}{x-i} \right] + C \sum_{i=-k}^{-m-1} \frac{1}{(x-i)^2} + C \sum_{i=m+1}^k \frac{1}{(x-i)^2} \rightarrow 0, \end{aligned}$$

as $m, k \rightarrow +\infty$, for some constant $C > 0$, and

$$\begin{aligned} & \sum_{i=-k}^k \mathcal{I}_1[\phi, x_i] - \sum_{i=-m}^m \mathcal{I}_1[\phi, x_i] \\ & \geq -\frac{\delta|p_0|}{\pi} \left[\sum_{i=-k}^{-m-1} \frac{1}{x-i} + \sum_{i=m+1}^k \frac{1}{x-i} \right] - C \sum_{i=-k}^{-m-1} \frac{1}{(x-i)^2} - C \sum_{i=m+1}^k \frac{1}{(x-i)^2} \rightarrow 0, \end{aligned}$$

as $m, k \rightarrow +\infty$. Then $\sum_{i=-n}^n \mathcal{I}_1[\phi, x_i]$ is a Cauchy sequence, i.e. it converges.

Let us consider now $\sum_{i=-n}^n \mathcal{I}_1[\psi, x_i]$. By (3.11), (3.1), (3.2) and (3.5) we get

$$\begin{aligned} & \sum_{i=-k}^k \mathcal{I}_1[\psi, x_i] - \sum_{i=-m}^m \mathcal{I}_1[\psi, x_i] \\ & \leq \tilde{C} \left[\sum_{i=-k}^{-m-1} \frac{1}{x-i} + \sum_{i=m+1}^k \frac{1}{x-i} \right] + C \sum_{i=-k}^{-m-1} \frac{1}{(x-i)^2} + C \sum_{i=m+1}^k \frac{1}{(x-i)^2}, \end{aligned}$$

and

$$\begin{aligned} & \sum_{i=-k}^k \mathcal{I}_1[\psi, x_i] - \sum_{i=-m}^m \mathcal{I}_1[\psi, x_i] \\ & \geq \tilde{C} \left[\sum_{i=-k}^{-m-1} \frac{1}{x-i} + \sum_{i=m+1}^k \frac{1}{x-i} \right] - C \sum_{i=-k}^{-m-1} \frac{1}{(x-i)^2} - C \sum_{i=m+1}^k \frac{1}{(x-i)^2}, \end{aligned}$$

for some $\tilde{C} \in \mathbb{R}$ and $C > 0$, which ensures the convergence of $\sum_{i=-n}^n \mathcal{I}_1[\psi, x_i]$.

Proof of Lemma 4.1

Let $\frac{1}{\delta|p_0|} \geq 2$. We first remark that if $z > n + \frac{1}{2}$, then $z_i = \frac{z-i}{\delta|p_0|} \geq 1$ for $i = -n, \dots, n$ and by (3.1) and (3.5) we have

$$\begin{aligned} s_{\delta,n}^L(z) &= \frac{L\delta}{\alpha} + n + 1 + \sum_{i=-n}^n [\phi(z_i) - 1 + \delta\psi(z_i)] \\ &\leq \frac{L\delta}{\alpha} + n + 1 + \sum_{i=-n}^n \left[-\left(\frac{1}{\alpha\pi} - \delta K_2\right) \frac{\delta|p_0|}{z-i} + (K_1 + \delta K_3) \frac{\delta^2|p_0|^2}{(z-i)^2} \right], \end{aligned}$$

and

$$s_{\delta,n}^L(z) \geq \frac{L\delta}{\alpha} + n + 1 + \sum_{i=-n}^n \left[-\left(\frac{1}{\alpha\pi} - \delta K_2\right) \frac{\delta|p_0|}{z-i} - (K_1 + \delta K_3) \frac{\delta^2|p_0|^2}{(z-i)^2} \right].$$

By Claim 1, the quantities $\sum_{i=-n}^n \frac{1}{z-i}$ and $\sum_{i=-n}^n \frac{1}{(z-i)^2}$ are uniformly bounded on \mathbb{R} by a constant independent of n . Hence, we get

$$(5.5) \quad n - C \leq s_{\delta,n}^L(z) \leq n + C \quad \text{if } z > n + \frac{1}{2}.$$

The same argument shows that

$$(5.6) \quad -n - C \leq s_{\delta,n}^L(z) \leq -n + C \quad \text{if } z < -n - \frac{1}{2}.$$

If $|z| < n - \frac{1}{2}$, then $n > |j_0|$, where j_0 is the closest integer to z , and as in the proof of (ii) of Proposition 2.1 (see Step 1 there), we get

$$s_{\delta,n}^L(z) - z \leq \frac{L\delta}{\alpha} + \frac{3}{2} + \delta\|\psi\|_\infty + \sum_{\substack{i=-n \\ i \neq j_0}}^n \left[-\left(\frac{1}{\alpha\pi} - \delta K_2\right) \frac{\delta|p_0|}{z-i} + (K_1 + \delta K_3) \frac{\delta^2|p_0|^2}{(z-i)^2} \right],$$

and

$$s_{\delta,n}^L(z) - z \geq \frac{L\delta}{\alpha} - \frac{1}{2} - \delta\|\psi\|_\infty + \sum_{\substack{i=-n \\ i \neq j_0}}^n \left[-\left(\frac{1}{\alpha\pi} - \delta K_2\right) \frac{\delta|p_0|}{z-i} - (K_1 + \delta K_3) \frac{\delta^2|p_0|^2}{(z-i)^2} \right].$$

Then, again by Claim 1

$$(5.7) \quad -C \leq s_{\delta,n}^L(z) - z \leq C \quad \text{if } |z| < n - \frac{1}{2}.$$

Now, let $i_0 \in \mathbb{Z}$ be the closest integer to x , let us assume $n > |i_0| + 1 + a$. We have

$$\begin{aligned} \int_{|y| \geq a} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy) &= \int_{a \leq |y| < n-1-|i_0|} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy) \\ &\quad + \int_{n-1-|i_0| \leq |y| \leq n+1+|i_0|} [\dots] \mu(dy) + \int_{|y| > n+1+|i_0|} [\dots] \mu(dy). \end{aligned}$$

If $|y| < n - 1 - |i_0|$, then $|x+y| < n - \frac{1}{2}$ and by (5.7)

$$\begin{aligned} \int_{a \leq |y| < n-1-|i_0|} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy) &\leq \int_{a \leq |y| \leq n-1-|i_0|} (y+2C) \mu(dy) \\ &= \int_{a \leq |y| \leq n-1-|i_0|} 2C \mu(dy) \leq \frac{2C}{a}, \end{aligned}$$

and

$$\int_{a \leq |y| < n-1-|i_0|} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy) \geq -\frac{2C}{a}.$$

Then

$$(5.8) \quad \lim_{a \rightarrow +\infty} \lim_{n \rightarrow +\infty} \int_{a \leq |y| \leq n-1-|i_0|} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy) = 0.$$

Next, since $|s_{\delta,n}^L(z)| \leq Cn$ for any $z \in \mathbb{R}$, we have

$$(5.9) \quad \begin{aligned} &\left| \int_{n-1-|i_0| \leq |y| \leq n+1+|i_0|} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)] \mu(dy) \right| \\ &\leq Cn \int_{n-1-|i_0| \leq |y| \leq n+1+|i_0|} \mu(dy) = \tilde{C} \frac{n(|i_0|+1)}{n^2 - (|i_0|+1)^2} \rightarrow 0 \quad \text{as } n \rightarrow +\infty. \end{aligned}$$

Finally, if $y > n + 1 + |i_0|$, then $x + y > n + \frac{1}{2}$, while if $y < -n - 1 - |i_0|$, then $x + y < -n - \frac{1}{2}$. Hence, using (5.5) and (5.6), we obtain

$$\begin{aligned} & \int_{|y|>n+1+|i_0|} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)]\mu(dy) \\ &= \int_{y>n+1+|i_0|} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)]\mu(dy) + \int_{y<-n-1-|i_0|} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)]\mu(dy) \\ &\leq \int_{y>n+1+|i_0|} [n + C - s_{\delta,n}^L(x)]\mu(dy) + \int_{y<-n-1-|i_0|} [-n + C - s_{\delta,n}^L(x)]\mu(dy) \\ &= \int_{|y|>n+1+|i_0|} [C - s_{\delta,n}^L(x)]\mu(dy), \end{aligned}$$

and

$$\int_{|y|>n+1+|i_0|} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)]\mu(dy) \geq \int_{|y|>n+1+|i_0|} [-C - s_{\delta,n}^L(x)]\mu(dy).$$

We deduce that

$$\lim_{n \rightarrow +\infty} \int_{|y|>n+1+|i_0|} [s_{\delta,n}^L(x+y) - s_{\delta,n}^L(x)]\mu(dy) = 0.$$

Hence, by the previous limit, (5.8) and (5.9), we derive (4.1). This ends the proof of Lemma 4.1.

REFERENCES

- [1] P. BILER, G. KARCH, R. MONNEAU Nonlinear diffusion of dislocation density and self-similar solutions, *Communications in Mathematical Physics*, **294** (2010), no. 1, 145-168.
- [2] X. CABRÉ AND J. SOLÀ-MORALES, Layer solutions in a half-space for boundary reactions, *Comm. Pure Appl. Math.*, **58** (2005) no. 12, 1678-1732.
- [3] A. DE MASI, E. ORLANDI, E. PRESUTTI AND L. TRIOLO, Motion by curvature by scaling nonlocal evolution equations, *J. Statist. Phys.*, **73** (1993), 543-570.
- [4] C. DENOUAL, Dynamic dislocation modeling by combining Peierls Nabarro and Galerkin methods, *Phys. Rev. B*, **70** (2004), 024106.
- [5] J. DRONIOU AND C. IMBERT, Fractal first order partial differential equations, *Archive for Rational Mechanics and Analysis*, **182** (2006), no. 2, 299-331.
- [6] N. FORCADEL, C. IMBERT AND R. MONNEAU, Homogenization of fully overdamped Frenkel-Kontorova models, *Journal of Differential Equations*, **246** (2009), no. 1, 1057-1097.
- [7] M. GONZÁLEZ AND R. MONNEAU, Slow motion of particle systems as a limit of a reaction-diffusion equation with half-Laplacian in dimension one, preprint hal-00497492.
- [8] A. K. HEAD Dislocation group dynamics III. Similarity solutions of the continuum approximation, *Phil. Magazine*, **26**, (1972), 65-72.
- [9] J. R. HIRTH AND L. LOTHE, Theory of dislocations, Second Edition. Malabar, Florida: Krieger, 1992.
- [10] P. L. LIONS, G. C. PAPANICOLAOU AND S. R. S. VARADHAN, Homogenization of Hamilton-Jacobi equations, unpublished, 1986.
- [11] R. MONNEAU AND S. PATRIZI, Homogenization of the Peierls-Nabarro model for dislocation dynamics, *preprint*.
- [12] A.B. MOVCHAN, R. BULLOUGH, J.R. WILLIS, Stability of a dislocation: discrete model, *Eur. J. Appl. Math.* **9** (1998), 373-396.
- [13] F.R.N. NABARRO, Fifty-year study of the Peierls-Nabarro stress, *Material Science and Engineering A* **234-236** (1997), 67-76.

- [14] H. WEI, Y. XIANG, P. MING, A Generalized Peierls-Nabarro Model for Curved Dislocations Using Discrete Fourier Transform, *Communications in computational physics* 4(2) (2008), 275-293.

E-mail address: monneau@cermics.enpc.fr

E-mail address: spatrizi@math.ist.utl.pt