ON THE EVANS-KRYLOV THEOREM

LUIS CAFFARELLI AND LUIS SILVESTRE

The Evans-Krylov theorem consists of the a priori estimate:

Theorem 1. Smooth solutions, u, of a uniformly elliptic, fully non-linear convex equation $F(D^2u) = 0$ in the unit ball B_1 , of \mathbb{R}^n have a $C^{2,\alpha}$ interior a priori estimate

 $||u||_{C^{2,\alpha}(B_{1/2})} \le C ||u||_{C^{1,1}(B_1)}$

with the constant C depending only on the ellipticity of F.

The importance of the Evans-Krylov theorem is that it allows us to solve the Dirichlet problem for fully nonlinear equations by the method of continuity (rendering classical solutions).

This theorem was proved independently by N. Krylov [4] and L. C. Evans [3]. In this note, motivated by our work on integral fully nonlinear equations [1], we provide a more direct presentation of their proof (although the underlying key ideas are the same).

We recall the two opposite components in Krylov-Safonov Harnack inequality, the proof of which can be found in [2] (Theorem 4.8).

a) (The weak L^{ε} estimate) If v is a non-negative supersolution of

$$a_{ij}(x)D_{ij}v \le 0$$

in B_1 , with $\lambda I \leq a_{ij} \leq \Lambda I$ then

$$|\{v>t\inf_{B_{1/2}}v\}\cap B_{1/4}|\leq C(\lambda,\Lambda)t^{-\varepsilon}$$

b) (the oscillation lemma) If v is a subsolution of $a_{ij}(x)D_{ij}v \ge 0$ in B_1 and $v \le 1$, then

$$\sup_{B_{1/2}} v \le C(\lambda, \Lambda) |\{v > 0\} \cap B_{3/4}|$$

In case of harmonic functions, these are just consequences of the mean value theorem.

We also recall that convexity of F as a function of D^2u , implies that any pure second derivative, $u_{\sigma\sigma}$, of u and thus any linear combination

$$\ell(x) = \sum_{j} u_{\sigma_j \sigma_j}(x)$$

is a supersolution of the linearized operator

$$a_{ij}(x)D_{ij}\ell(x) \le 0$$

 $(a_{ij}(x) = F_{ij}(D^2u(x)).$

Finally, the uniform ellipticity of F implies that for any two points x_1, x_2 in B_1 ,

(1)
$$\operatorname{tr}[D^2 u(x_2) - D^2 u(x_1)]^+ \approx \operatorname{tr}[D^2 u(x_2) - D^2 u(x_1)]^-$$

At this point, we define for any subspace V

$$w(x,V) = \Delta_V u(x) - \Delta_V u(0)$$

 $(\Delta_V u(x))$ is the Laplacian of u at the point x when restricted to the affine variety x + V.

Note that for each fixed V, w is an $\ell(x)$ as above and satisfies the L^{ε} estimate. Also, note that the *positive* and *negative* part of the laplacian can be expressed as

$$\max_{V} w(x, V) = \operatorname{tr}[D^{2}u(x) - D^{2}u(0)]^{+},$$
$$\min_{V} w(x, V) = -\operatorname{tr}[D^{2}u(x) - D^{2}u(0)]^{-}$$

By rescaling dyadically and iterating it is enough to prove the following lemma:

Lemma 2. There exists a $\theta > 0$, $\theta = \theta(\lambda, \Lambda)$, such that if for all V, for all x in B_1 ,

$$w(x,V) \ge -1$$

Then for all V, for all x in $B_{1/2}$,

$$w(x,V) \ge -1 + \theta.$$

Indeed, this will imply by iteration, that the laplacian is Hölder continuous. Noew we prove the lemma.

Proof. Assume that $w(x_0, V_0) \leq -1 + \theta$ for some V_0 and x_0 in $B_{1/2}$ (θ , small, to be chosen). We will then find a contradiction. Since $w(\cdot, V) + 1$ is a nonnegative supersolution the L^{ε} lemma applies and

$$w(x,V) + 1 \le \theta^{1/2}$$

in a set Ω that covers almost all of $B_{1/4}$, i.e.,

$$|B_{1/4} \setminus \Omega| \le C\theta^{\varepsilon/2}$$

We notice that in Ω , $1 - \theta^{1/2} \leq -w(x, V) \leq \operatorname{tr}[D^2 u(x) - D^2 u(0)]^- \leq 1$. On the other hand, we know that

$$w(x,V) + w(x,V^{\perp}) = \Delta u(x) - \Delta u(0) = \operatorname{tr}[D^2 u(x) - D^2 u(0)]^+ - \operatorname{tr}[D^2 u(x) - D^2 u(0)]^-.$$

Thus, we also have $0 \leq \operatorname{tr}[D^2 u(x) - D^2 u(0)]^+ - w(x, V^{\perp}) \leq \theta^{1/2}$ for $x \in \Omega$. Moreover, for θ small, by (1),

$$-w(x,V) \approx \operatorname{tr}[D^2 u(x) - D^2 u(0)]^- \approx \operatorname{tr}[D^2 u(x) - D^2 u(0)]^+ \approx w(x,V^{\perp}).$$

Thus, there is a constant $c(\lambda, \Lambda) > 0$ such that $w(x, V^{\perp}) \ge c(\lambda, \Lambda)$ in Ω . We now examine the function $v = (c(\lambda, \Lambda) - w(x, V^{\perp}))^+$ in $B_{1/4}$, for which the oscillation lemma applies and satisfies

a) $0 \le v \le 2$ b) $v(0) = c(\lambda, \Lambda)$ c) v = 0 in Ω .

For θ small (i.e., for Ω almost all of $B_{1/4}$) this contradicts the oscillation lemma since $c(\lambda, \Lambda)$ is a fixed positive constant for θ small. This completes the proof.

References

- [1] L. Caffarelli and L. Silvestre. The evans-krylov theorem for non local fully non linear equations. Preprint.
- [2] L.A. Caffarelli and X. Cabre. Fully nonlinear elliptic equations. American Mathematical Society, 1995.
- [3] Lawrence C. Evans. Classical solutions of fully nonlinear, convex, second-order elliptic equations. Comm. Pure Appl. Math., 35(3):333–363, 1982.
- [4] N. V. Krylov. Boundedly inhomogeneous elliptic and parabolic equations. Izv. Akad. Nauk SSSR Ser. Mat., 46(3):487–523, 670, 1982.

Luis Caffarelli, Department of Mathematics, University of Texas at Austin, 1 University Station – C1200, Austin, TX 78712-0257

 $E\text{-}mail\ address: \texttt{caffarel@math.utexas.edu}$

Luis Silvestre, Mathematics Department, University of Chicago, Chicago, IL 60637 $E\text{-}mail\ address:\ luis@math.uchicago.edu$