

SELF-PROPAGATING HIGH TEMPERATURE SYNTHESIS (SHS) IN THE HIGH ACTIVATION ENERGY REGIME

R. MONNEAU AND G.S. WEISS

ABSTRACT. We derive the precise limit of SHS in the high activation energy scaling suggested by B.J. Matkowsky-G.I. Sivashinsky in 1978 and by A. Bayliss-B.J. Matkowsky-A.P. Aldushin in 2002. In the time-increasing case the limit turns out to be the Stefan problem for supercooled water with spatially inhomogeneous coefficients.

Although the present paper leaves open mathematical questions concerning the convergence, our precise form of the limit problem suggest a strikingly simple explanation for the numerically observed pulsating waves.

1. INTRODUCTION

The system

$$(1) \quad \begin{aligned} \partial_t u - \Delta u &= v f(u) \\ \partial_t v &= -v f(u), \end{aligned}$$

where u is the normalized temperature, v is the normalized concentration of the reactant and the non-negative nonlinearity f describes the reaction kinetics, is a simple but widely used model for solid combustion (i.e. the case of the Lewis number being $+\infty$). In particular it is being used to model the industrial process of Self-propagating High temperature Synthesis (SHS). In the case of high activation energy interesting phenomena like the instability of planar waves, fingering and helical waves are observed.

Since the seventies (and possibly even earlier) it has been argued that the problem is for high activation energy related to a Stefan problem describing the freezing of supercooled water (see [21], [10, p. 57]). In [21] B.J. Matkowsky-G.I. Sivashinsky derived a formal singular limit containing a jump condition for the temperature on the interface. Later the Stefan problem for supercooled water – the intuitive limit – became the basis for numerous papers focusing on stability analysis of (1),

Received

2000 *Mathematics Subject Classification*. Primary 80A25, 80A22, 35K55, 35R35.

Key words and phrases. SHS, solid combustion, singular limit, high activation energy, Stefan problem, ill-posed problem, free boundary, hysteresis.

G.S. Weiss has been partially supported by the Grant-in-Aid 15740100 of the Japanese Ministry of Education and partially supported by a fellowship of the Max Planck Society. Both authors thank the Max Planck Institute for Mathematics in the Sciences for the hospitality during their stay in Leipzig.

fingering, helical waves etc. (see for example [11],[12],[9],[14],[13],[15],[8],[1],[2]). Surprisingly there are few *mathematical* results on the subject: In [20] E. Logak-V. Loubeau proved existence of a planar wave in one-space dimension and gave a rigorous proof for convergence as the activation energy goes to infinity. Instability of the planar wave for a special linearization (and high activation energy) is due to [4].

In the present paper we argue that the SHS system converges to the irreversible Stefan problem for supercooled water. As the initial data of the reactant concentration enters the equation as the activation energy goes to infinity, our result also suggests a *surprisingly simple explanation for the numerically observed pulsating waves* (cf. [1] and [2]), namely that they are caused by the spatial inhomogeneity v^0 (or Y^0 , respectively) in the below equation and are therefore mathematically related to the pulsating waves in [3].

In the time-increasing case we give a rigorous convergence proof in higher dimensions. For general initial data in one space-dimension see our forthcoming paper [22].

In the original setting by B.J. Matkowsky-G.I. Sivashinsky [21, equation (2)],

$$(2) \quad \begin{aligned} \partial_t u_N - \Delta u_N &= (1 - \sigma_N) N e^N v_N \exp(-N/u_N), \\ \partial_t v_N &= -N e^N v_N \exp(-N/u_N), \end{aligned}$$

each limit u_∞ of $u_N > 0$ as $N \rightarrow \infty$ satisfies for $(\sigma_N)_{N \in \mathbf{N}} \subset \subset [0, 1)$ (for $\sigma_N \uparrow 1, N \rightarrow \infty$ the limit in this scaling is the solution of the heat equation; cf. Section 5.1 and Theorem 4.1)

$$(3) \quad \partial_t u_\infty - v^0 \partial_t \chi = \Delta u_\infty \text{ in } (0, +\infty) \times \Omega,$$

where v^0 are the initial data of v_∞ and

$$\chi(t, x) \begin{cases} \in [0, 1], & \text{esssup}_{(0,t)} u_\infty(\cdot, x) \leq 1, \\ = 1, & \text{esssup}_{(0,t)} u_\infty(\cdot, x) > 1, \end{cases}$$

and in the time-increasing case,

$$\chi(t, x) \begin{cases} = 0, & u_\infty(t, x) < 1, \\ \in [0, 1], & u_\infty(t, x) = 1, \\ = 1, & u_\infty(t, x) > 1. \end{cases}$$

In the SHS system with another scaling and a temperature threshold (see [2, p. 109-110]),

$$(4) \quad \begin{aligned} \partial_t \theta_N - \Delta \theta_N &= (1 - \sigma_N) N Y_N \exp((N(1 - \sigma_N)(\theta_N - 1))/(\sigma_N + (1 - \sigma_N)\theta_N)) \chi_{\{\theta_N > \bar{\theta}\}}, \\ \partial_t Y_N &= -(1 - \sigma_N) N Y_N \exp((N(1 - \sigma_N)(\theta_N - 1))/(\sigma_N + (1 - \sigma_N)\theta_N)) \chi_{\{\theta_N > \bar{\theta}\}}, \end{aligned}$$

where $N(1 - \sigma_N) \gg 1, \sigma_N \in (0, 1)$ and $\bar{\theta} \in (0, 1)$, each limit θ_∞ of θ_N satisfies (cf. Section 5.2 and Theorem 4.1)

$$(5) \quad \partial_t \theta_\infty - Y^0 \partial_t \chi = \Delta \theta_\infty \text{ in } (0, +\infty) \times \Omega,$$

where Y^0 are the initial data of Y_∞ and

$$\chi(t, x) \begin{cases} \in [0, 1], & \text{esssup}_{(0,t)} \theta_\infty(\cdot, x) \leq 1, \\ = 1, & \text{esssup}_{(0,t)} \theta_\infty(\cdot, x) > 1, \end{cases}$$

and in the time-increasing case,

$$\chi(t, x) \begin{cases} = 0, & \theta_\infty(t, x) < 1, \\ \in [0, 1], & \theta_\infty(t, x) = 1, \\ = 1, & \theta_\infty(t, x) > 1. \end{cases}$$

To our knowledge this precise form of the limit problem, i.e. the equation with the discontinuous hysteresis term, has not been known. Even in the time-increasing case it does not coincide with the formal result in [21].

In the case that θ_∞ (or u_∞ , respectively) is increasing in time and v^0 (or Y^0 , respectively) is constant, our limit problem coincides with the Stefan problem for supercooled water, an extensively studied ill-posed problem (for a survey see [5]). As it is a forward-backward parabolic equation it is not clear whether one should expect uniqueness (see [6, Remark 7.2] for an example of non-uniqueness in a related problem).

On the positive side, much more is known about the Stefan problem for supercooled water than the SHS system, e.g. existence of a finger ([16]), instability of the finger ([19]), one-phase solutions ([6]); those results, when combined with our convergence result, suggest that similar properties should be true for the SHS system.

It is interesting to observe that even in the time-increasing case our singular limit *selects certain solutions* of the Stefan problem for supercooled water. For example, $u(t) = (\kappa - 1)\chi_{\{t < 1\}} + \kappa\chi_{\{t > 1\}}$ is for each $\kappa \in (0, 1)$ a perfectly valid solution of the Stefan problem for supercooled water, but, as easily verified, it cannot be obtained from the ODE

$$\partial_t u_\epsilon(t) = -\partial_t \exp\left(-\frac{1}{\epsilon} \int_0^t \exp((1 - 1/(u_\epsilon(s) + 1))/\epsilon) ds\right) \text{ as } \epsilon \rightarrow 0.$$

2. NOTATION

Throughout this article \mathbf{R}^n will be equipped with the Euclidean inner product $x \cdot y$ and the induced norm $|x|$. $B_r(x)$ will denote the open n -dimensional ball of center x , radius r and volume $r^n \omega_n$. When the center is not specified, it is assumed to be 0.

When considering a set A , χ_A shall stand for the characteristic function of A , while ν shall typically denote the outward normal to a given boundary. The operator ∂_t will mean the partial derivative of a function in the time direction, Δ the Laplacian in the space variables and \mathcal{L}^n the n -dimensional Lebesgue measure.

Finally $\mathbf{W}_p^{2,1}$ denotes the parabolic Sobolev space as defined in [18].

3. PRELIMINARIES

In what follows, Ω is a bounded C^1 -domain in \mathbf{R}^n and

$$u_\epsilon \in \bigcap_{T \in (0, +\infty)} \mathbf{W}_2^{2,1}((0, T) \times \Omega)$$

is a strong solution of the equation

$$(6) \quad \begin{aligned} \partial_t u_\epsilon(t, x) - \Delta u_\epsilon(t, x) &= -v_\epsilon^0(x) \partial_t \exp\left(-\frac{1}{\epsilon} \int_0^t g_\epsilon(u_\epsilon(s, x)) ds\right), \\ u_\epsilon(0, \cdot) &= u_\epsilon^0 \text{ in } \Omega, \nabla u_\epsilon \cdot \nu = 0 \text{ on } (0, +\infty) \times \partial\Omega; \end{aligned}$$

here g_ϵ is a non-negative function on \mathbf{R} satisfying:

0) g_ϵ is for each $\epsilon \in (0, 1)$ piecewise continuous with only one possible jump at z_0 , $g_\epsilon(z_0-) = g_\epsilon(z_0) = 0$ in case of a jump, and g_ϵ satisfies for each $\epsilon \in (0, 1)$ and for every $z \in \mathbf{R}$ the bound $g_\epsilon(z) \leq C_\epsilon(1 + |z|)$.

1) $g_\epsilon/\epsilon \rightarrow 0$ as $\epsilon \rightarrow 0$ on each compact subset of $(-\infty, 0)$.

2) for each compact subset K of $(0, +\infty)$ there is $c_K > 0$ such that $\min(g_\epsilon, c_K) \rightarrow c_K$ uniformly on K as $\epsilon \rightarrow 0$.

The initial data satisfy $0 \leq v_\epsilon^0 \leq C < +\infty$, v_ϵ^0 converges in $L^1(\Omega)$ to v^0 as $\epsilon \rightarrow 0$, $(u_\epsilon^0)_{\epsilon \in (0, 1)}$ is bounded in $L^2(\Omega)$, it is uniformly bounded from below by a constant u_{\min} , and it converges in $L^1(\Omega)$ to u^0 as $\epsilon \rightarrow 0$.

Remark 3.1. *Assumption 0) guarantees existence of a global strong solution for each $\epsilon \in (0, 1)$.*

4. THE HIGH ACTIVATION ENERGY LIMIT

Theorem 4.1. *The family $(u_\epsilon)_{\epsilon \in (0, 1)}$ is for each $T \in (0, +\infty)$ precompact in $L^1((0, T) \times \Omega)$, and each limit u of $(u_\epsilon)_{\epsilon \in (0, 1)}$ as a sequence $\epsilon_m \rightarrow 0$, satisfies in the sense of distributions the initial-boundary value problem*

$$(7) \quad \begin{aligned} \partial_t u - v^0 \partial_t \chi &= \Delta u \text{ in } (0, +\infty) \times \Omega, \\ u(0, \cdot) &= u^0 + v^0 H(u^0) \text{ in } \Omega, \nabla u \cdot \nu = 0 \text{ on } (0, +\infty) \times \partial\Omega, \\ \text{where } \chi(t, x) &\begin{cases} \in [0, 1], & \text{esssup}_{(0, t)} u(\cdot, x) \leq 0, \\ = 1, & \text{esssup}_{(0, t)} u(\cdot, x) > 0, \end{cases} \end{aligned}$$

and H is the maximal monotone graph

$$H(z) \begin{cases} = 0, & z < 0, \\ \in [0, 1], & z = 0, \\ = 1, & z > 0. \end{cases}$$

Moreover, χ is increasing in time and u is a supercaloric function.

If $(u_\epsilon)_{\epsilon \in (0, 1)}$ satisfies $\partial_t u_\epsilon \geq 0$ in $(0, T) \times \Omega$, then u is a solution of the Stefan problem for supercooled water, i.e.

$$\partial_t u - v^0 \partial_t H(u) = \Delta u \text{ in } (0, +\infty) \times \Omega.$$

Remark 4.2. *Note that assumption 1) is only needed to prove the second statement “If”.*

Proof. Step 0 (Uniform Bound from below):

Since u_ϵ is supercaloric, it is bounded from below by the constant u_{\min} .

Step 1 ($L^2((0, T) \times \Omega)$ -Bound):

The time-integrated function $v_\epsilon(t, x) := \int_0^t u_\epsilon(s, x) ds$, satisfies

$$(8) \quad \partial_t v_\epsilon(t, x) - \Delta v_\epsilon(t, x) = w_\epsilon(t, x) + u_\epsilon^0(x)$$

where w_ϵ is a measurable function satisfying $0 \leq w_\epsilon \leq C$. Consequently

$$\begin{aligned} \int_0^T \int_\Omega (\partial_t v_\epsilon)^2 + \frac{1}{2} \int_\Omega |\nabla v_\epsilon|^2(T) &= \int_0^T \int_\Omega (w_\epsilon + u_\epsilon^0) \partial_t v_\epsilon \\ &\leq \frac{1}{2} \int_0^T \int_\Omega (\partial_t v_\epsilon)^2 + \frac{T}{2} \int_\Omega (C + |u_\epsilon^0|)^2, \end{aligned}$$

implying

$$(9) \quad \int_0^T \int_\Omega u_\epsilon^2 \leq T \int_\Omega (C + |u_\epsilon^0|)^2.$$

Step 2 ($L^2((0, T) \times \Omega)$ -Bound for $\nabla \min(u_\epsilon, M)$):

For

$$G_M(z) := \begin{cases} z^2/2, & z < M, \\ Mz - M^2/2, & z \geq M, \end{cases}$$

and any $M \in \mathbf{N}$,

$$\begin{aligned} &\int_\Omega G_M(u_\epsilon) - G_M(u_\epsilon^0) + \int_0^T \int_\Omega |\nabla \min(u_\epsilon, M)|^2 \\ &= \int_0^T \int_\Omega -v_\epsilon^0 \min(u_\epsilon, M) \partial_t \exp\left(-\frac{1}{\epsilon} \int_0^t g_\epsilon(u_\epsilon(s, x)) ds\right). \end{aligned}$$

As $\partial_t \exp\left(-\frac{1}{\epsilon} \int_0^t g_\epsilon(u_\epsilon(s, x)) ds\right) \leq 0$, we know that $\partial_t \exp\left(-\frac{1}{\epsilon} \int_0^t g_\epsilon(u_\epsilon(s, x)) ds\right)$ is bounded in $L^\infty(\Omega; L^1((0, T)))$, and

$$\begin{aligned} &\int_0^T \int_\Omega -v_\epsilon^0 \min(u_\epsilon, M) \partial_t \exp\left(-\frac{1}{\epsilon} \int_0^t g_\epsilon(u_\epsilon(s, x)) ds\right) \\ &\leq C \int_{\Omega(0, T)} \sup \max(\min(u_\epsilon, M), 0) \leq CM \mathcal{L}^n(\Omega). \end{aligned}$$

Step 3 (Compactness):

Let $\chi_M : \mathbf{R} \rightarrow \mathbf{R}$ be a smooth non-increasing function satisfying $\chi_{(-\infty, M-1)} \leq \chi_M \leq \chi_{(-\infty, M)}$ and let Φ_M be the primitive such that $\Phi_M(z) = z$ for $z \leq M-1$ and $\Phi_M \leq M$. Moreover, let $(\phi_\delta)_{\delta \in (0, 1)}$ be a family of mollifiers, i.e. $\phi_\delta \in C_0^{0,1}(\mathbf{R}^n; [0, +\infty))$ such that $\int \phi_\delta = 1$ and $\text{supp } \phi_\delta \subset B_\delta(0)$. Then, if we extend u_ϵ and v_ϵ^0 by the value 0 to the whole of $(0, +\infty) \times \mathbf{R}^n$, we obtain by the homogeneous Neumann data of u_ϵ that

$$\begin{aligned} &\partial_t (\Phi_M(u_\epsilon) * \phi_\delta)(t, x) \\ &= \left(\left(\chi_M(u_\epsilon) \left(\chi_\Omega \Delta u_\epsilon - v_\epsilon^0 \partial_t \exp\left(-\frac{1}{\epsilon} \int_0^t g_\epsilon(u_\epsilon(s, x)) ds\right) \right) \right) * \phi_\delta \right)(t, x) \end{aligned}$$

$$\begin{aligned}
&= \int_{\mathbf{R}^n} \chi_M(u_\epsilon)(t, y) \left(\chi_\Omega(y) \Delta u_\epsilon(t, y) \right. \\
&\quad \left. - v_\epsilon^0(y) \partial_t \exp\left(-\frac{1}{\epsilon} \int_0^t g_\epsilon(u_\epsilon(s, y)) ds\right) \right) \phi_\delta(x - y) dy \\
&= \int_{\mathbf{R}^n} \phi_\delta(x - y) \left(-\chi'_M(u_\epsilon(t, y)) \chi_\Omega(y) |\nabla u_\epsilon(t, y)|^2 \right. \\
&\quad \left. - \chi_M(u_\epsilon(t, y)) v_\epsilon^0(y) \partial_t \exp\left(-\frac{1}{\epsilon} \int_0^t g_\epsilon(u_\epsilon(s, y)) ds\right) \right. \\
&\quad \left. + \chi_M(u_\epsilon(t, y)) \chi_\Omega(y) \nabla u_\epsilon(t, y) \cdot \nabla \phi_\delta(x - y) \right) dy .
\end{aligned}$$

Consequently

$$\int_0^T \int_{\mathbf{R}^n} |\partial_t (\Phi_M(u_\epsilon) * \phi_\delta)| \leq C_1(\Omega, C, M, \delta, T)$$

and

$$\int_0^T \int_{\mathbf{R}^n} |\nabla (\Phi_M(u_\epsilon) * \phi_\delta)| \leq C_2(\Omega, M, \delta, T) .$$

It follows that $(\Phi_M(u_\epsilon) * \phi_\delta)_{\epsilon \in (0,1)}$ is for each (M, δ, T) precompact in $L^1((0, T) \times \mathbf{R}^n)$.

On the other hand

$$\begin{aligned}
\int_0^T \int_{\mathbf{R}^n} |\Phi_M(u_\epsilon) * \phi_\delta - \Phi_M(u_\epsilon)| &\leq C_3 \left(\delta^2 \int_0^T \int_{\Omega} |\nabla \Phi_M(u_\epsilon)|^2 \right)^{\frac{1}{2}} \\
&+ 2(M - u_{\min}) T \mathcal{L}^n(B_\delta(\partial\Omega)) \leq C_4(C, \Omega, u_{\min}, M, T) \delta .
\end{aligned}$$

Combining this estimate with the precompactness of $(\Phi_M(u_\epsilon) * \phi_\delta)_{\epsilon \in (0,1)}$ we obtain that $\Phi_M(u_\epsilon)$ is for each (M, T) precompact in $L^1((0, T) \times \mathbf{R}^n)$. Thus, by a diagonal sequence argument, we may take a sequence $\epsilon_m \rightarrow 0$ such that $\Phi_M(u_{\epsilon_m}) \rightarrow z_M$ a.e. in $(0, +\infty) \times \mathbf{R}^n$ as $m \rightarrow \infty$, for every $M \in \mathbf{N}$. At a.e. point of the set $\{z_M < M - 1\}$, u_{ϵ_m} converges to z_M . At each point (t, x) of the remainder $\bigcap_{M \in \mathbf{N}} \{z_M \geq M - 1\}$, the value $u_{\epsilon_m}(t, x)$ must for large m (depending on (M, t, x)) be larger than $M - 2$. But that means that on the set $\bigcap_{M \in \mathbf{N}} \{z_M \geq M - 1\}$, the sequence $(u_{\epsilon_m})_{m \in \mathbf{N}}$ converges a.e. to $+\infty$. It follows that $(u_{\epsilon_m})_{m \in \mathbf{N}}$ converges a.e. in $(0, +\infty) \times \Omega$ to a function $z : (0, +\infty) \times \Omega \rightarrow \mathbf{R} \cup \{+\infty\}$. But then, as $(u_{\epsilon_m})_{m \in \mathbf{N}}$ is for each $T \in (0, +\infty)$ bounded in $L^2((0, T) \times \Omega)$, $(u_{\epsilon_m})_{m \in \mathbf{N}}$ converges by Vitali's theorem (stating that a.e. convergence and a non-concentration condition in L^p imply in bounded domains L^p -convergence) for each $p \in [1, 2)$ in $L^p((0, T) \times \Omega)$ to the weak L^2 -limit u of $(u_{\epsilon_m})_{m \in \mathbf{N}}$. It follows that $\mathcal{L}^{n+1}(\bigcap_{M \in \mathbf{N}} \{z_M \geq M - 1\}) = \mathcal{L}^{n+1}(\{u = +\infty\}) = 0$.

Step 4 (Identification of the Limit Equation in $\text{esssup}_{(0,t)} u > 0$):

Let us consider $(t, x) \in (0, +\infty) \times \Omega$ such that $u_{\epsilon_m}(s, x) \rightarrow u(s, x)$ for a.e. $s \in (0, t)$ and $u(\cdot, x) \in L^2((0, t))$. In the case $\text{esssup}_{(0,t)} u(\cdot, x) > 0$, we obtain by Egorov's theorem and assumption 2) that $\exp\left(-\frac{1}{\epsilon_m} \int_0^t g_{\epsilon_m}(u_{\epsilon_m}(s, x)) ds\right) \rightarrow 0$ as $m \rightarrow \infty$.

Step 5 (The case $\partial_t u_\epsilon \geq 0$):

Let (t, x) be such that $u_{\epsilon_m}(t, x) \rightarrow u(t, x) = \lambda < 0$: Then by assumption 1),

$$\exp\left(-\frac{1}{\epsilon_m} \int_0^t g_{\epsilon_m}(u_{\epsilon_m}(s, x)) ds\right) \geq \exp\left(-t \frac{\max_{[u_{\min}, \lambda/2]} g_{\epsilon_m}}{\epsilon_m}\right) \rightarrow 1 \text{ as } m \rightarrow \infty .$$

Remark 4.3. 1) For a more general result in one space-dimension see the forthcoming paper [22].

2) We also obtain a rigorous convergence result in the case of (higher dimensional) traveling waves with suitable conditions at infinity. In this case our $L^2(W^{1,2})$ -estimate (Step 2) implies a no-concentration property of the time-derivative.

5. APPLICATIONS

Although the limit equation is an ill-posed problem, the convergence to the limit seems to be robust with respect to perturbations of the ϵ -system and the scaling: here we mention two examples of different systems leading to the same limit. Other examples can be found in mathematical biology (see [17] and [23]).

5.1. The Matkowsky-Sivashinsky scaling

We apply our result to the scaling in [21, equation (2)], i.e.

$$(10) \quad \begin{aligned} \partial_t u_N - \Delta u_N &= (1 - \sigma_N) N v_N \exp(N(1 - 1/u_N)), \\ \partial_t v_N &= -N v_N \exp(N(1 - 1/u_N)), \end{aligned}$$

where the normalized temperature u_N and the normalized concentration v_N are non-negative, $(\sigma_N)_{N \in \mathbf{N}} \subset \subset [0, 1]$ (in the case $\sigma_N \uparrow 1, N \rightarrow \infty$ the limit equation in the scaling as it would be the heat equation, but we could still apply our result to $u_N/(1 - \sigma_N)$) and the activation energy $N \rightarrow \infty$.

Setting $u_{\min} := -1, \epsilon := 1/N, u_\epsilon := u_N - 1$ and

$$g_\epsilon(z) := \begin{cases} \exp((1 - 1/(z + 1))/\epsilon), & z > -1 \\ 0, & z \leq -1 \end{cases}$$

and integrating the equation for v_N in time, we see that the assumptions of Theorem 4.1 are satisfied and we obtain that each limit u_∞, σ_∞ of u_N, σ_N satisfies

$$(11) \quad \partial_t u_\infty - (1 - \sigma_\infty) v^0 \partial_t \chi = \Delta u_\infty \text{ in } (0, +\infty) \times \Omega,$$

where

$$\chi(t, x) \begin{cases} \in [0, 1], & \text{esssup}_{(0,t)} u_\infty(\cdot, x) \leq 1, \\ = 1, & \text{esssup}_{(0,t)} u_\infty(\cdot, x) > 1, \end{cases}$$

and in the time-increasing case,

$$(12) \quad \partial_t u_\infty - (1 - \sigma_\infty) v^0 \partial_t H(u_\infty) = \Delta u_\infty \text{ in } (0, +\infty) \times \Omega,$$

$$u_\infty(0, \cdot) = u^0 + v^0 H(u^0) \text{ in } \Omega, \quad \nabla u_\infty \cdot \nu = 0 \text{ on } (0, +\infty) \times \partial\Omega,$$

where v^0 are the initial data of v_∞ . Moreover, χ is increasing in time and u_∞ is a supercaloric function.

5.2. SHS in another scaling with temperature threshold

Here we consider (cf. [2, p. 109-110]), i.e.

$$(13) \quad \begin{aligned} & \partial_t \theta_N - \Delta \theta_N \\ &= (1 - \sigma_N) N Y_N \exp((N(1 - \sigma_N)(\theta_N - 1))/(\sigma_N + (1 - \sigma_N)\theta_N)) \chi_{\{\theta_N > \bar{\theta}\}}, \\ & \partial_t Y_N = -(1 - \sigma_N) N Y_N \exp((N(1 - \sigma_N)(\theta_N - 1))/(\sigma_N + (1 - \sigma_N)\theta_N)) \chi_{\{\theta_N > \bar{\theta}\}} \end{aligned}$$

where $N(1 - \sigma_N) \gg 1$, $\sigma_N \in (0, 1)$ and the constant $\bar{\theta} \in (0, 1)$ is a threshold parameter at which the reaction sets in.

Setting $u_{\min} = -1$, $\epsilon := 1/(N(1 - \sigma_N))$, $\kappa(\epsilon) := 1 - \sigma_N$, $u_\epsilon := \theta_N - 1$,

$$g_\epsilon(z) := \begin{cases} \exp((z/(\kappa(\epsilon)z + 1))/\epsilon), & z > \bar{\theta} - 1 \\ 0, & z \leq \bar{\theta} - 1 \end{cases}$$

and integrating the equation for Y_N in time, we see that the assumptions of Theorem 4.1 are satisfied and we obtain that each limit u_∞ of u_N satisfies

$$(14) \quad \begin{aligned} & \partial_t u_\infty - v^0 \partial_t \chi = \Delta u_\infty \text{ in } (0, +\infty) \times \Omega, \\ & \chi(t, x) \begin{cases} \in [0, 1], & \text{esssup}_{(0,t)} u_\infty(\cdot, x) \leq 1, \\ = 1, & \text{esssup}_{(0,t)} u_\infty(\cdot, x) > 1, \end{cases} \end{aligned}$$

and in the time-increasing case,

$$(15) \quad \begin{aligned} & \partial_t u_\infty - v^0 \partial_t H(u_\infty) = \Delta u_\infty \text{ in } (0, +\infty) \times \Omega, \\ & u_\infty(0, \cdot) = u^0 + v^0 H(u^0) \text{ in } \Omega, \quad \nabla u_\infty \cdot \nu = 0 \text{ on } (0, +\infty) \times \partial\Omega, \end{aligned}$$

where v^0 are the initial data of v_∞ . Moreover, χ is increasing in time and u_∞ is a supercaloric function.

6. OPEN QUESTIONS

The most pressing task is of course to study the existence or non-existence of “peaking” (cf. Figure 1) of the solution in the negative phase (for the case of one




Figure 1. Is it possible for the solution to have a tiny peak traveling at high speed?.

space dimension see the forthcoming paper [22]). A related question is whether

$(u_\epsilon)_{\epsilon \in (0,1)}$ is bounded in L^∞ in the case of uniformly bounded initial data. Although this seems obvious, it is not obvious how to prevent concentration close to the interface.

Uniqueness for the limit problem (the irreversible Stefan problem for supercooled water) in general seems unlikely. One might however ask whether time-global uniqueness holds in the case that u is strictly increasing in the x_1 -direction. By the result in [7] for the ill-posed Hele-Shaw problem, time-local uniqueness is likely to be true here, too.

Acknowledgment. We thank Stephan Luckhaus, Mayan Mimura, Stefan Müller, and Juan J.L. Velázquez for discussions.

REFERENCES

1. A. P. Aldushin, A. Bayliss, and B. J. Matkowsky. Dynamics in layer models of solid flame propagation. *Phys. D*, 143(1-4):109–137, 2000.
2. A. Bayliss, B. J. Matkowsky, and A. P. Aldushin. Dynamics of hot spots in solid fuel combustion. *Phys. D*, 166(1-2):104–130, 2002.
3. Henri Berestycki and François Hamel. Front propagation in periodic excitable media. *Comm. Pure Appl. Math.*, 55(8):949–1032, 2002.
4. Alexis Bonnet and Elisabeth Logak. Instability of travelling waves in solid combustion for high activation energy. *Preprint*.
5. J. N. Dewynne. A survey of supercooled Stefan problems. In *Mini-Conference on Free and Moving Boundary and Diffusion Problems (Canberra, 1990)*, volume 30 of *Proc. Centre Math. Appl. Austral. Nat. Univ.*, pages 42–56. Austral. Nat. Univ., Canberra, 1992.
6. Emmanuele DiBenedetto and Avner Friedman. The ill-posed Hele-Shaw model and the Stefan problem for supercooled water. *Trans. Amer. Math. Soc.*, 282(1):183–204, 1984.
7. Jean Duchon and Raoul Robert. Évolution d’une interface par capillarité et diffusion de volume. I. Existence locale en temps. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 1(5):361–378, 1984.
8. M. Frankel, L. K. Gross, and V. Roytburd. Thermo-kinetically controlled pattern selection. *Interfaces Free Bound.*, 2(3):313–330, 2000.
9. M. Frankel, V. Roytburd, and G. Sivashinsky. A sequence of period doublings and chaotic pulsations in a free boundary problem modeling thermal instabilities. *SIAM J. Appl. Math.*, 54(4):1101–1112, 1994.
10. M. L. Frankel. On the nonlinear evolution of a solid-liquid interface. *Phys. Lett. A*, 128(1-2):57–60, 1988.
11. M. L. Frankel. On the nonlinear evolution of a solid-liquid interface. *Phys. Lett. A*, 128(1-2):57–60, 1988.
12. M. L. Frankel. On a free boundary problem associated with combustion and solidification. *RAIRO Modél. Math. Anal. Numér.*, 23(2):283–291, 1989.
13. Michael L. Frankel and Victor Roytburd. A free boundary problem modeling thermal instabilities: stability and bifurcation. *J. Dynam. Differential Equations*, 6(3):447–486, 1994.
14. Michael L. Frankel and Victor Roytburd. A free boundary problem modeling thermal instabilities: well-posedness. *SIAM J. Math. Anal.*, 25(5):1357–1374, 1994.
15. Michael L. Frankel and Victor Roytburd. On a free boundary model related to solid-state combustion. *Comm. Appl. Nonlinear Anal.*, 2(3):1–22, 1995.
16. G.P. Ivantsov. *Dokl. Akad. Nauk SSSR*, 58:567–569, 1947.
17. K. Kawasaki, A. Mochizuki, M. Matsushita, T. Umeda, and N. Shigesada. Modeling spatio-temporal patterns generated by bacillus subtilis. *J. theor. Biol.*, 188:177–185, 1997.

18. O. A. Ladyženskaja, V. A. Solonnikov, and N. N. Ural'ceva. *Linear and quasilinear equations of parabolic type*. Translated from the Russian by S. Smith. Translations of Mathematical Monographs, Vol. 23. American Mathematical Society, Providence, R.I., 1967.
19. J.S. Langer. Instabilities and pattern formation in crystal growth. *Rev. Mod. Phys.*, 52:1–28, 1980.
20. Elisabeth Logak and Vincent Loubeau. Travelling wave solutions to a condensed phase combustion model. *Asymptotic Anal.*, 12(4):259–294, 1996.
21. B. J. Matkowsky and G. I. Sivashinsky. Propagation of a pulsating reaction front in solid fuel combustion. *SIAM J. Appl. Math.*, 35(3):465–478, 1978.
22. Régis Monneau and G.S. Weiss. Self-propagating high temperature synthesis in the high activation energy regime – the general problem in one space dimension. *In preparation*.
23. R. A. Satnoianu, P. K. Maini, F. S. Garduno, and J. P. Armitage. Travelling waves in a nonlinear degenerate diffusion model for bacterial pattern formation. *Discrete Contin. Dyn. Syst. Ser. B*, 1(3):339–362, 2001.

R. Monneau, Ecole Nationale des Ponts et Chaussées, CERMICS, 6 et 8 avenue Blaise Pascal, Cité Descartes Champs-sur-Marne, 77455 Marne-la-Vallée Cedex 2, France, *e-mail*: monneau@cermics.enpc.fr,
<http://cermics.enpc.fr/monneau/home.html>

G.S. Weiss, Graduate School of Mathematical Sciences, University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo, 153-8914 Japan, *current address*: until Sept. 2005 guest of the Max Planck Institute for Mathematics in the Sciences, Inselstr. 22, D-04103 Leipzig, German, *e-mail*: gw@ms.u-tokyo.ac.jp,
<http://www.ms.u-tokyo.ac.jp/gw>