

A reduced stochastic model for shock and detonation waves

Jean-Bernard Maillet¹, Laurent Soulard¹, [Gabriel STOLTZ](#)^{1,2}

¹ CEA/DAM (Bruyères-le-Châtel, France)

² CERMICS, ENPC (Marne-la-Vallée, France)


`http://cermics.enpc.fr/~stoltz/`

- (All atom) Hamiltonian dynamics

$$\begin{cases} dq = M^{-1}p dt \\ dp = -\nabla V(q) dt \end{cases}$$

- Hamiltonian $H(q, p) = \frac{1}{2}p^T M^{-1}p + V(q)$
- All the physics is contained in V !
- Shock obtained through a [piston compression](#)
- Bond order potentials such as REBO and ReaxFF are now routinely used and the simulations are qualitatively correct
- Problem = reachable time (ns) and space (μm) scales are not large enough... Ultimately, not all microscopic details are relevant!

A typical simulation



Shock wave in a Lennard-Jones fluid (Hamiltonian dynamics): piston compression + relaxation

Reducing the complexity of the system

- Replace **deterministic all atom** dynamics by a **stochastic** dynamics on the variables of interest
- General strategy (Mori-Zwanzig) → average over the unrelevant degrees of freedom to eliminate them: replace their influence by some **mean action (drift)** and **fluctuations** around the mean behavior (**random noise**)
- In this context:
 - 1D model of shock waves in crystalline solids^a
 - Replace a complex by molecule by a center of mass with some internal energy (unresolved internal modes)^b

^aG. Stoltz, *Nonlinearity* **18**, 1967-1985 (2005)

^bStrachan and Holian, *Phys. Rev. Lett.* (2005)

Reduced dynamics: the inert case

- Langevin dynamics (e.g. implicit solvents in biology)

$$\begin{cases} dq = M^{-1}p dt \\ dp = -\nabla V(q) dt - \gamma M^{-1}p dt + \sigma dW_t \end{cases}$$

- Fluctuation/dissipation relation

$$\sigma^2 = 2\gamma k_B \bar{T} = \frac{2\gamma}{\beta}$$

ensures that the canonical measure is preserved

- Cannot be used for the simulation of shock waves:
 - the dynamics is **not invariant** through a Galilean transform;
 - the temperature **is fixed** *a priori*.

(Almost) Dissipative Particle Dynamics

- Galilean invariance → **DPD philosophy**^{a,b}
- Friction depending on the relative velocities (with some cut-off):

$$\begin{cases} dq_i = \frac{p_i}{m_i} dt \\ dp_i = \sum_{j \neq i} -\nabla V(r_{ij}) dt - \gamma \chi^2(r_{ij}) v_{ij} dt + \sqrt{\frac{2\gamma}{\beta}} \chi(r_{ij}) dW_{ij} \end{cases}$$

- relative distances $r_{ij} = |q_i - q_j|$, relative velocities $v_{ij} = \frac{p_i}{m_i} - \frac{p_j}{m_j}$
- (radial) weight function χ with support $[0, r_c]$
- **antisymmetric** standard Brownian motions $W_{ij} = -W_{ji}$.
- No projection along the lines of center

^aHoogerbrugge and Koelman, *Europhys. Lett.* **19**(3), 155–160 (1992)

^bEspanol and Warren, *Europhys. Lett.*, **30**(4), 191–196, (1995)

(Almost) Dissipative Particle Dynamics (2)

- Preserve the **canonical measure** (cf. Fokker-Planck equation)
- **Ergodicity** in a 1D framework when density high enough^a
- Numerical integration through a **splitting** strategy^b:
 - Verlet step
 - loop over particles for a Verlet-like treatment of the dissipation

$$\forall i < j, \left\{ \begin{array}{l} p_i^{n+1/2} = p_i^n - \frac{1}{2} \gamma \chi^2(r_{ij}) v_{ij}^n + \frac{1}{2} \sigma \sqrt{\Delta t} \chi(r_{ij}) U_{ij}^n, \\ p_j^{n+1/2} = p_j^n + \frac{1}{2} \gamma \chi^2(r_{ij}) v_{ij}^n - \frac{1}{2} \sigma \sqrt{\Delta t} \chi(r_{ij}) U_{ij}^n, \\ p_i^{n+1} = p_i^{n+1/2} - \frac{1}{2} \gamma \chi^2(r_{ij}) v_{ij}^{n+1} + \frac{1}{2} \sigma \sqrt{\Delta t} \chi(r_{ij}) U_{ij}^n, \\ p_j^{n+1} = p_j^{n+1/2} + \frac{1}{2} \gamma \chi^2(r_{ij}) v_{ij}^{n+1} - \frac{1}{2} \sigma \sqrt{\Delta t} \chi(r_{ij}) U_{ij}^n, \end{array} \right.$$

^aShardlow and Yan, *Stochastics and dynamics* **6**(1) (2006)

^bT. Shardlow, *SIAM J. Sci. Comput.* **24**(4) (2003) 1267-1282

- Variation of temperature when the shock passes? Energy conservation: **transfers between external and internal modes** (DPDE^{a,b} philosophy)
- Replace a complex by molecule by^c
 - a center of mass, effective interactions $H(q, p)$
 - an internal energy ϵ = all the unresolved (internal) modes
- Evolution such that $dH(q, p) + \sum_i d\epsilon_i = 0$
- Microscopic state law: entropy $s = s(\epsilon)$, internal temperature defined from the entropy as $T_i = \left(\frac{\partial s_i}{\partial \epsilon_i} \right)^{-1}$
- Harmonic internal degrees of freedom: $T(\epsilon) = \frac{\epsilon}{C_v}$

^aAvalos and Mackie, *Europhys. Lett.* **40**, 141-146 (1997)

^bEspañol, *Europhys. Lett.* **40** 631-636 (1997)

^cStrachan and Holian, *Phys. Rev. Lett.* (2005)

- **Store energy** dissipated by the frictional forces in the internal energies

$$\left\{ \begin{array}{l} dq_i = \frac{p_i}{m_i} dt \\ dp_i = \sum_{j, j \neq i} -\nabla V(r_{ij}) dt - \gamma_{ij} \chi^2(r_{ij}) v_{ij} dt + \sigma_{ij} \chi(r_{ij}) dW_{ij}, \\ d\epsilon_i = \frac{1}{2} \sum_{j, j \neq i} \left(\chi^2(r_{ij}) \gamma_{ij} v_{ij}^2 - \frac{\sigma_{ij}^2}{2} \left(\frac{1}{m_i} + \frac{1}{m_j} \right) \chi^2(r_{ij}) \right) dt \\ \quad - \sigma_{ij} \chi(r_{ij}) v_{ij} \cdot dW_{ij}, \end{array} \right.$$

- Fluctuation-dissipation relation depending on the **internal temperatures**:

$$\sigma_{ij} = \sigma^2 \text{ and } \gamma_{ij} = \frac{\sigma^2}{\beta_{ij}} \text{ with } \beta_{ij} = \frac{1}{2k_B} \left(\frac{1}{T_i} + \frac{1}{T_j} \right) \text{ to preserve the measure}$$

$$d\mu(q, p, \epsilon) = \frac{1}{Z} e^{-\beta(H(q,p)+f(\epsilon))} dq dp d\epsilon$$

where $f(\epsilon) = \epsilon - \bar{T}s(\epsilon)$ is a free energy

- Existence of invariants: **total energy** and **total momentum**

- Numerical scheme based on **splitting**

- Hamiltonian part $\left\{ \begin{array}{l} dq = M^{-1}p dt, \\ dp = -\nabla V(q) dt \end{array} \right.$, velocity Verlet scheme


- stochastic part:

$$\forall i < j, \left\{ \begin{array}{l} dp_i = -\gamma_{ij}\chi^2(r_{ij})v_{ij} dt + \sigma\chi(r_{ij}) dW_{ij}, \\ dp_j = -dp_i, \\ d\epsilon_i = -\frac{1}{2}d\left(\frac{p_i^2}{2m_i} + \frac{p_j^2}{2m_j}\right), \\ d\epsilon_j = d\epsilon_i. \end{array} \right.$$

- Estimators of the thermodynamic temperature

$$k_B \langle T_{\text{kin}} \rangle = \beta^{-1}, \quad k_B \left(\left\langle \frac{1}{T_{\text{int}}} \right\rangle \right)^{-1} = \beta^{-1}$$

An example (PVDF)



Average temperature in a slice: \hat{T}_{kin} , \hat{T}_{int} , compared with \hat{T}_{kin} when $C_v = 0$.

Reduction: from 18 d.d.l. to 3, time step $\Delta t = 10^{-14}$ s !

Reduced dynamics: the reactive case

The progress variable

- One more parameter per particle: **progress variable** λ_i , describing the **progress along the free energy profile** associated with the decomposition process
- The interaction potential depends on the reaction rate. For instance, in the Lennard-Jones case,

$$V_{ij}(r_{ij}, \lambda_i, \lambda_j) = 4E_{ij} \left(\left(\frac{a_{ij}}{r_{ij}} \right)^{12} - \left(\frac{a_{ij}}{r_{ij}} \right)^6 \right),$$

with $E_{ij} = E \sqrt{(1 + k_E \lambda_i)(1 + k_E \lambda_j)}$, $a_{ij} = a (1 + k_a(\lambda_i + \lambda_j)/2)$.

- Reversible kinetics $AB \rightleftharpoons A_2 + B_2$, **depending on the temperature**

$$\frac{d\lambda_i}{dt} = \sum_{i \neq j} \omega(r_{ij}) [K_1(T_{ij}^{\text{int}})(1 - \lambda_j)(1 - \lambda_i) - K_2(T_{ij}^{\text{int}})\lambda_j \lambda_i]$$

- For instance, arrhénius form $K_i(T) = Z_i e^{-E_i/k_B T}$.

Treating the exothermicity

- Exothermicity of the reaction $\Delta E_{\text{exthm}} (= E_2 - E_1)$.
- Seek a dynamics such that $dH_{\text{tot}}(q, p, \epsilon, \lambda) = 0$ with

$$dH_{\text{tot}}(q, p, \epsilon, \lambda) = d \left[\sum_{1 \leq i < j \leq N} V(r_{ij}, \lambda_i, \lambda_j) + \sum_{i=1}^N \frac{p_i^2}{2m_i} + \epsilon_i + (1 - \lambda_i) \Delta E_{\text{exthm}} \right].$$

- Additional assumption: during the elementary step corresponding to exothermicity, the total energy of a given mesoparticle does not change:

$$d \left[\frac{1}{2} \sum_{i \neq j} V(r_{ij}, \lambda_i, \lambda_j) \right] + d \left(\frac{p_i^2}{2m_i} \right) + d\epsilon_i - \Delta E_{\text{exthm}} d\lambda_i = 0.$$

- Evolutions of momenta and internal energies balancing the variations in the total energy due to the variations of λ (exothermicity, changes in the potential energies) \rightarrow processes Z_i^p, Z_i^ϵ .

Treating the exothermicity (2)

- Distribution between internal energies and kinetic energies following some predetermined ratio $0 < c < 1$.
- For the internal energies (fix r , vary λ)

$$d\epsilon_i = -c \left(d \left[\frac{1}{2} \sum_{i \neq j} V(r_{ij}, \lambda_i, \lambda_j) \right] - \Delta E_{\text{exthm}} d\lambda_i \right).$$

- For the momenta, we consider a process Z_i^p such that $dp_i = dZ_i^p$ with

$$d \left(\frac{p_i^2}{2m} \right) = -(1 - c) \left(d \left[\frac{1}{2} \sum_{i \neq j} V(r_{ij}, \lambda_i, \lambda_j) \right] - \Delta E_{\text{exthm}} d\lambda_i \right).$$

In practice (2D case), for a variation δE_i^n due to the variations of $\{\lambda_i^n\}$,


$$p_i^{n+1} = p_i^n + \alpha^n (\cos \theta^n, \sin \theta^n), \quad \frac{(p_i^{n+1})^2}{2m_i} = \frac{(\tilde{p}_i^n)^2}{2m_i} + (1 - c) \delta E_i^n.$$

- Finally, the **reactive** DPDE dynamics is

$$\begin{aligned}dq_i &= \frac{p_i}{m_i} dt, \\dp_i &= \sum_{j, j \neq i} -\nabla_{q_i} V(r_{ij}, \lambda_i, \lambda_j) dt - \gamma_{ij} \chi^2(r_{ij}) v_{ij} dt + \sigma \chi(r_{ij}) dW_{ij} + dZ_i^p, \\d\epsilon_i &= \frac{1}{2} \sum_{j, j \neq i} \left(\chi^2(r_{ij}) \gamma_{ij} v_{ij}^2 - \frac{d\sigma^2}{2} \left(\frac{1}{m_i} + \frac{1}{m_j} \right) \chi^2(r_{ij}) \right) dt \\&\quad - \sigma \chi(r_{ij}) v_{ij} \cdot dW_{ij} + dZ_i^\epsilon, \\d\lambda_i &= \sum_{j \neq i} \omega_r(r_{ij}) [K_1(T_{ij})(1 - \lambda_i)(1 - \lambda_j) + K_2(T_{ij})\lambda_i \lambda_j] dt,\end{aligned}$$

- Numerical implementation: splitting of the dynamics as (inert) + (reaction)
- Integration of the reaction: update first λ_i , compute then the exothermicity (variations in the potential and liberated chemical energy), compute finally the new internal energies and velocities.

- Parameters inspired by the nitromethane example (replace CH_3NO_2 by a mesoparticle in a space of 2 dimensions).
- Classification of the parameters in five main categories
 - (Material parameters) molar mass $m = 80$ g/mol, Lennard-Jones potential with $E_{\text{LJ}} = 3 \times 10^{-21}$ J (melting 220 K), $a = 5$ Å, cut-off radius $r_{\text{cut}} = 15$ Å for the computation of forces. Changes of the material use $k_E = 0$ and $k_a = 0.2$ (pure expansion).
 - (Parameters of the inert dynamics) Microscopic state law is $\epsilon = C_v T$ with $C_v = 10 k_B$ (i.e., 20 d.o.f). Friction is $\gamma = 10^{-15}$ kg/s, dissipation weighting function $\chi(r) = (1 - r/r_c)$, with $r_c = r_{\text{cut}}$.
 - (Chemical kinetics) Prefactors $Z_1 = Z_2 = 10^{17}$ s⁻¹, activation energies $E_1/k_B = 15000$ K, exothermicity $\Delta E_{\text{exthm}} = 6.25$ eV. Weighting function $\omega(r) = \chi(r)$;
 - (Exothermicity) distribution fraction $c = 0.5$.
 - (Initial conditions) density $\rho = 1.06$ g/cm³, temperature $\bar{T} = 300$ K.



Velocity profiles in the material at different times (lower curve (red):
 $t = 1.2 \times 10^{-10}$ s; middle curve (black): $t = 1.6 \times 10^{-10}$ s; upper curve (blue):
 $t = 2 \times 10^{-10}$ s). Time-step $\Delta t = 2 \times 10^{-15}$ s.

- Systematic parametrization from small all atom simulations (potential, friction, microscopic state law $s = s(\epsilon)$, reaction constants, exothermicity)
- Dimensionality reduction allows to treat larger systems, for longer times
→ truly mesoscopic model? (polycrystalline materials)
- Hierarchy of models from discrete to continuum hydrodynamic equations (discretized with particle methods such as *Smoothed Particle Hydrodynamics*)
- References for this work:
 - G. STOLTZ, A reduced model for shock and detonation waves. I. The inert case, *Europhys. Lett.* **76**(5) (2006) 849-855.
 - J.-B. MAILLET, L. SOULARD AND G. STOLTZ, A reduced model for shock and detonation waves. II. The reactive case, accepted for publication in *Europhys. Lett.* (2007).